



ALGEBRAIC COMBINATORICS

Artur Rapp

Representation stability on the cohomology of complements of subspace arrangements

Volume 2, issue 4 (2019), p. 603-611.

[<http://alco.centre-mersenne.org/item/ALCO_2019__2_4_603_0>](http://alco.centre-mersenne.org/item/ALCO_2019__2_4_603_0)

© The journal and the authors, 2019.
Some rights reserved.



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is member of the
Centre Mersenne for Open Scientific Publishing
www.centre-mersenne.org



Representation stability on the cohomology of complements of subspace arrangements

Artur Rapp

ABSTRACT We study representation stability in the sense of Church and Farb of sequences of cohomology groups of complements of arrangements of linear subspaces in real and complex space as S_n -modules. We consider arrangements of linear subspaces defined by sets of diagonal equalities $x_i = x_j$ and invariant under the action of S_n which permutes the coordinates. We provide bounds for the point when stabilization occurs and an alternative proof of the fact that stabilization happens. The latter is a special case of very general stabilization results proved independently by Gadish and by Petersen; for the pure braid space the result is part of the work of Church and Farb. For the latter space, better stabilization bounds were obtained by Hersh and Reiner.

1. INTRODUCTION

In this paper, we consider arrangements of diagonal subspaces of \mathbb{R}^{dn} for natural numbers d and n . Let π be a set partition of $\{1, \dots, n\}$. Let W_π^d be the linear subspace of n -tuples (w_1, \dots, w_n) of points in \mathbb{R}^d such that $w_i = w_j$ whenever i and j are in the same block of π . For an integer partition λ we denote by \mathcal{A}_λ^d the arrangement of all subspaces W_π^d such that π is of type λ . More generally, set $\mathcal{A}_\Lambda^d = \cup_{\lambda \in \Lambda} \mathcal{A}_\lambda^d$ for every finite set Λ of integer partitions of n . The complement $\mathcal{M}_\Lambda^d = \mathbb{R}^{dn} \setminus \cup_{W \in \mathcal{A}_\Lambda^d} W$ is a real manifold. If $\Lambda = \{\lambda\}$, we write \mathcal{M}_λ^d for \mathcal{M}_Λ^d . All representations, homology groups and cohomology groups in this paper are taken with coefficients in \mathbb{C} . The action of the symmetric group S_n on n -tuples of points in \mathbb{R}^d by permuting the coordinates induces an S_n -representation on the reduced singular cohomology $\tilde{H}^i(\mathcal{M}_\Lambda^d)$. Formulas for these S_n -representations were determined by Sundaram and Welker in [8]. We look into representation stability in the sense of Church and Farb (see [2]) of these modules. Our main purpose is to prove that the sequence of modules stabilizes, and to obtain stabilization bounds. This is the content of Theorem 1.1. The fact that this sequence stabilizes is a special case of results of Gadish ([3, Theorem A]) and Petersen ([6, Theorem 4.15]). The case $\Lambda = \{(2, 1^{n-2})\}$ was proved by Church ([1, Theorem 1]) and for this case Hersh and Reiner provided better stabilization bounds ([4, Theorem 1.1]). For an integer partition λ we write $l(\lambda)$ for its length i.e. its number

Manuscript received 21st December 2017, revised 15th June 2018 and 4th January 2019, accepted 16th January 2019.

KEYWORDS. representation stability, subspace arrangement, symmetric functions.

ACKNOWLEDGEMENTS. I thank Nir Gadish and Dan Petersen for helpful comments and the anonymous referees for helpful comments and suggestions.

of parts. As in [4, Definition 2.5] let $\text{rank}(\lambda) := |\lambda| - l(\lambda)$ be the rank of λ . Note that set partitions of type λ have $\text{rank}(\lambda)$ as their poset rank in the partition lattice.

THEOREM 1.1. *Let Λ be a nonempty finite set of integer partitions of the number n_0 not containing (1^{n_0}) . For every $n \geq n_0$ let $\Lambda^{(n)}$ be the set of all integer partitions of n obtained from integer partitions in Λ by adding $n - n_0$ parts of size 1. Let $\text{rank}(\Lambda) = \min\{\text{rank}(\lambda) \mid \lambda \in \Lambda\}$. For every i and $d \geq 2$ the sequence $\{\tilde{H}^i(\mathcal{M}_{\Lambda^{(n)}}^d)\}_n$ stabilizes at $4(i + 1 - \text{rank}(\Lambda))/(d - 1)$.*

In Section 2, we provide the definition of representation stability and prove Theorem 1.1. In Section 3, we consider the special case $\mathcal{M}_{(k, 1^{n-k})}^d$ for $k \geq d + 1$. We prove that stability in this case starts earlier than in the bound given in Theorem 1.1.

2. PROOF OF THE MAIN THEOREM

An integer partition λ of a natural number n is a finite sequence $(\lambda_1, \lambda_2, \dots)$ with $\lambda_1 \geq \lambda_2 \geq \dots$ and $\sum_{i \geq 1} \lambda_i = n$. We sometimes denote λ by $(1^{m_1(\lambda)}, \dots, n^{m_n(\lambda)})$ where $m_i(\lambda)$ is the number of occurrences of the number i in λ for $1 \leq i \leq n$. Given an integer partition $\lambda = (\lambda_1, \lambda_2, \dots)$ of n we write $\lambda + \square$ to denote $(\lambda_1 + 1, \lambda_2, \lambda_3, \dots)$. Let V be an S_n -representation and $\sum_{\lambda \vdash n} a_\lambda S^\lambda$, $a_\lambda \in \mathbb{C}$, its decomposition into irreducible S_n -representations S^λ . Then we write

$$V + \square := \sum_{\lambda \vdash n} a_\lambda S^{\lambda + \square}.$$

We use the same notation when S_n -representations are replaced by symmetric functions (see [5] for background on symmetric functions). As in [5] we write s_λ for the Schur function indexed by the integer partition λ and ch for the Frobenius characteristic. Henceforth we simply refer to this as characteristic. The above equation becomes

$$\text{ch}(V) + \square := \sum_{\lambda \vdash n} a_\lambda s_{\lambda + \square}.$$

Now let $n_0 \in \mathbb{N}$. Let $\{V_n\}_{n \geq n_0}$ be a sequence of S_{n+n_0} -representations or a sequence of characteristics of S_{n+n_0} -representations. We say that this sequence stabilizes at $m \geq n_0$, if

$$V_n = V_{n-1} + \square$$

for all $n > m$. We say that the sequence stabilizes sharply at m , if m is the smallest integer such that

$$V_n = V_{n-1} + \square$$

for all $n > m$. The following lemma is a generalization of [4, Lemma 2.2].

For integer partitions ν, λ and μ with $\mu \subseteq \nu$, we write $LR_{\mu, \lambda}^\nu$ for the set of all Littlewood-Richardson tableaux of shape ν/μ and weight λ . A Littlewood-Richardson tableau T of shape ν/μ and weight λ is a semistandard skew tableau of shape ν/μ whose boxes are labeled with λ_1 1's, λ_2 2's etc. and concatenating the reversed rows of T from top to bottom yields a word w with the property: In every initial part of w the integer i occurs at least as often as $i + 1$ for every $i \geq 1$.

LEMMA 2.1. *Let λ and α be integer partitions. For every $n \geq \alpha_1$ we consider the integer partition $(n, \alpha) = (n, \alpha_1, \alpha_2, \dots)$. The sequence $\{s_{(n, \alpha)} s_\lambda\}_n$ stabilizes sharply at $\lambda_1 + \alpha_1$. In other words*


$$s_{(n, \alpha)} s_\lambda = s_{(n-1, \alpha)} s_\lambda + \square$$

if and only if $n > \lambda_1 + \alpha_1$.

Proof. Suppose $n > \lambda_1 + \alpha_1$. Let ν be an integer partition of $n + |\lambda| + |\alpha|$ with $(n, \alpha) \subseteq \nu$. By the Littlewood-Richardson rule (see [5]) the multiplicity of s_ν in $s_{(n,\alpha)}s_\lambda$ is $\#LR_{(n,\alpha),\lambda}^\nu$. Let ν' be the integer partition of $n - 1 + |\lambda| + |\alpha|$ obtained from ν by replacing ν_1 by $\nu_1 - 1$. We define the map

$$\phi : LR_{(n,\alpha),\lambda}^\nu \rightarrow LR_{(n-1,\alpha),\lambda}^{\nu'}$$

by the following procedure: Remove the first empty box in the first row of the tableau and then move all other boxes of the first row one place to the left. The two steps are illustrated below with $n = 5$, $\alpha = (1, 1)$, $\lambda = (3, 1)$ and $\nu = (6, 4, 1)$:



We want to show that the resulting tableau is indeed a Littlewood-Richardson tableau so that ϕ is well defined. The only condition that has to be checked is that, in the first two rows, we have no two 1's lying in the same column. But this follows from the inequality $\nu_1 \geq n$, since $n > \lambda_1 + \alpha_1$ implies that n is larger than the number α_1 of empty boxes in the second row plus the number of 1's in the second row. Note that ϕ has an inverse map: Given a tableau in $LR_{(n-1,\alpha),\lambda}^{\nu'}$ we move the first row one place to the right and put an empty box in the gap. So ϕ is bijective and $\#LR_{(n,\alpha),\lambda}^\nu = \#LR_{(n-1,\alpha),\lambda}^{\nu'}$. This shows that $\{s_{(n,\alpha)}s_\lambda\}_n$ stabilizes at $\lambda_1 + \alpha_1$ or sooner. Now let $n = \lambda_1 + \alpha_1$ and $\nu = (n, n, \lambda_2 + \alpha_2, \lambda_3 + \alpha_3, \dots)$. There is a Littlewood-Richardson tableau of shape $\nu/(n, \alpha)$ and weight λ : We look at the Ferrers diagram of ν and put λ_1 1's at the end of the second row, λ_2 2's at the end of the third row and so on. It follows that we have a Schur function s_ν with $\nu_1 = \nu_2$ and multiplicity greater than or equal to 1 in the decomposition of $s_{(n,\alpha)}s_\lambda$. This shows that $s_{(n,\alpha)}s_\lambda$ cannot equal $f + \square$ for any symmetric function f , completing the proof of sharpness. \square

Though the special case of Lemma 2.1 where $\alpha = ()$ ([4, Lemma 2.2]) suffices to prove our main results, Theorems 1.1 and 3.3, the general case might also be of interest as we show in Section 4. For a finite arrangement \mathcal{A} of linear subspaces of \mathbb{R}^{dn} , the intersection lattice $L_{\mathcal{A}}$ is the set of intersections of arbitrarily many elements of \mathcal{A} ordered by reverse inclusion. The least element $\hat{0}$ is \mathbb{R}^{dn} , the empty intersection, and the greatest element $\hat{1}$ is the intersection of all elements of \mathcal{A} . For a subset T of $L_{\mathcal{A}}$ the join sublattice of $L_{\mathcal{A}}$ generated by T consists of all intersections of arbitrarily many elements of T also ordered by reverse inclusion. If \mathcal{A} is the arrangement of diagonal subspaces given by equations of the form $w_i = w_j$ for $1 \leq i < j \leq n$, $w = (w_1, \dots, w_n) \in (\mathbb{R}^d)^n$, the intersection lattice $L_{\mathcal{A}}$ is isomorphic to the lattice Π_n . For a set partition $\pi \in \Pi_n$, we also write π for the corresponding subspace of \mathbb{R}^{dn} . If $\pi \in \Pi_n$ is a set partition into the subsets B_1, \dots, B_l of $\{1, \dots, n\}$, we write $\pi = B_1 | \dots | B_l$. In this notation, we have $\hat{0} = \{1\} | \{2\} | \dots | \{n\}$. The set partition $\pi = B_1 | \dots | B_l$ is said to be finer than $\pi' = C_1 | \dots | C_m$, if for every $1 \leq i \leq l$ there is a $1 \leq j \leq m$ such that $B_i \subseteq C_j$. We may reorder the sets B_1, \dots, B_l such that $\#B_1 \geq \dots \geq \#B_l$. The integer partition $(\#B_1, \dots, \#B_l)$ is then called the type of π . If Λ is a set of integer partitions of n , then Π_Λ is the join sublattice of Π_n generated by all set partitions of type λ for all $\lambda \in \Lambda$.

Proof of Theorem 1.1. By [8, Theorem 2.5(ii)] we have

$$\tilde{H}^i(\mathcal{M}_{\Lambda(n)}^d) = \bigoplus_{\pi \in (\Pi_{\Lambda(n)}^{>0})/S_n} \text{Ind}_{(S_n)_\pi}^{S_n} (\tilde{H}_{\text{codim}(\pi)-i-2}(\hat{0}, \pi) \otimes \tilde{H}_{\text{codim}(\pi)-1}(S^{dn-1} \cap \pi^\perp)).$$

$(\Pi_{\Lambda^{(n)}}^{\hat{0}})/S_n$ is a set of representatives of the action of S_n on $\Pi_{\Lambda^{(n)}}$ excluding $\hat{0}$. $(S_n)_\pi$ is the stabilizer subgroup of π . $\tilde{H}_j([\hat{0}, \pi])$ is the reduced simplicial homology on the order complex $\Delta([\hat{0}, \pi])$ in degree $j \geq -1$. $\text{codim}(\pi)$ is the codimension of π as a real subspace of \mathbb{R}^{dn} and S^{dn-1} is the $(dn - 1)$ -dimensional sphere. If π is of type $\mu = (1^{m_1(\mu)}, 2^{m_2(\mu)}, \dots) \vdash n$, then its stabilizer $(S_n)_\pi$ is the product of wreath products $\prod_j S_{m_j(\mu)}[S_j]$ and $\text{codim}(\pi) = d(n - l(\mu)) = d \cdot \text{rank}(\mu)$. The length of a chain in Π_n from $\hat{0}$ to π is less than or equal to $\sum_{j=1}^{l(\mu)} (\mu_j - 1) = n - l(\mu) = \text{rank}(\mu)$. Since the atoms in Π_Λ are of shape λ for $\lambda \in \Lambda$, the length of a chain in Π_Λ from $\hat{0}$ to π is less than or equal to $\text{rank}(\mu) - \text{rank}(\Lambda) + 1$ and contributes to homology in degree less than or equal to $\text{rank}(\mu) - \text{rank}(\Lambda) - 1$. It follows that if the homology $\tilde{H}_{\text{codim}(\pi)-i-2}([\hat{0}, \pi])$ is not zero, then

$$-1 \leq d \cdot \text{rank}(\mu) - i - 2 \leq \text{rank}(\mu) - \text{rank}(\Lambda) - 1$$

and then

$$(i + 1)/d \leq \text{rank}(\mu) \leq (i + 1 - \text{rank}(\Lambda))/(d - 1).$$

Let $\tilde{\mu}$ be the integer partition obtained from μ by removing the parts of size 1. The rank of μ and the rank of $\tilde{\mu}$ are the same. From [4, Proposition 2.8], we have $\text{rank}(\tilde{\mu}) + 1 \leq |\tilde{\mu}| \leq 2 \cdot \text{rank}(\tilde{\mu})$. This yields

$$1 + (i + 1)/d \leq |\tilde{\mu}| \leq 2(i + 1 - \text{rank}(\Lambda))/(d - 1).$$

The subgroup $S_{m_1(\mu)}[S_1] \cong S_{m_1(\mu)}$ acts trivially on $\tilde{H}_{\text{codim}(\pi)-i-2}([\hat{0}, \pi])$. The coordinates of vectors in the space π^\perp which correspond to the singletons of π are zero. It follows that the above copy of $S_{m_1(\mu)}$ acts trivially on $\tilde{H}_{\text{codim}(\pi)-1}(S^{dn-1} \cap \pi^\perp)$. Let $S^{(m_1(\mu))}$ be the trivial $S_{m_1(\mu)}$ -module. We get the following isomorphism of $\prod_{j \geq 1} S_{m_j(\mu)}[S_j]$ -modules:

$$\begin{aligned} \tilde{H}_{\text{codim}(\pi)-i-2}([\hat{0}, \pi]) \otimes \tilde{H}_{\text{codim}(\pi)-1}(S^{dn-1} \cap \pi^\perp) \\ \cong S^{(m_1(\mu))} \otimes (\tilde{H}_{\text{codim}(\pi)-i-2}([\hat{0}, \pi]) \otimes \tilde{H}_{\text{codim}(\pi)-1}(S^{dn-1} \cap \pi^\perp)). \end{aligned}$$

We consider the interval $[\hat{0}, \pi]$ in $\Pi_{\Lambda^{(n)}}$. The atoms in $[\hat{0}, \pi]$ have at least $n - n_0$ singletons. If we delete $\min\{n - |\tilde{\mu}|, n - n_0\}$ many singletons from π , after renumbering we can view $[0, \pi]$ as an interval in $\Pi_{\Lambda^{(max\{|\tilde{\mu}|, n_0\})}}$. We may also ignore the coordinates of vectors in π^\perp which correspond to the singletons of π . We have $\text{codim}(\pi) = d \cdot \text{rank}(\tilde{\mu})$. It follows that the $\prod_{j \geq 2} S_{m_j(\mu)}[S_j]$ -module

$$\tilde{H}_{\text{codim}(\pi)-i-2}([\hat{0}, \pi]) \otimes \tilde{H}_{\text{codim}(\pi)-1}(S^{dn-1} \cap \pi^\perp)$$

does not depend on n and we write $V_{\tilde{\mu}}$ for it. Using the transitivity of induction on $\prod_{j \geq 1} S_{m_j(\mu)}[S_j] \leq S_{m_1(\mu)} \times S_{n-m_1(\mu)} \leq S_n$ we get:

$$\begin{aligned} \text{Ind}_{\prod_{j \geq 1} S_{m_j(\mu)}[S_j]}^{S_n} (S^{m_1(\mu)} \otimes V_{\tilde{\mu}}) &= \text{Ind}_{S_{m_1(\mu)} \times S_{n-m_1(\mu)}}^{S_n} \left(S^{m_1(\mu)} \otimes \text{Ind}_{\prod_{j \geq 2} S_{m_j(\mu)}[S_j]}^{S_{n-m_1(\mu)}} (V_{\tilde{\mu}}) \right) \\ &= \text{Ind}_{S_{n-|\tilde{\mu}|} \times S_{|\tilde{\mu}|}}^{S_n} \left(S^{n-|\tilde{\mu}|} \otimes \text{Ind}_{\prod_{j \geq 2} S_{m_j(\tilde{\mu})}[S_j]}^{S_{|\tilde{\mu}|}} (V_{\tilde{\mu}}) \right). \end{aligned}$$

Let

$$f_{\tilde{\mu}} := \text{ch} \left(\text{Ind}_{\prod_{j \geq 2} S_{m_j(\tilde{\mu})}[S_j]}^{S_{|\tilde{\mu}|}} (V_{\tilde{\mu}}) \right).$$

We have

$$\text{ch} \left(\text{Ind}_{S_{n-|\tilde{\mu}|} \times S_{|\tilde{\mu}|}}^{S_n} \left(S^{n-|\tilde{\mu}|} \otimes \text{Ind}_{\prod_{j \geq 2} S_{m_j(\tilde{\mu})}[S_j]}^{S_{|\tilde{\mu}|}} (V_{\tilde{\mu}}) \right) \right) = h_{n-|\tilde{\mu}|} f_{\tilde{\mu}}$$

where $h_{n-|\tilde{\mu}|} = s_{(n-|\tilde{\mu}|)}$. It follows that the characteristic of $\tilde{H}^i(\mathcal{M}_{\Lambda(n)}^d)$ is

$$\sum_{\tilde{\mu}} h_{n-|\tilde{\mu}|} f_{\tilde{\mu}}$$

where the summation runs over all integer partitions $\tilde{\mu}$ with no parts 1, and $1 + (i + 1)/d \leq |\tilde{\mu}| \leq 2(i + 1 - \text{rank}(\Lambda))/(d - 1)$.

From Lemma 2.1, it follows that the sequence stabilizes at a number larger than $2|\tilde{\mu}|$ for every $\tilde{\mu}$ occurring in the sum. This is fulfilled at $4(i + 1 - \text{rank}(\Lambda))/(d - 1)$. \square

3. IMPROVED STABILITY BOUNDS FOR k -EQUAL ARRANGEMENTS

We consider the sequence $\{\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^d)\}_n$ for $k \geq 2$. Theorem 1.1 states that stabilization occurs at $4(i + 2 - k)/(d - 1)$. First we have a closer look at the special case $k = 2$. In this case, stabilization occurs at $4i/(d - 1)$. Compare this to the known results in the literature which focus on the case $k = 2$: By [1, Theorem 1] we have stabilization at $2i$ for $d \geq 3$ and stabilization at $4i$ for $d = 2$. By [4, Theorem 1.1] we have the following for $i \geq 1$. The sequence is zero from the beginning, if $d - 1$ does not divide i . Otherwise it stabilizes sharply at $3i/(d - 1)$ for odd $d \geq 3$ and it stabilizes sharply at $3i/(d - 1) + 1$ for even $d \geq 2$.

Now we consider $\{\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^d)\}_n$ for general $k \geq 2$. The stability of this sequence was also considered by Gadish ([3, Example 6.11]) as an example of his general results. We want to determine smaller upper bounds than the ones given in Theorem 1.1 where stabilization occurs for $k \geq d + 1$. Let $h_n = s_{(n)}$ be the complete homogeneous symmetric function, $e_n = s_{(1^n)}$ the elementary symmetric function and ω the involutive ring homomorphism of the ring of symmetric functions with $\omega(h_n) = e_n$. We write π_n for the characteristic of $\tilde{H}^{n-3}(\Delta(\Pi_n))$ and $l_n = \omega(\pi_n)$. For symmetric functions f and g we write $f[g]$ for the plethysm of these two functions.

THEOREM 3.1. [8, Theorem 4.4(iii)] *Let $d \geq 2$, $k \geq 2$, $i \geq 0$ and $n \geq 1$. Let $U_k := \sum_{j \geq k} s_{(j-k+1, 1^{k-1})}$. For every $r, t \geq 1$ and $q \geq 0$ such that $i = (d-1)(n-r-q) + t(k-2)$ let $\psi_{n,q,r,t}$ be*

$$\begin{cases} \omega\left(\omega^k\left(e_r\left[\sum_{j \geq 1} l_j\right]\right)\right) |_{\text{deg } t[U_k]} |_{\text{deg } n-q} h_q & \text{if } d \text{ is even} \\ \left(\left(h_r\left[\sum_{j \geq 1} l_j\right]\right) |_{\text{deg } t[U_k]}\right) |_{\text{deg } n-q} h_q & \text{if } d \text{ is odd and } k \text{ is even} \\ \left(\left((-1)^t h_r\left[\sum_{j \geq 1} (-1)^j \pi_j\right]\right) |_{\text{deg } t[U_k]}\right) |_{\text{deg } n-q} h_q & \text{if } d \text{ and } k \text{ are odd.} \end{cases}$$

Then the characteristic of the S_n -representation on $\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^d)$ is

$$\sum_{\substack{r,t \geq 1, q \geq 0: \\ i = (d-1)(n-r-q) + t(k-2)}} \psi_{n,q,r,t}.$$

LEMMA 3.2. *Let $d \geq 2$, $k \geq d + 1$, $i \geq 0$ and $n \geq 1$. Let $r, t \geq 1$ and $q \geq 0$ be such that $i = (d - 1)(n - r - q) + t(k - 2)$. Let $U_k := \sum_{j \geq k} s_{(j-k+1, 1^{k-1})}$ and $\psi_{n,q,r,t}$ be*

$$\begin{cases} \omega\left(\omega^k\left(e_r\left[\sum_{j \geq 1} l_j\right]\right)\right) |_{\text{deg } t[U_k]} |_{\text{deg } n-q} h_q & \text{if } d \text{ is even} \\ \left(\left(h_r\left[\sum_{j \geq 1} l_j\right]\right) |_{\text{deg } t[U_k]}\right) |_{\text{deg } n-q} h_q & \text{if } d \text{ is odd and } k \text{ is even} \\ \left(\left((-1)^t h_r\left[\sum_{j \geq 1} (-1)^j \pi_j\right]\right) |_{\text{deg } t[U_k]}\right) |_{\text{deg } n-q} h_q & \text{if } d \text{ and } k \text{ are odd.} \end{cases}$$

Then

- (i) $\psi_{n,q,r,t} = \psi_{n-1,q-1,r,t} + \square$ if $q > n/2$ and $n \geq 2$.
- (ii) $\psi_{n,q,r,t} = \psi_{n-1,q-1,r,t} + \square$ if d is even, $q > tk$ and $n \geq 2$.

- (iii) $\psi_{n,q,r,t} = 0$ if $r > t$ or $t > n/k$.
- (iv) $\psi_{n,q,r,t} = 0$ if $q \leq n/2$ and $n > \frac{2i}{d-1}$.
- (v) $\psi_{n,q,r,t} = 0$ if $k \geq d + 2$, $q \leq tk$ and $n > \frac{ki}{k-d-1}$.

Proof. (i) We have $\psi_{n,q,r,t} = f_{n-q}h_q$ for a symmetric function f_{n-q} of degree $n - q$ and $h_q = s_{(q)}$. From Lemma 2.1, we get

$$\psi_{n,q,r,t} = f_{n-q}h_q = f_{n-q}h_{q-1} + \square = f_{(n-1)-(q-1)}h_{q-1} + \square = \psi_{n-1,q-1,r,t} + \square$$

if $q > n - q$ or equivalently $q > n/2$.

(ii) If d is even, then $\psi_{n,q,r,t} = \omega(f_t[U_k])|_{deg\ n-q}h_q$ for a symmetric function f_t of degree t . The first column of every Schur function in $U_k = \sum_{j \geq k} s_{(j-k+1, 1^{k-1})}$ has length k . From [4, Proposition 4.3 (d)] it follows that for every s_λ with $\lambda \vdash n - q$ occurring in the Schur function decomposition of $\omega(f_t[U_k])|_{deg\ n-q}$ the first row of λ has length less than or equal to tk . If $q > tk$, it follows from Lemma 2.1 that

$$\begin{aligned} \psi_{n,q,r,t} &= \omega(f_t[U_k])|_{deg\ n-q}h_q \\ &= \omega(f_t[U_k])|_{deg\ n-q}h_{q-1} + \square \\ &= \omega(f_t[U_k])|_{deg\ (n-1)-(q-1)}h_{q-1} + \square = \psi_{n-1,q-1,r,t} + \square. \end{aligned}$$

(iii) If $r > t$ the terms $e_r[\sum_{j \geq 1} l_j]$, $h_r[\sum_{j \geq 1} l_j]$ and $(-1)^t h_r[\sum_{j \geq 1} (-1)^j \pi_j]$ only have terms of degree greater than t . Then the whole term $\psi_{n,q,r,t}$ is zero. U_k only has terms of degree greater than or equal to k . Then $f_t[U_k]$ for a symmetric function f_t of degree t has only terms of degree greater than or equal to tk . If $t > n/k$ then $tk > n \geq n - q$ and again $\psi_{n,q,r,t}$ is zero.

(iv) Suppose $\psi_{n,q,r,t} \neq 0$. We have to show that $q > n/2$ or $n \leq \frac{2i}{d-1}$. Suppose $q \leq n/2$. From $\psi_{n,q,r,t} \neq 0$ and (ii) we get $r \leq t$. From $q \leq n/2$ and $i = (d - 1)(n - r - q) + t(k - 2)$ we get

$$\frac{i}{1-d} + n/2 + \frac{t(k-2)}{d-1} \leq \frac{i}{1-d} + n - q + \frac{t(k-2)}{d-1} = r.$$

Using $r \leq t$ we get

$$\frac{i}{1-d} + n/2 + \frac{t(k-2)}{d-1} \leq t$$

and simplifying yields

$$n/2 \leq \frac{i}{d-1} + \frac{t(d+1-k)}{d-1}.$$

Using $k \geq d + 1$ we get

$$n \leq \frac{2i}{d-1}.$$

(v) Let $k \geq d+2$. Suppose $\psi_{n,q,r,t} \neq 0$ and $q \leq tk$. We have to show that $n \leq \frac{ki}{k-d-1}$. From $q \leq tk$, $i = (d - 1)(n - r - q) + t(k - 2)$ and $r \leq t$ by (iii) we get

$$\frac{i}{1-d} + n - tk + \frac{t(k-2)}{d-1} \leq \frac{i}{1-d} + n - q + \frac{t(k-2)}{d-1} = r \leq t.$$

It follows that

$$\frac{i}{1-d} + n - tk + \frac{t(k-2)}{d-1} \leq t$$

and then

$$n \leq \frac{i}{d-1} + t \left(k + \frac{k-2}{1-d} + 1 \right).$$

From (iii) we know $t \leq n/k$. It follows that

$$n \leq \frac{i}{d-1} + \frac{n}{k} \left(k + \frac{k-2}{1-d} + 1 \right)$$

and then

$$n \binom{k-2}{d-1} \leq \frac{ki}{d-1}.$$

Using $k \geq d+2$ we get

$$n \leq \frac{\frac{ki}{d-1}}{\frac{k-2}{d-1} - 1} = \frac{ki}{k-d-1}. \quad \square$$

THEOREM 3.3. *Let $d \geq 2$, $k \geq d+1$ and $i \geq 0$. The sequence $\{\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^d)\}_n$ stabilizes at $\frac{2i}{d-1}$. If d is even and $k \geq d+2$, the sequence stabilizes at $\frac{ki}{k-d-1}$.*

Proof. From Theorem 3.1, we have that the characteristic of the S_n -representation on $\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^d)$ is

$$\sum_{\substack{r,t \geq 1, q \geq 0: \\ i=(d-1)(n-r-q)+t(k-2)}} \psi_{n,q,r,t}$$

where $\psi_{n,q,r,t}$ is as in the previous lemma. If $q > n/2$ then we get

$$\psi_{n,q,r,t} = \psi_{n-1,q-1,r,t} + \square$$

from Lemma 3.2(i). From Lemma 3.2(iv) we get $\psi_{n,q,r,t} = 0$ if $q \leq n/2$ and $n > \frac{2i}{d-1}$. Putting these facts together we get for $n > \frac{2i}{d-1}$:

$$\begin{aligned} \sum_{\substack{r,t \geq 1, q \geq 0: \\ i=(d-1)(n-r-q)+t(k-2)}} \psi_{n,q,r,t} &= \sum_{\substack{r,t \geq 1, q \geq 1: \\ i=(d-1)(n-r-q)+t(k-2)}} \psi_{n,q,r,t} \\ &= \sum_{\substack{r,t \geq 1, q \geq 1: \\ i=(d-1)(n-r-q)+t(k-2)}} \psi_{n-1,q-1,r,t} + \square \\ &= \sum_{\substack{r,t \geq 1, q \geq 0: \\ i=(d-1)(n-1-r-q)+t(k-2)}} \psi_{n-1,q,r,t} + \square. \end{aligned}$$

Now let d be even and $k \geq d+2$. If d is even and $q > tk$ we have

$$\psi_{n,q,r,t} = \psi_{n-1,q-1,r,t} + \square$$

from Lemma 3.2(ii) and $\psi_{n,q,r,t} = 0$ if $q \leq tk$ and $n > \frac{ki}{k-d-1}$ from Lemma 3.2(v). For $n > \frac{ki}{k-d-1}$ the same computation as above yields the stability property. \square

In Table 3.5, we give a list of sharp stability bounds for these representations.

QUESTION 3.4. *Is there an explicit formula for the sharp stability bound of $\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^d)$ for general k, d, i ?*

TABLE 3.5 (Sharp stability bounds for $\tilde{H}^i(\mathcal{M}_{(k,1^{n-k})}^2)$). *If k is fixed and i grows, the sequence of bounds appears to increase by 1 in most of the steps especially at the beginning and with large k . Later, there also appear steps with bound differences 2 or 3.*

$k = 3$:

i	3	4	5	6	7	8	9	10	11	12	13	14
bound	6	7	8	11	13	14	16	18	20	21	23	25

$k = 4$:

i	5	6	7	8	9	10	11	12	13	14	15	16
bound	8	9	10	11	12	15	17	18	19	20	22	24

$k = 5 :$

i	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$bound$	10	11	12	13	14	15	16	19	21	22	23	24	25	26

$k = 6 :$

i	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
$bound$	12	13	14	15	16	17	18	19	20	23	25	26	27	28	29

$k = 7 :$

i	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
$bound$	14	15	16	17	18	19	20	21	22	23	24	27	29	30	31	32	33

$k = 8 :$

i	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
$bound$	16	17	18	19	20	21	22	23	24	25	26	27	28	31	33	34	35

$k = 9 :$

i	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
$bound$	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	35	37

4. STABILITY IN THE HOMOLOGY OF k -EQUAL PARTITION LATTICES

We showed in Lemma 2.1 that for integer partitions α and λ the sequence $\{s_{(n,\alpha)}s_\lambda\}_n$ stabilizes at $\alpha_1 + \lambda_1$. In this section we give an application of this fact in a situation where α is not the empty partition. For every $2 \leq k \leq n$ we consider the lattice $\Pi_{(k,1^{n-k})}$ of set partitions all of whose block sizes are 1 or greater than or equal to k ordered by reverse refinement. We have $\Pi_{(2,1^{n-2})} = \Pi_n$. Note that $\Pi_{k,1^{n-k}}$ is the intersection lattice of the subspace arrangement with complement $\mathcal{M}_{(k,1^{n-k})}^d$. We recall the following result on the homology of the order complex of $\Pi_{k,1^{n-k}}$:

THEOREM 4.1 ([7, Corollary 3.6]).

- (i) Let $3 \leq k \leq n$ and $1 \leq t \leq \lfloor n/k \rfloor$. The characteristic of $\tilde{H}_{n-3-t(k-2)}(\Pi_{(k,1^{n-k})})$ tensored with the sign representation is given by the degree n term in

$$\omega^k(l_t) \left[\sum_{j \geq k} s_{(j-k+1,1^{k-1})} \right].$$

- (ii) Let $2 = k \leq n$. The characteristic of $\tilde{H}_{n-3}(\Pi_n)$ tensored with the sign representation is given by the degree n term in

$$\sum_{t=1}^{\lfloor n/2 \rfloor} l_t \left[\sum_{j \geq 2} s_{(j-1,1)} \right].$$

By [7, Lemma 3.2] the term $\omega^k(l_t) \left[\sum_{j \geq k} s_{(j-k+1,1^{k-1})} \right] |_{\deg n}$ decomposes into the sum

$$\sum_{\lambda} \phi_{k,t,n,\lambda}$$

where

$$\phi_{k,t,n,\lambda} = \omega^k(l_t) |_{\prod_{i \geq 1: m_i > 0} S_{m_i}} \left[\bigotimes_{j \geq k} s_{(j-k+1,1^{k-1})} \right]$$

and the sum runs over all partitions $\lambda = (\lambda_1, \dots, \lambda_t) = (n^{m_n}, \dots, k^{m_k})$ of n with t parts and all parts greater than or equal to k . Now we apply Lemma 2.1 to $\{\phi_{k,t,n,\lambda}\}_{\lambda_1}$ where $\lambda_2, \dots, \lambda_t, k$ and t are fixed and λ_1 and $n = n(\lambda_1) = \sum_{i \geq 1} \lambda_i$ grow.

PROPOSITION 4.2. *The sequence $\{\phi_{k,t,n,\lambda}\}_{\lambda_1}$ stabilizes at $\lambda_1 = k + \sum_{i \geq 2} \lambda_i$.*

Proof. If $\lambda_1 > \lambda_2$ then $m_{\lambda_1} = 1$ and the restriction $\omega^k(l_t)|_{\prod_{i \geq 1: m_i > 0} S_{m_i}}$ is the tensor product of the trivial S_1 -module and a $\prod_{i \geq 1: i \neq \lambda_1, m_i > 0} S_{m_i}$ -module. We get $\phi_{k,t,n,\lambda} = s_{(\lambda_1 - k + 1, 1^{k-1})} f$ for a symmetric function f of degree $\sum_{i \geq 2} \lambda_i$. It follows from Lemma 2.1 that $\{\phi_{k,t,n,\lambda}\}_{\lambda_1}$ stabilizes at $\lambda_1 = k + \sum_{i \geq 2} \lambda_i$. \square

REFERENCES

- [1] Thomas Church, *Homological stability for configuration spaces of manifolds*, Invent. Math. **188** (2012), no. 2, 465–504.
- [2] Thomas Church and Benson Farb, *Representation theory and homological stability*, Adv. Math. **245** (2013), 250–314.
- [3] Nir Gadish, *Representation stability for families of linear subspace arrangements*, Adv. Math. **322** (2017), 341–377.
- [4] Patricia Hersh and Victor Reiner, *Representation stability for cohomology of configuration spaces in \mathbb{R}^d* , Int. Math. Res. Not. IMRN **2017** (2017), no. 5, 1433–1486, With an appendix written jointly with Steven Sam.
- [5] Ian Grant Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications.
- [6] Dan Petersen, *A spectral sequence for stratified spaces and configuration spaces of points*, Geom. Topol. **21** (2017), no. 4, 2527–2555.
- [7] Sheila Sundaram and Michelle Wachs, *The homology representations of the k -equal partition lattice*, Trans. Amer. Math. Soc. **349** (1997), no. 3, 935–954.
- [8] Sheila Sundaram and Volkmar Welker, *Group actions on arrangements of linear subspaces and applications to configuration spaces*, Trans. Amer. Math. Soc. **349** (1997), no. 4, 1389–1420.

ARTUR RAPP, Philipps-Universität Marburg, Fachbereich Mathematik und Informatik, Hans-Meerweinstr. 6, 35032 Marburg, Germany
E-mail : rapp202@mathematik.uni-marburg.de