



ALGEBRAIC COMBINATORICS


Sebastian König

The decomposition of 0-Hecke modules associated to quasisymmetric Schur functions

Volume 2, issue 5 (2019), p. 735-751.

http://alco.centre-mersenne.org/item/ALCO_2019__2_5_735_0

© The journal and the authors, 2019.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is member of the
Centre Mersenne for Open Scientific Publishing
www.centre-mersenne.org



The decomposition of 0-Hecke modules associated to quasisymmetric Schur functions

Sebastian König

ABSTRACT Recently Tewari and van Willigenburg constructed modules of the 0-Hecke algebra that are mapped to the quasisymmetric Schur functions by the quasisymmetric characteristic. These modules have a natural decomposition into a direct sum of certain submodules. We show that the summands are indecomposable by determining their endomorphism rings.

1. INTRODUCTION

Since the 19th century mathematicians have been interested in the Schur functions s_λ and their various properties. For example, they form an orthonormal basis of Sym , the Hopf algebra of symmetric functions and they are the images of the irreducible complex characters of the symmetric groups under the characteristic map [13]. The symmetric functions are contained in the Hopf algebra $QSym$ of quasisymmetric functions defined by Gessel in 1984 [6]. An introduction to $QSym$ can be found in [7].

There is a representation theoretic interpretation of $QSym$ as well. The 0-Hecke algebra $H_n(0)$ over a field \mathbb{K} is a deformation of the group algebra $\mathbb{K}\mathfrak{S}_n$ of the symmetric group obtained by replacing the generators $(i, i+1)$ of \mathfrak{S}_n by projections π_i satisfying the same braid relations. Norton discovered a great deal of the representation theory of the 0-Hecke algebras in [12]. For a more combinatorial approach see [9].

Let $\mathcal{G}_0(H_n(0))$ denote the Grothendieck group of the finitely generated $H_n(0)$ -modules and $\mathcal{G} := \bigoplus_{n \geq 0} \mathcal{G}_0(H_n(0))$. Duchamp, Krob, Leclerc and Thibon established the connection to $QSym$ by defining an algebra isomorphism $Ch: \mathcal{G} \rightarrow QSym$ called quasisymmetric characteristic [5, 10].

As Sym is contained in $QSym$, one may ask whether there are quasisymmetric analogues of the Schur functions. One such analogue due to Haglund, Luoto, Mason and van Willigenburg is given by the quasisymmetric Schur functions \mathcal{S}_α [8]. They form a basis of $QSym$ and nicely refine the Schur functions via

$$s_\lambda = \sum_{\tilde{\alpha}=\lambda} \mathcal{S}_\alpha$$

where λ is a partition and the sum runs over all compositions α that rearrange λ [8] (see Section 2.2 for definitions). Bessenrodt, Luoto and van Willigenburg defined skew quasisymmetric Schur functions $\mathcal{S}_{\alpha//\beta}$ and proved a Littlewood–Richardson rule for expressing them in the basis of quasisymmetric Schur functions [3].

Manuscript received 11th December 2017, revised 6th September 2018, accepted 5th January 2019.

KEYWORDS. 0-Hecke algebra, composition tableau, quasisymmetric function, Schur function.

Another basis of $QSym$ sharing properties with the Schur functions is formed by the dual immaculate functions of Berg, Bergeron, Saliola, Serrano and Zabrocki [1]. Indecomposable 0-Hecke modules whose images under Ch are the dual immaculate functions were defined in [2].

Tewari and van Willigenburg constructed modules \mathcal{S}_α of the 0-Hecke algebra that are mapped to \mathcal{S}_α by Ch [14]. Each \mathcal{S}_α has a \mathbb{K} -basis parametrized by a set of tableaux. By using an equivalence relation, they divided this set into equivalence classes, obtained a submodule $\mathcal{S}_{\alpha,E}$ of \mathcal{S}_α for each such equivalence class E and decomposed \mathcal{S}_α as $\mathcal{S}_\alpha = \bigoplus_E \mathcal{S}_{\alpha,E}$. In the same way they defined and decomposed skew modules $\mathcal{S}_{\alpha//\beta}$ whose image under Ch is $\mathcal{S}_{\alpha//\beta}$.

This article is mainly concerned with the modules \mathcal{S}_α and $\mathcal{S}_{\alpha,E}$. In [14], for a special equivalence class E_α it was shown that $\mathcal{S}_{\alpha,E_\alpha}$ is indecomposable. Yet, the question of the indecomposability of the modules $\mathcal{S}_{\alpha,E}$ in general remained open. The goal of this paper is to answer this question. We show that for each $\mathcal{S}_{\alpha,E}$ the ring of $H_n(0)$ -endomorphisms is $\mathbb{K} \text{id}$ so that $\mathcal{S}_{\alpha,E}$ is indecomposable. As a consequence, $\mathcal{S}_\alpha = \bigoplus_E \mathcal{S}_{\alpha,E}$ is a decomposition into indecomposable submodules.

The structure of the paper is as follows. In Section 2 we present the combinatorial and algebraic background and then review the modules $\mathcal{S}_{\alpha//\beta}$ and $\mathcal{S}_{\alpha//\beta,E}$. Section 3 is devoted to a related $H_n(0)$ -operation on chains of a composition poset. From this we obtain an argument crucial for proving the main results in Section 4.

2. BACKGROUND

We set $\mathbb{N} := \{1, 2, \dots\}$ and always assume that $n \in \mathbb{N}$. For $a, b \in \mathbb{Z}$ we define the *discrete interval* $[a, b] := \{c \in \mathbb{Z} \mid a \leq c \leq b\}$ and may use the shorthand $[a] := [1, a]$. Throughout this paper \mathbb{K} denotes an arbitrary field. For a set X , $\text{span}_{\mathbb{K}} X$ is the \mathbb{K} -vector space with basis X .

2.1. SYMMETRIC GROUPS AND 0-HECKE ALGEBRAS. The *symmetric group* \mathfrak{S}_n is the group of all permutations of the set $[n]$. We proceed by reviewing \mathfrak{S}_n as a Coxeter group. More details can be found in [4].

As a Coxeter group \mathfrak{S}_n is generated by the adjacent transpositions $s_i := (i, i + 1)$ for $i = 1, \dots, n - 1$ which satisfy

$$\begin{aligned} s_i^2 &= 1, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, \\ s_i s_j &= s_j s_i \text{ if } |i - j| \geq 2. \end{aligned}$$

The latter two relations are called *braid relations*. Let $\sigma \in \mathfrak{S}_n$. We can write σ as a product $\sigma = s_{j_k} \cdots s_{j_1}$. If k is minimal among such expressions, $s_{j_k} \cdots s_{j_1}$ is a *reduced word* for σ and $\ell(\sigma) := k$ is the *length* of σ .

The *support* of σ is $\text{supp}(\sigma) = \{i \in [n - 1] \mid s_i \text{ appears in a reduced word of } \sigma\}$. One assertion of the *word property* of \mathfrak{S}_n [4, Theorem 3.3.1] is that a reduced word of σ can be transformed into any other reduced word of σ by applying a sequence of braid relations. Thus, for each reduced word of σ the set of indices occurring in it is $\text{supp}(\sigma)$.

Let $\sigma, \tau \in \mathfrak{S}_n$. The *left weak order* \leq_L is the partial order on \mathfrak{S}_n given by

$$\sigma \leq_L \tau \iff \begin{aligned} \tau &= s_{i_k} \cdots s_{i_1} \sigma, \\ \ell(s_{i_r} \cdots s_{i_1} \sigma) &= \ell(\sigma) + r \text{ for } r = 1, \dots, k. \end{aligned}$$

The following Proposition 2.1 gathers immediate consequences of the definition.

PROPOSITION 2.1. *Let $\sigma, \tau \in \mathfrak{S}_n$.*

- (1) *We have $\sigma \leq_L \tau$ if and only if $\ell(\tau\sigma^{-1}) = \ell(\tau) - \ell(\sigma)$.*

(2) If $\sigma \leq_L \tau$ then the reduced words for $\tau\sigma^{-1}$ are in bijection with saturated chains in the left weak order poset (\mathfrak{S}_n, \leq_L) from σ to τ via

$$s_{i_k} \cdots s_{i_1} \longleftrightarrow \sigma \leq_L s_{i_1}\sigma \leq_L s_{i_2}s_{i_1}\sigma \leq_L \cdots \leq_L s_{i_k} \cdots s_{i_1}\sigma = \tau.$$

(3) The poset (\mathfrak{S}_n, \leq_L) is graded by the length function.

THEOREM 2.2 ([4, Corollary 3.2.2]). Let $\sigma, \tau \in \mathfrak{S}_n$. The interval in left weak order $[\sigma, \tau] := \{\rho \in \mathfrak{S}_n \mid \sigma \leq_L \rho \leq_L \tau\}$ is a graded lattice with rank function $\rho \mapsto \ell(\rho\sigma^{-1})$.

Next, we define the 0-Hecke algebra $H_n(0)$. We use the presentation as in [14] and refer to [11, Ch. 1] for details.

DEFINITION 2.3. The 0-Hecke algebra $H_n(0)$ is the unital associative \mathbb{K} -algebra generated by the elements $\pi_1, \pi_2, \dots, \pi_{n-1}$ subject to relations

$$\begin{aligned} \pi_i^2 &= \pi_i, \\ \pi_i\pi_{i+1}\pi_i &= \pi_{i+1}\pi_i\pi_{i+1}, \\ \pi_i\pi_j &= \pi_j\pi_i \text{ if } |i - j| \geq 2. \end{aligned}$$

Note that the π_1, \dots, π_{n-1} are projections satisfying the same braid relations as the s_1, \dots, s_{n-1} . Another set of generators of $H_n(0)$ is given by elements $\bar{\pi}_i$ for $i = 1, \dots, n - 1$ such that $\bar{\pi}_i^2 = -\pi_i$ and the braid relations hold. We do not use them here but they appear in some of the references.

Let $\sigma \in \mathfrak{S}_n$. We define $\pi_\sigma := \pi_{j_k} \cdots \pi_{j_1}$ where $s_{j_k} \cdots s_{j_1}$ is a reduced word for σ . The word property ensures that this is well defined. Multiplication is given by

$$\pi_i\pi_\sigma = \begin{cases} \pi_{s_i\sigma} & \text{if } \ell(s_i\sigma) > \ell(\sigma) \\ \pi_\sigma & \text{if } \ell(s_i\sigma) < \ell(\sigma) \end{cases}$$


for $i = 1, \dots, n - 1$. As a consequence, $\{\pi_\tau \mid \tau \in \mathfrak{S}_n\}$ spans $H_n(0)$ over \mathbb{K} . One can also show that it is a \mathbb{K} -basis of $H_n(0)$.

2.2. COMPOSITIONS AND COMPOSITION TABLEAUX. A *composition* $\alpha = (\alpha_1, \dots, \alpha_l)$ is a finite sequence of positive integers. The *length* and the *size* of α are given by $\ell(\alpha) := l$ and $|\alpha| := \sum_{i=1}^l \alpha_i$, respectively. The α_i 's are called *parts* of α . If α has size n , α is called *composition of n* and we write $\alpha \vDash n$. A *partition* is a composition whose parts are weakly decreasing. We write $\lambda \vdash n$ if λ is a partition of size n . For a composition α we denote the partition obtained by sorting the parts of α in decreasing order by $\tilde{\alpha}$. The *empty composition* \emptyset is the unique composition of length and size 0.

EXAMPLE 2.4. For $\alpha = (1, 4, 3) \vDash 8$, we have $\tilde{\alpha} = (4, 3, 1) \vdash 8$.

A *cell* (i, j) is an element of $\mathbb{N} \times \mathbb{N}$. A finite set of cells is called *diagram*. Diagrams are visualized in English notation. That is, for each cell (i, j) of a diagram we draw a box at position (i, j) in matrix coordinates. The *diagram* of $\alpha \vDash n$ is the set $\{(i, j) \in \mathbb{N} \times \mathbb{N} \mid i \leq \ell(\alpha), j \leq \alpha_i\}$. So, we display the diagram of α by putting α_i boxes in row i where the top row has index 1. We often identify α with its diagram.

EXAMPLE 2.5.




Next, we will consider standard composition tableaux and a related poset of compositions which arose in [3].

DEFINITION 2.6. The composition poset \mathcal{L}_c is the set of all compositions together with the partial order \leq_c given as the transitive closure of the following covering relation. For compositions α and $\beta = (\beta_1, \dots, \beta_l)$

$$\beta <_c \alpha \iff \begin{aligned} &\alpha = (1, \beta_1, \dots, \beta_l) \text{ or} \\ &\alpha = (\beta_1, \dots, \beta_k + 1, \dots, \beta_l) \text{ and } \beta_i \neq \beta_k \text{ for all } i < k. \end{aligned}$$

In other words, β is covered by α in \mathcal{L}_c if and only if the diagram of α can be obtained from the diagram of β by adding a box as the new first row or appending a box to a row which is the topmost row of its length in β .


EXAMPLE 2.7.



Let α and β be two compositions such that $\beta \leq_c \alpha$. In this situation we always assume that the diagram of β is moved to the bottom of the diagram of α , and we define the skew composition diagram (or skew shape) $\alpha // \beta$ to consist of all cells of α which are not contained in β .

Moreover, we define $\text{osh}(\alpha // \beta) = \alpha$ and $\text{ish}(\alpha // \beta) = \beta$ as the outer and the inner shape of $\alpha // \beta$, respectively. The size of a skew shape is $|\alpha // \beta| := |\alpha| - |\beta|$. We call $\alpha // \beta$ straight if $\beta = \emptyset$. In this case the skew composition diagram $\alpha // \beta$ is nothing but the ordinary composition diagram α .

EXAMPLE 2.8. In the following the cells of the inner shape are gray.




Note that $\beta \leq_c \alpha$ implies $\beta_{\ell(\beta)-i} \leq \alpha_{\ell(\alpha)-i}$ for $i = 0, \dots, \ell(\beta) - 1$. One could define skew shapes for all pairs of compositions fulfilling this containment condition. Anyway, we demand \leq_c rather than containment since with the latter one allows skew shapes for which standard composition tableaux (which we will define next) do not exist. For instance, the compositions $\beta = (1, 1)$ and $\alpha = (1, 2)$ satisfy the containment condition but $\beta \not\leq_c \alpha$. Even if $\alpha // \beta$ were a skew shape, there would be no standard composition tableau of this shape (because of the triple rule stated below).

Let D be a diagram. A tableau T of shape D is a map $T: D \rightarrow \mathbb{N}$. It is visualized by filling each $(i, j) \in D$ with $T(i, j)$.

DEFINITION 2.9. Let $\alpha // \beta$ be a skew shape of size n . A standard composition tableau (SCT) of shape $\alpha // \beta$ is a bijective filling $T: \alpha // \beta \rightarrow [n]$ satisfying the following conditions:

- (1) The entries are decreasing in each row from left to right.
- (2) The entries are increasing in the first column from top to bottom.
- (3) (Triple rule). Set $T(i, j) := \infty$ for all $(i, j) \in \beta$. If $(j, k) \in \alpha // \beta$ and $(i, k-1) \in \alpha$ such that $j > i$ and $T(j, k) < T(i, k-1)$ then $(i, k) \in \alpha$ and $T(j, k) < T(i, k)$.

Let $a := T(j, k)$, $b := T(i, k-1)$ be two entries of an SCT T occurring in adjacent columns. Then the triple rule can be visualized as follows by considering the positions of entries in T :



The set of standard composition tableaux of shape $\alpha//\beta$ is denoted with $\text{SCT}(\alpha//\beta)$. For an SCT T we write $\text{sh}(T)$ for its shape. The notions of outer and inner shape are carried over from $\text{sh}(T)$ to T . We call T *straight* if its shape is straight.

EXAMPLE 2.10. An SCT is shown below.

$$T = \begin{array}{|c|c|c|c|} \hline 2 & & & \\ \hline & 5 & 4 & 1 \\ \hline & & 3 & \\ \hline \end{array}$$

We have $\text{osh}(T) = (1, 4, 3)$ and $\text{ish}(T) = (1, 2)$.

Standard composition tableaux encode saturated chains of \mathcal{L}_c in the following way.

PROPOSITION 2.11 (see [3, Proposition 2.11]). *Let $\alpha//\beta$ be a skew composition of size n . For $T \in \text{SCT}(\alpha//\beta)$,*

$$\beta = \alpha^n \prec_c \alpha^{n-1} \prec_c \dots \prec_c \alpha^0 = \alpha$$

given by

$$(1) \quad \alpha^n = \beta, \quad \alpha^{k-1} = \alpha^k \cup T^{-1}(k) \quad \text{for } k = 1, \dots, n$$

is a saturated chain in \mathcal{L}_c . Moreover, we obtain a bijection from $\text{SCT}(\alpha//\beta)$ to the set of saturated chains in \mathcal{L}_c from β to α by mapping each tableau of $\text{SCT}(\alpha//\beta)$ to its corresponding chain given by (1).

EXAMPLE 2.12. The SCT from Example 2.10 corresponds to the chain from Example 2.7.

From the perspective of Proposition 2.11, the triple rule reflects the fact that by adding a cell to a row of a composition diagram, a covering relation in \mathcal{L}_c is established if and only if the row in question is the topmost row of its length.

Some of the upcoming notions already played a role in [14]. Let (i, j) and (i', j') be two cells. Define $r(i, j) := i$ and $c(i, j) := j$ the *row* and the *column* of (i, j) , respectively. We say that (i, j) *attacks* (i', j') and write $(i, j) \rightsquigarrow (i', j')$ if $j = j'$ and $i \neq i'$ or $j = j' - 1$ and $i < i'$. That is, the two cells are distinct and either they appear in the same column or they appear in adjacent columns such that (i', j') is located strictly below and right of (i, j) . We call (i, j) the *left neighbor* of (i', j') if $i = i'$ and $j = j' - 1$.

Let T be an SCT and $i, j \in T$ be two entries. We refer to the *row* and the *column* of i in T by $r_T(i) := r(T^{-1}(i))$ and $c_T(i) := c(T^{-1}(i))$, respectively. We say that i *attacks* j in T and write $i \rightsquigarrow_T j$ if $T^{-1}(i) \rightsquigarrow T^{-1}(j)$. Note that $i \rightsquigarrow_T j$ implies $i \neq j$. If $T^{-1}(j)$ is the left neighbor of $T^{-1}(i)$ then we also call j the *left neighbor* of i .

For two sets of cells $C_1, C_2 \subseteq \mathbb{N}^2$ we say C_1 *attacks* C_2 and write $C_1 \rightsquigarrow C_2$ if there are cells $c_1 \in C_1$ and $c_2 \in C_2$ such that $c_1 \rightsquigarrow c_2$. If $c(c_1) \leq c(c_2)$ for all $c_1 \in C_1, c_2 \in C_2$ then C_1 is called *left* of C_2 . If $c(c_1) < c(c_2)$ for all $c_1 \in C_1, c_2 \in C_2$, C_1 is *strictly left* of C_2 . To simplify notation we may replace singletons by their respective element. For instance, given a cell c_1 we may write $c_1 \rightsquigarrow C_2$ instead of $\{c_1\} \rightsquigarrow C_2$.

In the same way we use these notions for sets of entries of an SCT.

EXAMPLE 2.13. Consider the standard composition tableau T from Example 2.10. We have $2 \rightsquigarrow_T 5, 3 \rightsquigarrow_T 4, 4 \rightsquigarrow_T 3, 5 \rightsquigarrow_T 3$ and $i \not\rightsquigarrow_T j$ for all other pairs of entries. Moreover, 3 is left of $\{1, 4\}$ in T and $2 \rightsquigarrow_T \{3, 5\}$.

DEFINITION 2.14. *Let T be an SCT of size n .*

- (1) $D(T) = \{i \in [n - 1] \mid c_T(i) \leq c_T(i + 1)\}$ is the descent set of T .
- (2) $AD(T) = \{i \in D(T) \mid i \rightsquigarrow_T i + 1\}$ is the set of attacking descents of T .
- (3) $nAD(T) = \{i \in D(T) \mid i \notin AD(T)\}$ is the set of non-attacking descents of T .

- (1') $D^c(T) = \{i \in [n - 1] \mid c_T(i + 1) < c_T(i)\} = [n - 1] \setminus D(T)$ is the ascent set of T .
- (2') $ND^c(T) = \{i \in D^c(T) \mid i + 1 \text{ is the left neighbor of } i\}$ is the set of neighborly ascents of T .

EXAMPLE 2.15. Let T be the tableau from Example 2.10. Then $D(T) = \{2, 3\}$, $AD(T) = \{3\}$, $D^c(T) = \{1, 4\}$ and $ND^c(T) = \{4\}$.

2.3. 0-HECKE MODULES OF STANDARD COMPOSITION TABLEAUX. In this subsection we consider the skew 0-Hecke modules $\mathbf{S}_{\alpha//\beta}$ and $\mathbf{S}_{\alpha//\beta,E}$ introduced in [14] and review related results. This includes the special cases \mathbf{S}_α and $\mathbf{S}_{\alpha,E}$.

THEOREM 2.16 ([14, Theorem 9.8]). Let $\alpha//\beta$ be a skew composition of size n . Then $\mathbf{S}_{\alpha//\beta} := \text{span}_{\mathbb{K}} \text{SCT}(\alpha//\beta)$ is an $H_n(0)$ -module with respect to the following action. For $T \in \text{SCT}(\alpha//\beta)$ and $i = 1, \dots, n - 1$,

$$\pi_i T = \begin{cases} T & \text{if } i \notin D(T) \\ 0 & \text{if } i \in AD(T) \\ s_i T & \text{if } i \in nAD(T) \end{cases}$$

where $s_i T$ is the tableau obtained from T by interchanging i and $i + 1$.

The module \mathbf{S}_α is called *straight* if $\alpha = \alpha//\beta$ is a composition. Even though the main results of this paper are only for straight modules, we consider the more general concept of skew modules here as they naturally arise in the context of the 0-Hecke action on chains of \mathcal{L}_c in Section 3.

EXAMPLE 2.17. Consider the SCT $T = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline 6 & 5 & 4 & 3 \\ \hline 8 & 7 & 2 & \\ \hline \end{array}$. Then $D(T) = \{1, 2, 6\}$,

$$\pi_i T = \begin{cases} T & \text{for } i = 3, 4, 5, 7 \\ 0 & \text{for } i = 6 \\ s_i T & \text{for } i = 1, 2, \end{cases} \quad s_1 T = \begin{array}{|c|c|c|c|} \hline 2 & & & \\ \hline 6 & 5 & 4 & 3 \\ \hline 8 & 7 & 1 & \\ \hline \end{array} \quad \text{and} \quad s_2 T = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline 6 & 5 & 4 & 2 \\ \hline 8 & 7 & 3 & \\ \hline \end{array}.$$

We now decompose $\mathbf{S}_{\alpha//\beta}$ as in [14]. To do this we use an equivalence relation. Let $\alpha//\beta$ be a skew composition of size n and $T_1, T_2 \in \text{SCT}(\alpha//\beta)$. The equivalence relation \sim on $\text{SCT}(\alpha//\beta)$ is given by

$$T_1 \sim T_2 \iff \text{in each column the relative orders of entries in } T_1 \text{ and } T_2 \text{ coincide.}$$

For example, the tableaux shown in Figure 1(a) form an equivalence class under \sim as well as the tableaux shown in Figure 1(b). We denote the *set of equivalence classes* under \sim on $\text{SCT}(\alpha//\beta)$ by $\mathcal{E}(\alpha//\beta)$.

For $E \in \mathcal{E}(\alpha//\beta)$ define $\mathbf{S}_{\alpha//\beta,E} = \text{span}_{\mathbb{K}} E$. It is easy to see that the definition of the 0-Hecke action on standard composition tableaux in Theorem 2.16 implies that $\mathbf{S}_{\alpha//\beta,E}$ is an $H_n(0)$ -submodule of $\mathbf{S}_{\alpha//\beta}$. Thus we have the following.

PROPOSITION 2.18 ([14, Lemma 6.6]). Let $\alpha//\beta$ be a skew composition. Then we have $\mathbf{S}_{\alpha//\beta} = \bigoplus_{E \in \mathcal{E}(\alpha//\beta)} \mathbf{S}_{\alpha//\beta,E}$ as $H_n(0)$ -modules.

The main result of this paper is that the $H_n(0)$ -endomorphism ring of each straight module $\mathbf{S}_{\alpha,E}$ is $\mathbb{K} \text{id}$ and, therefore, we obtain a decomposition of \mathbf{S}_α into indecomposable submodules from Proposition 2.18.

Let $\alpha//\beta$ be a skew composition of size n and $E \in \mathcal{E}(\alpha//\beta)$. We continue by studying E and its module $\mathbf{S}_{\alpha//\beta,E}$ more deeply. First, we consider a partial order \preceq on E .

It will turn out that (E, \preceq) is a graded lattice. Afterwards, we prepare two technical results, Corollary 2.24 and Proposition 2.25, on the 0-Hecke action on standard composition tableaux for later use.

Suppose $T_1, T_2 \in E$. In [14, Section 4] it is shown that a partial order \preceq on E is given by

$$T_1 \preceq T_2 \iff \exists \sigma \in \mathfrak{S}_n \text{ such that } \pi_\sigma T_1 = T_2.$$

We refer to the poset (E, \preceq) simply by E . Two examples are shown in Figure 1. The following theorem summarizes results of [14, Section 6].




FIGURE 1. Two posets each of them given by an equivalence class of standard composition tableaux and the corresponding partial order \preceq . Each covering relation is labeled with the 0-Hecke generator π_i realizing it.

THEOREM 2.19. *Let $\alpha//\beta$ be a skew composition, $E \in \mathcal{E}(\alpha//\beta)$ and $T \in E$.*

- (1) *The tableau T is minimal according to \preceq if and only if $D^c(T) = ND^c(T)$. There is a unique tableau $T_{0,E} \in E$ which satisfies these conditions called source tableau of E .*
- (2) *The tableau T is maximal according to \preceq if and only if $D(T) = AD(T)$. There is a unique tableau $T_{1,E} \in E$ which satisfies these conditions called sink tableau of E .*

In particular, $\mathcal{S}_{\alpha//\beta,E}$ is a cyclic module generated by $T_{0,E}$.

Source and sink tableaux can be observed in Figure 1. Next, we establish a connection between E and an interval of the left weak order. To do this we use the notion of *column words*. Given $T \in \text{SCT}(\alpha//\beta)$ and $j \geq 1$, let w_j be the word obtained by reading the entries in the j th column of T from top to bottom. Then $\text{col}_T = w_1 w_2 \cdots$ is the *column word* of T . Clearly, col_T can be regarded as an element of \mathfrak{S}_n (in one-line notation).

EXAMPLE 2.20. The column word of the tableau $T_{0,E}$ from Figure 1(a) is given by $\text{col}_{T_{0,E}} = 16857423 \in \mathfrak{S}_8$.

LEMMA 2.21 ([14, Lemma 4.4]). *Let T_1 be an SCT, $i \in nAD(T_1)$ and $T_2 = \pi_i T_1$. Then $\text{col}_{T_2} = s_i \text{col}_{T_1}$ and $\ell(\text{col}_{T_2}) = \ell(\text{col}_{T_1}) + 1$. That is, col_{T_2} covers col_{T_1} in left weak order.*

The following statement is similar to [14, Lemma 4.3].

LEMMA 2.22. *Let T_1 and T_2 be two standard composition tableaux such that $\pi_{i_p} \cdots \pi_{i_1} T_1 = T_2$. Then there is a subsequence j_q, \dots, j_1 of i_p, \dots, i_1 such that*

- (1) $T_2 = \pi_{j_q} \cdots \pi_{j_1} T_1$,
- (2) $s_{j_q} \cdots s_{j_1}$ is a reduced word for $\text{col}_{T_2} \text{col}_{T_1}^{-1}$.

In particular, $T_2 = \pi_{\text{col}_{T_2} \text{col}_{T_1}^{-1}} T_1$.

Proof. It follows from the definition of the 0-Hecke operation that we can find a subsequence j_q, \dots, j_1 of i_p, \dots, i_1 of minimal length such that $T_2 = \pi_{j_q} \cdots \pi_{j_1} T_1$. If $q = 0$ then $T_2 = T_1$ and the result is trivial. If $q = 1$ set $i := j_1$. Then by the minimality of q , $T_2 \neq T_1$ and thus $i \in nAD(T_1)$. Now Lemma 2.21 shows that s_i is a reduced word for $\text{col}_{T_2} \text{col}_{T_1}^{-1}$. If $q > 1$ use the case $q = 1$ iteratively. \square

THEOREM 2.23 ([14, Theorem 6.18]). *Let $\alpha//\beta$ be a skew composition, $E \in \mathcal{E}(\alpha//\beta)$ and $I = [\text{col}_{T_{0,E}}, \text{col}_{T_{1,E}}]$ an interval in left weak order. Then the map $\text{col}: E \rightarrow I$, $T \mapsto \text{col}_T$ is a poset isomorphism. In particular, E is a graded lattice with rank function $\delta: T \mapsto \ell(\text{col}_T \text{col}_{T_{0,E}}^{-1})$.*

Actually, Theorem 2.19, Lemma 2.21 and Lemma 2.22 are everything needed to prove Theorem 2.23 as in [14]. They imply that col (and its inverse) map maximal chains to maximal chains. Note that it follows from Theorem 2.23 and Proposition 2.1 that for $T_1 \preceq T_2$ saturated chains from T_1 to T_2 correspond to reduced words for $\text{col}_{T_2} \text{col}_{T_1}^{-1}$.

COROLLARY 2.24. *Let T_1 and T_2 be two standard composition tableaux of size n and $\sigma \in \mathfrak{S}_n$ such that $T_2 = \pi_\sigma T_1$. Then T_1 and T_2 belong to the same equivalence class under \sim . Let δ be the rank function of that class. Then*

- (1) $\delta(T_2) - \delta(T_1) = \ell(\text{col}_{T_2} \text{col}_{T_1}^{-1})$,
- (2) $\delta(T_2) - \delta(T_1) \leq \ell(\sigma)$ where we have equality if and only if $\sigma = \text{col}_{T_2} \text{col}_{T_1}^{-1}$.

Proof. Since $T_2 = \pi_\sigma T_1$, $T_2 \sim T_1$. We obtain part (1) from the above discussion. Part (2) is a consequence of (1) and Lemma 2.22. \square

We finish this section by preparing another result for Section 4.

PROPOSITION 2.25. *Let T be an SCT, $i, j \in T$ be such that $i < j$ and $\square = T^{-1}(i)$. If in T i is located left of $[i + 1, j]$ and does not attack $[i + 1, j]$ then*

- (1) $T' := \pi_{j-1} \cdots \pi_{i+1} \pi_i T$ is an SCT,
- (2) $s_{j-1} \cdots s_{i+1} s_i$ is a reduced word for $\text{col}_{T'} \text{col}_T^{-1}$,
- (3) $T'(\square) = j$.

Proof. Let T be an SCT and $i, j \in T$ such that $i < j$, i is located left of $[i + 1, j]$ and $i \not\rightarrow [i + 1, j]$. Set $\square = T^{-1}(i)$. We do an induction on $m := j - i$. If $m = 1$ then $i \in nAD(T)$ and $T' = \pi_i T$. Thus, (1) and (3) hold by the definition of the 0-Hecke action and (2) is a consequence of Lemma 2.21.

Now, let $m > 1$. Since by assumption i is located left of $[i + 1, j]$ and $i \not\rightarrow [i + 1, j]$, we can apply the induction hypothesis on i and $j - 1$ and obtain that $T'' := \pi_{j-2} \cdots \pi_{i+1} \pi_i T$ is an SCT, $s_{j-2} \cdots s_{i+1} s_i$ is a reduced word for $\text{col}_{T''} \text{col}_T^{-1}$ and $T''(\square) = j - 1$. Since the operators $\pi_{j-2}, \dots, \pi_{i+1}, \pi_i$ are unable to move j , we have $T''^{-1}(j) = T^{-1}(j)$. By choice of i and j , $\square \not\rightarrow T^{-1}(j) = T''^{-1}(j)$ and \square is left of $T''^{-1}(j)$. Thus, $j - 1 \in nAD(T'')$ so that $T' = \pi_{j-1} \pi_{j-2} \cdots \pi_i T = \pi_{j-1}(T'')$ is an SCT and $T'(\square) = j$. It follows from Lemma 2.21 that $\text{col}_{T'} \text{col}_T^{-1} = s_{j-1} \text{col}_{T''} \text{col}_T^{-1} = s_{j-1} s_{j-2} \cdots s_i$ and that $s_{j-1} s_{j-2} \cdots s_i$ is a reduced word. \square

3. A 0-HECKE ACTION ON CHAINS OF THE COMPOSITION POSET

In Proposition 2.11 a bijection between saturated chains in the composition poset \mathcal{L}_c and standard composition tableaux is given. In this section we study the 0-Hecke action on these chains induced by this bijection. The main result of this section, Proposition 3.8, will be useful in Section 4. We begin with some notation.

DEFINITION 3.1. Let T be an SCT of shape $\alpha // \beta$ and size n , $m \in [0, n]$ and $\beta = \alpha^n \prec_c \alpha^{n-1} \prec_c \cdots \prec_c \alpha^0 = \alpha$ the chain in \mathcal{L}_c corresponding to T . The SCT of shape $\alpha^m // \beta$ corresponding to the chain $\alpha^n \prec_c \alpha^{n-1} \prec_c \cdots \prec_c \alpha^m$ is denoted by $T^{>m}$.

EXAMPLE 3.2. For $T =$

1		
		3
	2	

 we have $T^{>2} =$

			1

.

The following Lemma shows how we can obtain $T^{>m}$ directly from T .

LEMMA 3.3. Let T be an SCT of size n and shape $\alpha // \beta$, $\beta = \alpha^n \prec_c \alpha^{n-1} \prec_c \cdots \prec_c \alpha^0 = \alpha$ the chain in \mathcal{L}_c corresponding to T and $m \in [0, n]$.

- (1) $\alpha^m = \text{osh}(T^{>m})$.
- (2) We obtain $T^{>m}$ from T by removing the cells containing $1, \dots, m$ and subtracting m from the remaining entries.

Proof. Part (1) is an immediate consequence of Definition 3.1. By Proposition 2.11, we obtain $T^{>m}$ by successively adding cells with entries $n - m, \dots, 1$ to the inner shape β at exactly the same positions where we would add $n, \dots, m + 1$ to β in order to obtain T from its corresponding chain. This implies part (2). \square

With the first part of Lemma 3.3 we can access the compositions within a chain of a given SCT. We use the following preorder to describe how the 0-Hecke action affects these compositions.

DEFINITION 3.4.

- (1) For a composition $\alpha = (\alpha_1, \dots, \alpha_l)$ of n and $j \in \mathbb{N}$ we define $|\alpha|_j = \#\{i \in [l] \mid \alpha_i \geq j\}$.
- (2) On the set of compositions of size n we define the preorder \trianglelefteq by

$$\alpha \trianglelefteq \beta \iff \sum_{j=1}^k |\beta|_j \leq \sum_{j=1}^k |\alpha|_j \text{ for all } k \geq 1.$$

Moreover, set $\alpha \triangleleft \beta \iff \alpha \trianglelefteq \beta$ and $\alpha \neq \beta$.

Note that $|\alpha|_j$ is the number of cells in the j th column of the diagram of α . Obviously \trianglelefteq is reflexive and transitive. It is not antisymmetric since for example $(2, 1) \trianglelefteq (1, 2)$ and $(1, 2) \trianglelefteq (2, 1)$. In general, for $\alpha, \beta \vDash n$ we have

$$\alpha \trianglelefteq \beta \text{ and } \beta \trianglelefteq \alpha \iff |\alpha|_j = |\beta|_j \text{ for all } j = 1, 2, \dots \iff \tilde{\alpha} = \tilde{\beta}.$$

EXAMPLE 3.5.



If we restrict \trianglelefteq to partitions, we obtain the well known dominance order appearing, for example, in [13]. However, \trianglelefteq on partitions may seem to be reversed to the dominance order. This is because in the definition above we are considering the number of cells in columns rather than in rows as usual.

LEMMA 3.6. Let $\alpha // \beta$ be a skew composition of size n and $T_1, T_2 \in \text{SCT}(\alpha // \beta)$ be such that $T_2 = \pi_i T_1$ for an $i \in nAD(T_1)$. Then


$$\begin{aligned} \text{osh}(T_2^{>i}) &\triangleleft \text{osh}(T_1^{>i}), \\ \text{osh}(T_2^{>m}) &= \text{osh}(T_1^{>m}) \text{ for } m \in [0, n], m \neq i. \end{aligned}$$

Proof. Assume T_1, T_2 and i as in the assertion. Then T_2 is the tableau obtained from T_1 by swapping the entries i and $i + 1$ of T_1 . Let $m \in [0, n]$.

If $m \neq i$ then either $\{i, i + 1\} \subseteq [1, m]$ or $\{i, i + 1\} \cap [1, m] = \emptyset$. Therefore, $T_1^{-1}([1, m]) = T_2^{-1}([1, m])$ and so from the perspective of Lemma 3.3 we remove the same set of cells from T_1 to obtain $T_1^{>m}$ as we remove from T_2 to obtain $T_2^{>m}$. That is, $\text{sh}(T_1^{>m}) = \text{sh}(T_2^{>m})$.

If $m = i$, set $(r_k, c_k) := T_1^{-1}(k)$ for $k = i, i + 1$, $\gamma_1 := \text{osh}(T_1^{>i})$ and $\gamma_2 := \text{osh}(T_2^{>i})$. We assume that all composition diagrams appearing here are moved to the bottom of α . Observe that as $T_2 = s_i T_1$, one obtains $\text{sh}(T_2^{>i})$ from $\text{sh}(T_1^{>i})$ by moving the cell (r_{i+1}, c_{i+1}) to the position (r_i, c_i) . Since $\text{ish}(T_2^{>i}) = \beta = \text{ish}(T_1^{>i})$, we obtain γ_2 from γ_1 by this movement. Moreover, $i \in nAD(T_1)$ implies $c_i < c_{i+1}$. That is, we obtain γ_2 from γ_1 by moving a cell strictly to the left. By the definition of \trianglelefteq , this means that $\gamma_2 \triangleleft \gamma_1$. \square

EXAMPLE 3.7. The $H_n(0)$ -action on tableaux and the corresponding chains of the composition poset is shown below.



Let $\alpha // \beta$ be a skew composition, $E \in \mathcal{E}(\alpha // \beta)$ and $T_1, T_2 \in E$ be such that $T_1 \preceq T_2$. Recall that for each saturated chain from T_1 to T_2 in E the index set of the 0-Hecke operators establishing the covering relations within the chain is $\text{supp}(\text{col}_{T_2} \text{col}_{T_1}^{-1})$.

As a consequence of Lemma 3.6 we obtain a criterion for determining whether an operator π_i appears in the saturated chains from T_1 to T_2 or not.

PROPOSITION 3.8. *Let $\alpha // \beta$ be a skew composition of size n , $i \in [n - 1]$, $E \in \mathcal{E}(\alpha // \beta)$ and $T_1, T_2 \in E$ be such that $T_1 \preceq T_2$. Then $i \in \text{supp}(\text{col}_{T_2} \text{col}_{T_1}^{-1})$ if and only if $\text{sh}(T_2^{>i}) \neq \text{sh}(T_1^{>i})$.*

Proof. Lemma 3.6 applied to each covering relation in a saturated chain from T_1 to T_2 in E and the fact that \preceq is a preorder imply

$$i \in \text{supp}(\text{col}_{T_2} \text{col}_{T_1}^{-1}) \iff \text{osh}(T_2^{>i}) \neq \text{osh}(T_1^{>i}).$$

From this we obtain the claim since $\text{ish}(T_1^{>i}) = \beta = \text{ish}(T_2^{>i})$. □

4. THE ENDOMORPHISM RING OF $\mathcal{S}_{\alpha, E}$

For each $\alpha \vDash n$ there is an equivalence class $E_\alpha \in \mathcal{E}(\alpha)$ such that for all $T \in E_\alpha$ the entries increase in each column from top to bottom [14, Section 8]. In [14], Tewari and van Willigenburg showed that $\mathcal{S}_{\alpha, E_\alpha}$ is indecomposable.

In this section, we show for all $E \in \mathcal{E}(\alpha)$ that $\text{End}_{H_n(0)}(\mathcal{S}_{\alpha, E}) = \mathbb{K} \text{id}$ and hence $\mathcal{S}_{\alpha, E}$ is indecomposable; this extends the result of Tewari and van Willigenburg to the general case. By Proposition 2.18 we then have the desired decomposition of \mathcal{S}_α . In contrast, skew modules $\mathcal{S}_{\alpha // \beta, E}$ can be decomposable (see Example 4.13 at the end of the section).

We fix some notation that we use in the entire section unless stated otherwise. Let $\alpha \vDash n$, $E \in \mathcal{E}(\alpha)$ and $T_0 := T_{0, E}$ be the source tableau of E . Moreover, let $f \in \text{End}_{H_n(0)}(\mathcal{S}_{\alpha, E})$, $v := f(T_0)$ and $v = \sum_{T \in E} a_T T$ be the expansion of v in the \mathbb{K} -basis E . Since $\mathcal{S}_{\alpha, E}$ is cyclically generated by T_0 , f is already determined by v . The support of v is given by $\text{supp}(v) = \{T \in E \mid a_T \neq 0\}$. Our goal is to show that T_0 is the only tableau that may occur in $\text{supp}(v)$ since then $f = a_{T_0} \text{id} \in \mathbb{K} \text{id}$. We begin with a property holding for $\text{supp}(v)$ that also appeared in the proof of [14, Theorem 7.8].

LEMMA 4.1. *If $T \in \text{supp}(v)$ then $D(T) \subseteq D(T_0)$.*

Proof. Let $T_* \in E$ be such that $D(T_*) \not\subseteq D(T_0)$. Then there is an $i \in D(T_*) \cap D^c(T_0)$. Because $i \in D^c(T_0)$, $\pi_i v = f(\pi_i T_0) = v$. Thus, a_{T_*} is the coefficient of T_* in $\pi_i v = \sum_{T \in E} a_T \pi_i T$. But this coefficient is 0 since $\pi_i T \neq T_*$ for all $T \in E$. To see this, assume that there is a $T \in E$ such that $\pi_i T = T_*$. Then we obtain a contradiction as

$$T_* \neq \pi_i T_* = \pi_i^2 T = \pi_i T = T_*. \quad \square$$

Thanks to Lemma 4.1 it remains to show $a_T = 0$ for all $T \in E$ such that $T \neq T_0$ and $D(T) \subseteq D(T_0)$. So fix such a tableau T . In order to determine a_T , we will use a 0-Hecke operator π_σ where $\sigma = s_{j-1} \cdots s_i$ and i and j are given by

$$(2) \quad \begin{aligned} i &= \max \{k \in [n] \mid T^{-1}(k) \neq T_0^{-1}(k)\}, \\ j &= \min \{k \in [n] \mid k > i \text{ and } i \rightsquigarrow_{T_0} k\}. \end{aligned}$$

That is, i is the greatest entry whose position in T differs from that in T_0 and j is the smallest entry in T_0 which is greater than i and attacked by i in T_0 . At this point it

is not clear that j is well defined since the defining set could be empty. However, the next two lemmas will show that there always exists an element in this set.

EXAMPLE 4.2. Consider the equivalence class E from Figure 1(a). Then $T_0 = T_{0,E}$ and there is exactly one other tableau T in E with $D(T) \subseteq D(T_0)$:

$$T_0 = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline 6 & 5 & 4 & 3 \\ \hline 8 & 7 & 2 & \\ \hline \end{array} \xrightarrow{\pi_1} T = \begin{array}{|c|c|c|c|} \hline 2 & & & \\ \hline 6 & 5 & 4 & 3 \\ \hline 8 & 7 & 1 & \\ \hline \end{array}$$

Defining i and j for T as in (2), we obtain $i = 2$ and $j = 4$. Note that $2 \in D(T_0)$. This property holds in general by the following result.

LEMMA 4.3. Let $T \in E$ be such that $T \neq T_0$ and $D(T) \subseteq D(T_0)$ and set

$$i = \max \{k \in [n] \mid T^{-1}(k) \neq T_0^{-1}(k)\}.$$

Then $i \in D(T_0)$.

Proof. Let T, T_0 and i be given as in the assertion. We introduce indices such that $D(T_0) = \{d_1 < d_2 < \dots < d_m\}$ and set $d_0 := 0, d_{m+1} := n$. Moreover, define $I_k := [d_{k-1} + 1, d_k]$ for $k = 1, \dots, m + 1$. Recall that since T_0 is a source tableau, $D^c(T_0) = ND^c(T_0)$ by Theorem 2.19. That is, $a+1$ is the left neighbor of a for each ascent a of T_0 . Therefore, we have $I_k \setminus \{d_k\} \subset ND^c(T_0)$ and conclude that $T_0^{-1}(I_k)$ is a connected horizontal strip (a one-row diagram which contains all cells between its leftmost and rightmost cell) for $k = 1, \dots, m + 1$.

Set $\square_k := T_0^{-1}(k)$ for $k = 1, \dots, n$ and let x be the index such that $T(\square_x) = i$. Since T_0 and T are straight, the ordering conditions of standard composition tableaux imply $T^{-1}(n) = (\ell(\alpha), 1) = T_0^{-1}(n)$. Therefore $i \neq n$ and we now show $i \notin D^c(T_0)$.

Assume for sake of contradiction that $i \in D^c(T_0)$. Let $l \in [m + 1]$ be such that $i \in I_l$. Since $i \in D^c(T_0)$, $i < d_l$ and $i + 1 \in I_l$. The strip $T_0^{-1}(I_l)$ looks as follows:

$$(3) \quad \square_{d_l} \square_{d_l-1} \cdots \square_{i+1} \square_i \cdots \square_{d_{l-1}+1}.$$

By choice of i , we have

$$(4) \quad T(\square_k) = k \text{ for } k = i + 1, \dots, n \text{ and } T(\square_i) < i.$$

Since entries decrease in rows of T , (3) implies

$$(5) \quad T(\square_k) < i \text{ for } k = d_{l-1} + 1, \dots, i.$$

Combining (4) and (5) we obtain

$$(6) \quad x \leq d_{l-1}.$$

We deal with two cases depending on $c_T(i)$. In both cases we will end up with a contradiction.

CASE 1. $c_T(i) \leq c_{T_0}(d_{l-1} + 1)$. It follows from $D(T) \subseteq D(T_0)$ that $i \in D^c(T)$ and thus $c_T(i + 1) < c_T(i)$. Using $c_{T_0}(i) = c_{T_0}(i + 1) + 1 = c_T(i + 1) + 1$, we obtain $c_{T_0}(i) \leq c_T(i) \leq c_{T_0}(d_{l-1} + 1)$. Then there is a $y \in [d_{l-1} + 1, i]$ such that \square_x and \square_y are in the same column. On the one hand, we obtain from (5) that $T(\square_y) < i = T(\square_x)$. On the other hand, the choice of y and (6) imply $y > d_{l-1} \geq x$ and hence $T_0(\square_y) = y > x = T_0(\square_x)$. That is, in the column of \square_x and \square_y the relative order of entries in T differs from that in T_0 . So $T \not\sim T_0$ which contradicts the assumption $T, T_0 \in E$.

CASE 2. $c_T(i) > c_{T_0}(d_{l-1} + 1)$. This case is illustrated in Figure 2. Since by (6) $x \leq d_{l-1}$, there is a $1 \leq p \leq l - 1$ such that $x \in I_p$. The leftmost cell of the connected horizontal strip $T_0^{-1}(I_p)$ is \square_{d_p} . As entries decrease in rows of T from left to right, we have $T(\square_{d_p}) \geq T(\square_x) = i$. In addition, the choice of p and (4) imply that $T(\square_{d_p}) \leq i$. Thus, $d_p = x$.

From $d_p = x$ we obtain $d_p \neq d_{l-1}$ since

$$c_{T_0}(d_{l-1}) \leq c_{T_0}(d_{l-1} + 1) < c_T(i) = c_{T_0}(d_p)$$

where we use $d_{l-1} \in D(T_0)$ for the first inequality.

We claim that there exists an index $y \in [d_p + 1, d_{l-1} - 1]$ such that \square_y and \square_{d_p} are located in the same column. To prove the claim, assume for sake of contradiction that this is not the case. Then $d_p \in D(T_0)$ implies $c_{T_0}(d_p) < c_{T_0}(d_p + 1)$. Thus, it follows from $D^c(T_0) = ND^c(T_0)$ and induction that $c_{T_0}(d_p) < c_{T_0}(z)$ for all $z \in [d_p + 1, d_{l-1} - 1]$. As a consequence,

$$c_{T_0}(d_{l-1}) < c_{T_0}(d_p) < c_{T_0}(d_{l-1} - 1).$$

In other words, $d_{l-1} - 1$ is an ascent of T_0 but d_{l-1} is not the left neighbor of $d_{l-1} - 1$. This is a contradiction to the fact that T_0 is a source tableau and finishes the proof of the claim.

Now, let y be as claimed above. Then $y \in [d_p + 1, d_{l-1} - 1]$ and in particular $y \neq d_p = x$. Hence, (4) implies $T(\square_y) < i$ and so $T(\square_y) < i = T(\square_{d_p})$. On the other hand, $y \in [d_p + 1, d_{l-1} - 1]$ yields $T_0(\square_y) = y > d_p = T_0(\square_{d_p})$. As in Case 1, this is a contradiction to $T, T_0 \in E$. \square




FIGURE 2. An example for the positions of cells and entries in the tableau T from Case 2 of the proof of Lemma 4.3.

Note that the i appearing in the following Lemma is not the same as in (2).

LEMMA 4.4. For all $i \in D(T_0)$ there exists $k \in T_0$ such that $k > i$ and $i \rightsquigarrow_{T_0} k$.

Proof. Let $i \in D(T_0)$. Then $c_{T_0}(i) \leq c_{T_0}(i + 1)$ and thus $r_{T_0}(i) \neq r_{T_0}(i + 1)$. Since T_0 is straight by assumption, the cell $(r_{T_0}(i + 1), c_{T_0}(i))$ is contained in the shape of T_0 . Let k be the entry of T_0 in that cell. Then $i \rightsquigarrow_{T_0} k$ and $k \geq i + 1$ because entries decrease in rows. \square

Let T, i and j as in (2). Lemma 4.3 and Lemma 4.4 now show that j is well defined. We proceed by considering the relative positions of i and $[i + 1, j]$ first in T_0 and then

in T . This will allow us to deduce useful properties of the operator π_σ to be defined in Lemma 4.9. In the following Lemma, i is slightly more general than in (2).

LEMMA 4.5. *Let $i \in D(T_0)$ and set $j = \min\{k \in [n] \mid k > i \text{ and } i \rightsquigarrow_{T_0} k\}$. Then j is well defined and in T_0 i is located strictly left of $[i + 1, j - 1]$ and does not attack $[i + 1, j - 1]$.*

We illustrate Lemma 4.5 before we prove it.

EXAMPLE 4.6. For the source tableau from above

$$T_0 = \begin{array}{|c|c|c|c|} \hline & 1 & & \\ \hline 6 & 5 & 4 & \mathbf{3} \\ \hline 8 & 7 & \mathbf{2} & \\ \hline \end{array}$$

and $i = 2 \in D(T_0)$ we have $j = 4 = \min\{k \in [n] \mid k > i \text{ and } i \rightsquigarrow_{T_0} k\}$ and $\{3\} = [i + 1, j - 1]$. Note $2 \rightsquigarrow_{T_0} 4$ but $2 \not\rightsquigarrow_{T_0} 3$.

Proof of Lemma 4.5. First, it follows from Lemma 4.4 that j is well defined. We set $I := [i + 1, j - 1]$ and $c_l := c_{T_0}(l)$ for $l \in T_0$. By the minimality of j , we have $i \not\rightsquigarrow_{T_0} I$. It remains to show that i is strictly left of I or equivalently that $c_i < c_l$ for all $l \in I$. We may assume $I \neq \emptyset$ and use an induction argument to show this.

We begin with $i + 1$, the minimum of I . Since $i \in D(T_0)$, $c_i \leq c_{i+1}$. Moreover, $i + 1 \in I$ implies $i \not\rightsquigarrow_{T_0} i + 1$ and consequently $c_i < c_{i+1}$.

Now, let $l \in I$ such that $l > i + 1$ and $c_i < c_{l-1}$. If $l - 1 \in D(T_0)$ then $c_i < c_{l-1} \leq c_l$. If $l - 1 \in D^c(T_0)$ then $l - 1 \in ND^c(T_0)$ as T_0 is a source tableau. Thus $c_l = c_{l-1} - 1$ and $c_i \leq c_l$. Furthermore $c_i \neq c_l$ since $i \not\rightsquigarrow_{T_0} I \ni l$. Hence, $c_i < c_l$. \square

Let T, i and j as in (2). By definition, i attacks j in T_0 . In contrast, the next Lemma shows that i does not attack j in T . There, i and j are defined as in (2).

LEMMA 4.7. *Let $T \in E$ be such that $T \neq T_0$ and $D(T) \subseteq D(T_0)$. Define*

$$i = \max \{k \in [n] \mid T^{-1}(k) \neq T_0^{-1}(k)\},$$

$$j = \min \{k \in [n] \mid k > i \text{ and } i \rightsquigarrow_{T_0} k\}.$$

Then i and j are well defined and in T i appears strictly left of $[i + 1, j]$ and does not attack $[i + 1, j]$.

We first give an example and then the proof of Lemma 4.7.

EXAMPLE 4.8. Recall that in our running example $i = 2$ and $j = 4$ when defined for

$$T = \begin{array}{|c|c|c|c|} \hline & \mathbf{2} & & \\ \hline 6 & 5 & 4 & \mathbf{3} \\ \hline 8 & 7 & 1 & \\ \hline \end{array}$$

as in Lemma 4.7. Then $[i + 1, j] = \{3, 4\}$ and $2 \not\rightsquigarrow_T \{3, 4\}$.

Proof of Lemma 4.7. Lemma 4.3 yields $i \in D(T_0)$ and so Lemma 4.4 ensures that j is well defined. Set $\sigma := \text{col}_T \text{col}_{T_0}^{-1}$, $\square_k := T_0^{-1}(k)$ for $k = 1, \dots, n$ and let x be the index such that $T(\square_x) = i$.

By choice of i , we have $T^{>i} = T_0^{>i}$. So, $\text{sh}(T^{>k}) = \text{sh}(T_0^{>k})$ for $k = i, \dots, n$. Hence, from Proposition 3.8 we obtain

$$(7) \quad \text{supp}(\sigma) \subseteq [i - 1].$$

Let $s_{i_p} \cdots s_{i_1}$ be a reduced word for σ . Then $T = \pi_{i_p} \cdots \pi_{i_1} T_0$. From (7) we have $i_q \neq i$ for $q = 1, \dots, p$. Moreover, at least one π_{i_q} has to move i because the position of i in T differs from its position in T_0 . Hence, there is a q such that $i_q = i - 1$ since π_{i-1}

and π_i are the only operators that are able to move i . For two standard composition tableaux T_1 and T_2 such that $T_2 = \pi_{i-1}T_1 = s_{i-1}T_1$ we have that $i - 1 \in nAD(T_1)$ and thus $T_2^{-1}(i)$ is left of $T_1^{-1}(i)$ and $T_2^{-1}(i) \not\prec_T T_1^{-1}(i)$. So, by applying $\pi_{i_p} \cdots \pi_{i_1}$ to T_0 , i is moved (possibly multiple times) strictly to the left into a cell that does not attack \square_i . That is,

$$(8) \quad \square_x \text{ is located strictly left of } \square_i \text{ and } \square_x \not\prec \square_i.$$

It follows from the choice of i that the elements of $[i + 1, j - 1]$ have the same position in T and T_0 . By combining (8) and Lemma 4.5 we obtain:

$$(9) \quad \text{In } T \text{ } i \text{ is located strictly left of } [i + 1, j - 1] \text{ and } i \not\prec_T [i + 1, j - 1].$$

Recall that j has the same position in T and T_0 . It follows from (8) and $i \rightsquigarrow_{T_0} j$ that $c_T(i) < c_{T_0}(i) \leq c_{T_0}(j)$. Thus, i is strictly left of j in T .

It remains to show $i \not\prec_T j$. We have either $c_{T_0}(j) = c_{T_0}(i) + 1$ or $c_{T_0}(j) = c_{T_0}(i)$ since $i \rightsquigarrow_{T_0} j$.

CASE 1. $c_{T_0}(j) = c_{T_0}(i) + 1$. Then (8) implies $c_T(i) < c_{T_0}(i) < c_{T_0}(j) = c_T(j)$ and so $i \not\prec_T j$.

CASE 2. $c_{T_0}(j) = c_{T_0}(i)$. If $c_T(i) < c_{T_0}(i) - 1$ then $c_T(i) < c_T(j) - 1$ and so $i \not\prec_T j$. If $c_T(i) = c_{T_0}(i) - 1$ then i and j appear in adjacent columns of T and for $i \not\prec_T j$ we have to show that $r_T(j) < r_T(i)$. On the one hand, we have $1 \leq c_T(i) < c_{T_0}(i)$ so that i has a left neighbor $t > i$ in T_0 . In addition, from the first statement of Lemma 4.5 and $c_{T_0}(j) = c_{T_0}(i)$ we obtain that i is weakly left of $[i + 1, j]$ in T_0 . Thus, $t > j$ and hence $r_{T_0}(j) < r_{T_0}(i)$ because otherwise t, i and j would violate the triple rule in T_0 . On the other hand, $c_T(i) = c_{T_0}(i) - 1$ and $i \not\prec_T \square_i$ imply $r_{T_0}(i) < r_T(i)$. All in all, $r_T(j) = r_{T_0}(j) < r_{T_0}(i) < r_T(i)$ and thus $i \not\prec_T j$. \square

Next, we prove useful properties of the operators π_σ mentioned already above in (2).

LEMMA 4.9. *Keep the notation of Lemma 4.7 and set $\sigma = s_{j-1} \cdots s_{i+1}s_i$. Then*

- (1) $\pi_\sigma T_0 = 0$,
- (2) $\pi_\sigma T \in E$,
- (3) $\sigma = \text{col}_{\pi_\sigma T} \text{col}_T^{-1}$.

Proof. First observe that $s_{j-1} \cdots s_{i+1}s_i$ is a reduced word, i.e. $\pi_\sigma = \pi_{j-1} \cdots \pi_{i+1}\pi_i$. Set $\square_k = T_0^{-1}(k)$ for $k = 1, \dots, n$.

We consider T_0 . Set $T' = \pi_{j-2} \cdots \pi_{i+1}\pi_i T_0$. We can apply Proposition 2.25 in T_0 to i and $[i + 1, j - 1]$ because of Lemma 4.3 and Lemma 4.5. By doing this, we obtain that $T' \in E$ and $T'(\square_i) = j - 1$. In addition, $T'(\square_j) = T_0(\square_j) = j$ as none of the operators $\pi_{j-2}, \dots, \pi_{i+1}$ moves j . Recall that j is defined such that $\square_i \rightsquigarrow \square_j$. Thus $j - 1 \in AD(T')$ and $\pi_\sigma T_0 = \pi_{j-1}T' = 0$.

Now consider T . Because of Lemma 4.7, we can apply Proposition 2.25 in T on i and $[i + 1, j]$. This immediately gives us (2) and (3). \square

EXAMPLE 4.10. Continuing our running example, we have $i = 2, j = 4$ and $\pi_\sigma = \pi_3\pi_2$. Moreover,

$$\begin{array}{l}
 T_0 = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline 6 & 5 & 4 & 3 \\ \hline 8 & 7 & 2 & \\ \hline \end{array} \xrightarrow{\pi_2} \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline 6 & 5 & 4 & 2 \\ \hline 8 & 7 & 3 & \\ \hline \end{array} \xrightarrow{\pi_3} 0, \\
 \\
 T = \begin{array}{|c|c|c|c|} \hline 2 & & & \\ \hline 6 & 5 & 4 & 3 \\ \hline 8 & 7 & 1 & \\ \hline \end{array} \xrightarrow{\pi_2} \begin{array}{|c|c|c|c|} \hline 3 & & & \\ \hline 6 & 5 & 4 & 2 \\ \hline 8 & 7 & 1 & \\ \hline \end{array} \xrightarrow{\pi_3} \begin{array}{|c|c|c|c|} \hline 4 & & & \\ \hline 6 & 5 & 3 & 2 \\ \hline 8 & 7 & 1 & \\ \hline \end{array}.
 \end{array}$$

We are ready to prove the main result of this paper now.

THEOREM 4.11. *Let $\alpha \vDash n$ and $E \in \mathcal{E}(\alpha)$. Then $\text{End}_{H_n(0)}(\mathbf{S}_{\alpha,E}) = \mathbb{K} \text{id}$. In particular, $\mathbf{S}_{\alpha,E}$ is an indecomposable $H_n(0)$ -module.*

Proof. For the second part, note that if $\text{End}_{H_n(0)}(\mathbf{S}_{\alpha,E}) = \mathbb{K} \text{id}$ then $\mathbf{S}_{\alpha,E}$ is indecomposable.

To prove the first part, let $f \in \text{End}_{H_n(0)}(\mathbf{S}_{\alpha,E})$, $v := f(T_0)$ and $v = \sum_{T \in E} a_T T$ as before. We show $\text{supp}(v) \subseteq \{T_0\}$ since this and the fact that $\mathbf{S}_{\alpha,E}$ is cyclically generated by T_0 imply $f = a_{T_0} \text{id} \in \mathbb{K} \text{id}$.

If $v = 0$, this is clear so that we can assume $v \neq 0$. Recall that by Theorem 2.23 E is a graded poset. We denote its rank function with δ . Let $T_* \in \text{supp}(v)$ be of maximal rank in $\text{supp}(v)$. Assume for sake of contradiction that $T_* \neq T_0$. Then Lemma 4.1 yields $D(T_*) \subseteq D(T_0)$. Hence, Lemma 4.9 provides the existence of $\sigma \in \mathfrak{S}_n$ such that $\pi_\sigma T_* \in E$, $\pi_\sigma T_0 = 0$ and $\sigma = \text{col}_{\pi_\sigma T_*} \text{col}_{T_*}^{-1}$.

We claim that if $T \in \text{supp}(v)$ and $\pi_\sigma T = \pi_\sigma T_*$ then $T = T_*$. To see this, let $T \in \text{supp}(v)$ be such that $\pi_\sigma T = \pi_\sigma T_*$. Then

$$\ell(\sigma) \geq \delta(\pi_\sigma T) - \delta(T) = \delta(\pi_\sigma T_*) - \delta(T) \geq \delta(\pi_\sigma T_*) - \delta(T_*) = \ell(\sigma)$$

where Corollary 2.24 is used to establish the first inequality and the last equality. Hence, $\ell(\sigma) = \delta(\pi_\sigma T) - \delta(T)$ and another application of Corollary 2.24 yields $\text{col}_{\pi_\sigma T_*} \text{col}_T^{-1} = \sigma$. But then

$$\text{col}_{\pi_\sigma T_*} \text{col}_T^{-1} = \sigma = \text{col}_{\pi_\sigma T_*} \text{col}_{T_*}^{-1}$$

so that $\text{col}_T = \text{col}_{T_*}$ and thus $T = T_*$ as claimed.

The claim implies that the coefficient of $\pi_\sigma T_*$ in $\pi_\sigma v = \sum_{T \in \text{supp}(v)} a_T \pi_\sigma T$ is a_{T_*} . Yet, $\pi_\sigma v = f(\pi_\sigma T_0) = 0$ and hence $a_{T_*} = 0$ which contradicts the assumption $T_* \in \text{supp}(v)$ and completes the proof of $\text{supp}(v) \subseteq \{T_0\}$. \square

Combining Theorem 4.11 with Proposition 2.18, we obtain the desired decomposition of \mathbf{S}_α .

COROLLARY 4.12. *Let $\alpha \vDash n$. Then $\mathbf{S}_\alpha = \bigoplus_{E \in \mathcal{E}(\alpha)} \mathbf{S}_{\alpha,E}$ is a decomposition into indecomposable submodules.*

EXAMPLE 4.13. In general, Theorem 4.11 does not hold for skew modules $\mathbf{S}_{\alpha//\beta,E}$. The two tableaux

$$T_0 = \begin{array}{|c|c|c|} \hline 1 & & \\ \hline & & 2 \\ \hline \end{array} \xrightarrow{\pi_1} T_1 = \begin{array}{|c|c|c|} \hline 2 & & \\ \hline & & 1 \\ \hline \end{array}$$

form an equivalence class E . Let $n = 2$ and $\alpha//\beta = \text{sh}(T_0)$. Observe that we obtain an idempotent $H_n(0)$ -endomorphism φ by setting $\varphi(T_0) = \varphi(T_1) = T_1$. Clearly, φ is none of the trivial idempotents $0, \text{id} \in \text{End}_{H_n(0)}(\mathbf{S}_{\alpha//\beta,E})$. Thus, $\text{End}_{H_n(0)}(\mathbf{S}_{\alpha//\beta,E}) \neq \mathbb{K} \text{id}$. Moreover, we obtain a decomposition

$$\mathbf{S}_{\alpha//\beta,E} = \varphi(\mathbf{S}_{\alpha//\beta,E}) \oplus (\text{id} - \varphi)(\mathbf{S}_{\alpha//\beta,E}) = \text{span}_{\mathbb{K}}(T_1) \oplus \text{span}_{\mathbb{K}}(T_0 - T_1)$$

into two submodules of dimension 1.

This example also illustrates how the argumentation of this section can fail when it is applied to skew modules. Note that $D(T_1) \subseteq D(T_0)$. So, we may try to set

$$i = \max \{k \in [n] \mid T_1^{-1}(k) \neq T_0^{-1}(k)\},$$

$$j = \min \{k \in [n] \mid k > i \text{ and } i \rightsquigarrow_{T_0} k\}.$$

as before. But then $i = 2$ so that j does not exist.

Acknowledgements. The new results presented in this article are part of my PhD research, and I would like to thank my supervisor Christine Bessenrodt for her support especially during the work on this paper. I would also like to thank the referees for their valuable comments. This includes the referees of FPSAC 2018 who reviewed the accompanying extended abstract.

REFERENCES

- [1] C. Berg, N. Bergeron, F. Saliola, L. Serrano, and M. Zabrocki, *A lift of the Schur and Hall–Littlewood bases to non-commutative symmetric functions*, *Canad. J. Math.* **66** (2014), 525–565.
- [2] ———, *Indecomposable modules for the dual immaculate basis of quasi-symmetric functions*, *Proc. Amer. Math. Soc.* **143** (2015), 991–1000.
- [3] C. Bessenrodt, K. Luoto, and S. van Willigenburg, *Skew quasisymmetric Schur functions and noncommutative Schur functions*, *Adv. Math.* **226** (2011), 4492–4532.
- [4] A. Björner and F. Brenti, *Combinatorics of Coxeter groups*, Springer Science & Business Media, 2006.
- [5] G. Duchamp, D. Krob, B. Leclerc, and J.-Y. Thibon, *Fonctions quasi-symétriques, fonctions symétriques non commutatives et algèbres de Hecke à $q = 0$* , *C. R. Acad. Sci. Paris Sér. I Math.* **322** (1996), 107–112.
- [6] I. Gessel, *Multipartite P -partitions and inner products of skew Schur functions*, in *Combinatorics and Algebra* (Boulder, Colo., 1983), *Contemp. Math.*, vol. 34, Amer. Math. Soc., Providence, RI, 1984, pp. 289–317.
- [7] D. Grinberg and V. Reiner, *Hopf algebras in combinatorics*, <https://arxiv.org/abs/1409.8356>, 2014.
- [8] J. Haglund, K. Luoto, S. Mason, and S. van Willigenburg, *Quasisymmetric Schur functions*, *J. Combin. Theory Ser. A* **118** (2011), 463–490.
- [9] J. Huang, *A tableau approach to the representation theory of 0-Hecke algebras*, *Ann. Comb.* **20** (2016), no. 4, 831–868.
- [10] D. Krob and J.-Y. Thibon, *Noncommutative symmetric functions IV: Quantum linear groups and Hecke algebras at $q = 0$* , *J. Algebraic Combin.* **6** (1997), 339–376.
- [11] A. Mathas, *Iwahori–Hecke algebras and Schur algebras of the symmetric group*, vol. 15, American Mathematical Soc., 1999.
- [12] P. N. Norton, *0-Hecke algebras*, *J. Austral. Math. Soc. Ser. A* **27** (1979), no. 3, 337–357.
- [13] R. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge University Press, Cambridge, 1999.
- [14] V. Tewari and S. van Willigenburg, *Modules of the 0-Hecke algebra and quasisymmetric Schur functions*, *Adv. Math.* **285** (2015), 1025–1065.

SEBASTIAN KÖNIG, Leibniz Universität Hannover, Institute of Algebra, Number Theory and Discrete Mathematics, Welfengarten 1, 30167 Hannover, Germany
E-mail : sebastian.koenig@math.uni-hannover.de