



ALGEBRAIC COMBINATORICS

David Anderson & William Fulton

Vexillary signed permutations revisited

Volume 3, issue 5 (2020), p. 1041-1057.

[<http://alco.centre-mersenne.org/item/ALCO_2020__3_5_1041_0>](http://alco.centre-mersenne.org/item/ALCO_2020__3_5_1041_0)

© The journal and the authors, 2020.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is member of the
Centre Mersenne for Open Scientific Publishing
www.centre-mersenne.org



Vexillary signed permutations revisited

David Anderson & William Fulton

ABSTRACT We study the combinatorial properties of vexillary signed permutations, which are signed analogues of the vexillary permutations first considered by Lascoux and Schützenberger. We give several equivalent characterizations of vexillary signed permutations, including descriptions in terms of essential sets and pattern avoidance, and we relate them to the vexillary elements introduced by Billey and Lam.

INTRODUCTION

The class of *vexillary permutations* in S_n , first identified by Lascoux and Schützenberger [17, 18], plays a central role in the combinatorics of the symmetric group and the corresponding geometry of Schubert varieties and degeneracy loci. The name derives from the fact that the Schubert polynomial of a vexillary permutation is equal to a flagged Schur polynomial. This was given a geometric explanation in [11]: vexillary permutations correspond to degeneracy loci defined by simple rank conditions, whose classes are computed by variations of the Kempf–Laksov determinantal formula.

Degeneracy loci of other classical types are indexed by the group W_n of signed permutations. In the course of proving analogous Pfaffian formulas for such loci [2, 3], we constructed *vexillary signed permutations*, starting with the notion of a *triple*. A triple consists of three s -tuples of positive integers, $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$, with $\mathbf{k} = (0 < k_1 < \cdots < k_s)$, $\mathbf{p} = (p_1 \geq \cdots \geq p_s > 0)$, and $\mathbf{q} = (q_1 \geq \cdots \geq q_s > 0)$, satisfying $k_{i+1} - k_i \leq p_i - p_{i+1} + q_i - q_{i+1}$ for $1 \leq i \leq s - 1$. Given such a triple, one constructs a signed permutation $w = w(\tau)$ (see § 2 for the details). By definition, our vexillary signed permutations are the ones arising this way.

Vexillary permutations in S_n have many equivalent characterizations, some of which will be reviewed below; others may be found in [15]. The quickest one is via *pattern avoidance*: a permutation v is vexillary if and only if it avoids the pattern $[2\ 1\ 4\ 3]$, that is, there are no indices $a < b < c < d$ such that $v(b) < v(a) < v(d) < v(c)$. Another is that the Stanley symmetric function of a vexillary permutation is equal to a single Schur function. The latter property was taken as the starting point for Billey and Lam’s extension of “vexillary” to other Lie types: they defined three distinct classes of vexillary elements in types B, C, and D, whose Stanley functions are equal to single Schur P - or Q -functions [6]. Our starting point is the geometric property:

Manuscript received 4th June 2018, revised 25th March 2020, accepted 5th April 2020.

KEYWORDS. Signed permutation, vexillary permutation, degeneracy locus, essential set.

ACKNOWLEDGEMENTS. DA was partially supported by NSF Grant DMS-1502201 and a postdoctoral fellowship from the Instituto Nacional de Matemática Pura e Aplicada (IMPA).

the vexillary signed permutations considered in [2, 3] correspond to degeneracy loci defined by rank conditions of a particularly simple kind, and whose Schubert polynomials can be written as flagged Pfaffians. As often happens, properties that coincide in type A diverge in other types, it turns out that Billey and Lam’s type B vexillary elements are the same as our vexillary signed permutations, which do not depend on type.⁽¹⁾ See Remark 2.7 for more context.

The main goal of this article is to provide alternative characterizations of vexillary signed permutations. In addition to a signed pattern avoidance criterion, we give characterizations of vexillary signed permutations in terms of essential sets of rank conditions, embeddings in symmetric groups, and Stanley symmetric functions (the latter as a consequence of the coincidence with Billey–Lam’s type B vexillary elements).

The definition of a vexillary signed permutation $w(\boldsymbol{\tau})$, given combinatorially in § 2, comes with a geometric explanation in terms of degeneracy loci. Given an odd-rank vector bundle V on a variety X , equipped with a nondegenerate quadratic form and flags of isotropic subbundles $V \supset E_1 \supset E_2 \supset \dots$ and $V \supset F_1 \supset F_2 \supset \dots$, a signed permutation w determines a degeneracy locus $\Omega_w \subseteq X$, defined by imposing certain rank conditions $\dim(E_p \cap F_q) \geq k$. Given a triple $\boldsymbol{\tau} = (\mathbf{k}, \mathbf{p}, \mathbf{q})$, the vexillary signed permutation $w(\boldsymbol{\tau})$ is defined so that the rank conditions for the corresponding degeneracy locus are $\dim(E_{p_i} \cap F_{q_i}) \geq k_i$, for $1 \leq i \leq s$, and so that $w(\boldsymbol{\tau})$ is minimal (in Bruhat order) with this property. The inequalities required on $\mathbf{k}, \mathbf{p}, \mathbf{q}$ guarantee that these rank conditions are feasible: they come from the inclusion of vector spaces

$$(E_{p_{i+1}} \cap F_{q_{i+1}})/(E_{p_i} \cap F_{q_i}) \subseteq E_{p_{i+1}}/E_{p_i} \oplus F_{q_{i+1}}/F_{q_i}.$$

There is an equivalent interpretation for even-rank vector bundles with a symplectic form.

For degeneracy loci corresponding to ordinary permutations, a minimal list of non-redundant rank conditions is determined by the *essential set* of the permutation [11]. The essential set defined in [11] consists of pairs (p, q) appearing in rank conditions $\dim(E_p \cap F_q) \geq k$; the value of k is determined by a rank function associated to the permutation. Here we will modify the definitions slightly, for both ordinary and signed permutations, to include the value of the rank function. Our essential sets, introduced and studied in [1], consist of certain “basic triples” (k, p, q) ; the pairs (p, q) will be called *essential positions*. (Precise definitions are reviewed in § 1.)

One characterization of vexillary signed permutations is that the corresponding essential positions (p_i, q_i) may be ordered so that $p_1 \geq \dots \geq p_s > 0$ and $q_1 \geq \dots \geq q_s > 0$. Another is given via comparison with symmetric groups: the definition of the group of signed permutations leads naturally to an embedding $\iota: W_n \hookrightarrow S_{2n+1}$, and remarkably, a signed permutation w is vexillary if and only if $\iota(w)$ is. (The same statement applies to another natural embedding $\iota': W_n \hookrightarrow S_{2n}$.) We also give a characterization via *signed pattern avoidance*, analogous to the pattern avoidance criterion for S_n ; see § 3 for details.

THEOREM. *Let w be a signed permutation. The following are equivalent:*

- (i) w is vexillary, i.e. it is equal to $w(\boldsymbol{\tau})$ for some triple $\boldsymbol{\tau}$.
- (ii) The essential positions of w can be ordered $(p_1, q_1), \dots, (p_s, q_s)$, so that $p_1 \geq \dots \geq p_s > 0$ and $q_1 \geq \dots \geq q_s > 0$.
- (iii) $\iota(w)$ is vexillary, as a permutation in S_{2n+1} .

⁽¹⁾Since a motivating property of our vexillary signed permutations is the flagged Pfaffian formula for Schubert polynomials, they could be called *Pfaffian-vexillary*, when it is necessary to distinguish them from Billey and Lam’s vexillary elements.

- (iv) $\iota'(w)$ is vexillary, as a permutation in S_{2n} .
- (v) w avoids the nine signed patterns $[2\ 1]$, $[\bar{3}\ 2\ \bar{1}]$, $[\bar{4}\ \bar{1}\ \bar{2}\ 3]$, $[\bar{4}\ 1\ \bar{2}\ 3]$, $[\bar{3}\ \bar{4}\ \bar{1}\ \bar{2}]$, $[\bar{3}\ \bar{4}\ 1\ \bar{2}]$, $[2\ \bar{3}\ 4\ \bar{1}]$, $[2\ \bar{3}\ 4\ \bar{1}]$, and $[3\ \bar{4}\ \bar{1}\ \bar{2}]$.

The theorem is a consequence of Theorem 2.6 combined with Proposition 3.1 and Remark 3.2.

A triple τ determines a strict partition $\lambda(\tau)$ of length s , by setting $\lambda_{k_i} = p_i + q_i - 1$, and filling in the remaining λ_k minimally subject to the strict decreasing requirement; these partitions index the multi-Schur Pfaffians appearing as degeneracy locus formulas in [2]. In § 4, we use “labellings” of their (shifted) Young diagrams to describe how to move between vexillary signed permutations along chains in Bruhat order. We also show that the sum of the parts of $\lambda(\tau)$ computes the length of $w(\tau)$ (Corollary 4.5).

The pattern avoidance criterion (v) coincides with that of Billey and Lam in type B, and as a consequence, we can add another equivalent condition to the list in the above theorem: writing H_w for the Stanley symmetric function of type B (see, e.g. [5] or [10, (6.3)]), we have

- (vi) H_w is equal to a Schur P -function P_λ .

(In fact, Corollary 5.2 says $H_{w(\tau)} = P_{\lambda(\tau)}$.) The equivalence of (vi) with (v) is the content of [6, Theorem 14]. Billey and Lam also give pattern avoidance criteria for determining when other variants of Stanley symmetric functions are equal to P - and Q -functions, but our vexillary signed permutations form a strict subset of the those defined in [6] for types C and D.

Is there a geometric property enjoyed by Billey–Lam’s vexillary elements of type C or D? In [3], we find “flagged theta-polynomial” formulas for a larger class of signed permutations constructed from generalized triples (so they might be called *theta-vexillary*), but they are not the same as any of Billey–Lam’s vexillary elements.⁽²⁾

NOTATION. We refer to [7, § 8.1] for details on signed permutations, and review the notation and conventions here. We will consider permutations of (positive and negative) integers

$$\dots, \bar{n}, \dots, \bar{2}, \bar{1}, 0, 1, 2, \dots, n, \dots,$$

using the bar to denote a negative sign, and we take the natural order on them, as above. All permutations are finite, in the sense that $v(m) = m$ whenever $|m|$ is sufficiently large. We generally write permutations in S_{2n+1} using one-line notation, by listing the values $v(\bar{n})\ v(\bar{n}-1)\ \dots\ v(n)$.

A *signed permutation* is a permutation w with the property that for each i , $w(\bar{i}) = \overline{w(i)}$. A signed permutation is in W_n if $w(m) = m$ for all $m > n$; this is a group isomorphic to the hyperoctahedral group, the Weyl group of types B_n and C_n . When writing signed permutations in one-line notation, we only list the values on positive integers: $w \in W_n$ is represented as $w(1)\ w(2)\ \dots\ w(n)$. For example, $w = \bar{2}\ 1\ \bar{3}$ is a signed permutation in W_3 , and $w(\bar{3}) = 3$ since $w(3) = \bar{3}$. The *simple transpositions* s_0, \dots, s_{n-1} generate W_n , where for $i > 0$, right-multiplication by s_i exchanges entries in positions i and $i + 1$, and right-multiplication by s_0 replaces $w(1)$ with $\overline{w(1)}$. The

⁽²⁾Theta-vexillary signed permutations are rather more complicated to define than the vexillary ones we consider here. They include all Grassmannian signed permutations; our vexillary signed permutations include all 0-Grassmannian signed permutations, that is, ones coming from Grassmannians of maximal isotropic subspaces, but only some p -Grassmannian ones, for $p > 0$.

Answering a question we raised when an early version of this paper circulated, J. Lambert has carried out a detailed combinatorial study of these theta-vexillary signed permutations [16]. His results include a pattern-avoidance criterion which distinguishes these elements from previously studied ones.

length of a signed permutation w is the least number $\ell = \ell(w)$ such that $w = s_{i_1} \cdots s_{i_\ell}$. The longest element in W_n , denoted $w_\circ^{(n)}$, is $\bar{1} \bar{2} \cdots \bar{n}$, and has length n^2 .

The definition of W_n presents it as embedded in the symmetric group S_{2n+1} , considering the latter as the group of all permutations of the integers $\bar{n}, \dots, 0, \dots, n$. This is the odd embedding, and we write $\iota: W_n \hookrightarrow S_{2n+1}$ for emphasis when a signed permutation is considered as a full permutation. Specifically, ι sends $w = w(1) w(2) \cdots w(n)$ to the permutation

$$\overline{w(n)} \cdots \overline{w(2)} \overline{w(1)} 0 w(1) w(2) \cdots w(n)$$

in S_{2n+1} . Occasionally, we will refer to the even embedding $\iota': W_n \hookrightarrow S_{2n}$, defined similarly to ι by considering the even symmetric group as permutations of $\{\pm 1, \dots, \pm n\}$ and omitting the value $w(0) = 0$.

A permutation v has a descent at position i if $v(i) > v(i + 1)$; here i may be any integer. Equivalently, $\ell(vs_i) = \ell(v) - 1$. The same definition applies to signed permutations w , but we only consider descents at positions $i \geq 0$, following the convention of recording the values of w only on positive integers. A descent at 0 simply means that $w(1)$ is negative. Equivalently, $\ell(ws_i) = \ell(w) - 1$. For example, $w = \bar{2} 1 \bar{3}$ has descents at 0 and 2, while $\iota(w) = 3 \bar{1} 2 0 \bar{2} 1 \bar{3}$ has descents at $-3, -1, 0$, and 2.

1. DIAGRAMS AND ESSENTIAL SETS

A permutation $v \in S_{2n+1}$ can be represented in a $(2n + 1) \times (2n + 1)$ array of boxes, with rows and columns indexed by $\{\bar{n}, \dots, 0, \dots, n\}$, by placing a dot in position $(v(i), i)$ for $\bar{n} \leq i \leq n$; we refer to this as the permutation matrix of v . The diagram of v is the collection of boxes that remain after crossing out those weakly south or east of a dot. The rank function of v is defined as

$$(1) \quad r_v(p, q) = \#\{i \leq \bar{p} \mid v(i) \geq q\},$$

for $\bar{n} \leq p, q \leq n$. This is equal to the number of dots strictly south and weakly west of the box $(q - 1, \bar{p})$ in the permutation matrix of v .

The $(2n + 1)^2$ numbers $r_v(p, q)$ determine v , but in fact much less information is required to specify a permutation. A minimal list of rank conditions determining v was given in [11], by restricting attention to the southeast corners of the diagram.⁽³⁾ More precisely, following the conventions and terminology of [1] (which differ slightly from those of [11]), a pair (p, q) is an essential position if the box $(q - 1, \bar{p})$ is a southeast corner of the diagram of v . The essential set $\mathcal{Ess}(v)$ is the set of (k, p, q) such that (p, q) is an essential position and $k = r_v(p, q)$.

In formulas, a box (a, b) is a SE corner of the diagram of v if and only if

$$(2) \quad v(b) > a \geq v(b + 1) \quad \text{and}$$

$$(2') \quad v^{-1}(a) > b \geq v^{-1}(a + 1),$$

so the essential positions are those (p, q) such that $(q - 1, \bar{p})$ satisfies Equations (2) and (2'). This characterizes $\mathcal{Ess}(v)$ in terms of the descents of v and v^{-1} , and will be useful later.

Analogous diagrams and essential sets for signed permutations were described in [1]. We review the definitions and basic facts briefly here, since essential sets play a role in characterizing vexillary signed permutations.

⁽³⁾By analyzing the possibilities for boxes occurring as southeast corners of the diagram of a permutation, Eriksson and Linusson showed how to reconstruct v from a subset of the essential set [9]; however, the essential set is minimal in the sense of [11, Lemma 3.10].

For a signed permutation $w \in W_n$, the following simple lemma says that the essential positions of the corresponding permutation $\iota(w) \in S_{2n+1}$ are “symmetric about the origin” (see Figure 1).

LEMMA 1.1 ([1, Lemma 1.1]). *For $w \in W_n$, the essential set of $\iota(w) \in S_{2n+1}$ possesses the following symmetry: (k, p, q) is in $\mathcal{Ess}(\iota(w))$ if and only if $(k+p+q-1, \bar{p}+1, \bar{q}+1)$ is in $\mathcal{Ess}(\iota(w))$.*




FIGURE 1. Diagram and essential set for $v = \iota(\bar{2} \ 3 \ \bar{1})$, with circled corners illustrating the symmetry of Lemma 1.1.

This implies that half of $\mathcal{Ess}(\iota(w))$ suffices to determine the signed permutation w ; we will consider those corners appearing in the first n columns. In general, only a subset of these corners is needed, as shown in [1].

As with ordinary permutations, the essential set of a signed permutation is defined in terms of its diagram. The *permutation matrix* of $w \in W_n$ is a $(2n + 1) \times n$ array of boxes, with rows labelled $\bar{n}, \dots, 0, \dots, n$ and columns labelled $\bar{n}, \dots, \bar{1}$. A dot is placed in the boxes $(\overline{w(i)}, \bar{i})$ for $1 \leq i \leq n$. Additionally, for each dot, the box in the same column but opposite row is marked with an \times , as is each box to the right of this one. (That is, an \times is placed in each box (a, b) such that $a = \overline{w(i)}$ for some $i \leq b$.) The *extended diagram* of w is the set \tilde{D}_w of boxes that remain after crossing out those south or east of a dot in the permutation matrix; the *diagram* D_w is the subset of \tilde{D}_w not marked by an \times . The placement of the \times 's is closely related to the parametrization of Schubert cells described in [12], and the boxes of D_w are in natural bijection with the inversions of w . (The *inversions* of a signed permutation w are pairs $(i < j)$ such that $w(i) > w(j)$, together with pairs $(i \leq j)$ such that $w(i) + w(j) < 0$.) In particular, the number of boxes in D_w is equal to the length of w . See Figure 2 for an illustration.

The rank function of w is defined as

$$(3) \quad r_w(p, q) = \#\{i \geq p \mid w(i) \leq \bar{q}\}.$$

Since $w(\bar{i}) = \overline{w(i)}$, this is equivalent to

$$(4) \quad r_w(p, q) = \#\{i \leq \bar{p} \mid w(i) \geq q\}.$$

The *essential set* of a signed permutation w is the set of (k, p, q) such that $(q - 1, \bar{p})$ is a SE corner of the extended diagram \tilde{D}_w and $k = r_w(p, q)$, with two exceptions. First, if $p = 1$ and $q < 0$ (i.e. if the corner appears in the rightmost column and above the middle row in the permutation matrix), then (k, p, q) is not in $\mathcal{Ess}(w)$. Second, when $p > 1$ and $q > 0$, (k, p, q) is not in $\mathcal{Ess}(w)$ if there is another SE corner in box (\bar{q}, \bar{p}) , and $k = r_w(p, q) = r_w(p, \bar{q} + 1) - q + 1$.

The exceptions are easy to understand in the context of linear algebra; see [1] for more explanation. The first one comes from Lemma 1.1: such corners are artifacts of




FIGURE 2. Diagram and essential set of a signed permutation. (This is also the vexillary signed permutation for the triple $\tau = (1\ 3\ 4\ 5\ 8, 9\ 9\ 6\ 4\ 3, 12\ 9\ 8\ 8\ 5)$.)

restricting the permutation matrix to n columns, and they do not appear as corners of the diagram of $\iota(w)$. The second, more complicated exception never applies to the vexillary signed permutations to be defined in the next section. As we will see in Lemma 2.5, when w is vexillary, no SE corners of \tilde{D}_w lie above the middle row, except possibly in the rightmost column.

Just as with ordinary permutations, a signed permutation is determined by its essential set: w is the minimal element (in Bruhat order on W_n) such that $r_w(p, q) \geq k$ for all $(k, p, q) \in \mathcal{E}_{ss}(w)$. Furthermore, $\mathcal{E}_{ss}(w)$ is the smallest set with this property, choosing any $(k_0, p_0, q_0) \in \mathcal{E}_{ss}(w)$, there exists a $w' \neq w$ such that $r_{w'}(p_0, q_0) < k_0$ and $r_{w'}(p, q) \geq k$ for all $(k, p, q) \in \mathcal{E}_{ss}(w) \setminus \{(k_0, p_0, q_0)\}$. (This is the content of [1, Theorem 2.3].)

2. TRIPLES AND VEXILLARY SIGNED PERMUTATIONS

As defined in the introduction, a *triple* consists of three s -tuples of positive integers $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$, with

$$\begin{aligned} \mathbf{k} &= (0 < k_1 < \dots < k_s), \\ \mathbf{p} &= (p_1 \geq \dots \geq p_s > 0), \\ \mathbf{q} &= (q_1 \geq \dots \geq q_s > 0), \end{aligned}$$

satisfying

$$(*) \quad p_i - p_{i+1} + q_i - q_{i+1} \geq k_{i+1} - k_i$$

for $1 \leq i \leq s - 1$. The triple is *essential* if the inequalities (*) are strict. (These inequalities ensure that the rank conditions $\dim(E_{p_i} \cap F_{q_i}) \geq k_i$ are feasible, and strict inequalities ensure they are independent.) Every triple reduces to a unique essential triple, by successively removing each (k_i, p_i, q_i) such that equality holds in (*). Two triples are *equivalent* if they reduce to the same essential triple.

Given a triple, one forms a signed permutation $w(\tau)$ as follows.

- (1) Starting in the p_1 position, place k_1 consecutive negative entries, in increasing order, ending with $\overline{q_1}$. Mark these numbers as “used”.
- (i) For $1 < i \leq s$, starting in the p_i position, or the next available position to the right, fill the next available $k_i - k_{i-1}$ positions with negative entries chosen consecutively from the unused numbers, ending with at most $\overline{q_i}$. (Note: the sequence may have to “jump” over previously placed sequences.)⁽⁴⁾
- (s+1) Fill the remaining available positions with the unused positive numbers, in increasing order.

DEFINITION 2.1. A signed permutation $w \in W_n$ is *vexillary* if $w = w(\tau)$ for some triple $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$. By convention, the “empty” triple is a triple, so the identity element is vexillary.

A triple also defines a strict partition $\lambda(\tau)$, by setting $\lambda_{k_i} = p_i + q_i - 1$, and defining the other parts minimally so that the result is a strictly decreasing sequence of integers. (In formulas, $\lambda_k = p_i + q_i - 1 + k_i - k$ if $k_{i-1} < k \leq k_i$.)

For example, with $\tau = (1\ 3\ 4\ 5\ 8, 9\ 9\ 6\ 4\ 3, 12\ 9\ 8\ 8\ 5)$, the six steps in forming $w(\tau)$ produce

$$\begin{aligned} w &= \dots\dots\dots \overline{12}, \\ w &= \dots\dots\dots \overline{12}\ \overline{10}\ \overline{9}, \\ w &= \dots\dots\ \overline{8}\ \dots\ \overline{12}\ \overline{10}\ \overline{9}, \\ w &= \dots\ \overline{11}\ \cdot\ \overline{8}\ \dots\ \overline{12}\ \overline{10}\ \overline{9}, \\ w &= \dots\ \overline{7}\ \overline{11}\ \overline{6}\ \overline{8}\ \overline{5}\ \cdot\ \overline{12}\ \overline{10}\ \overline{9}, \\ w &= \mathbf{1}\ \mathbf{2}\ \overline{7}\ \overline{11}\ \overline{6}\ \overline{8}\ \overline{5}\ \mathbf{3}\ \overline{12}\ \overline{10}\ \overline{9}\ \mathbf{4}. \end{aligned}$$

The corresponding partition is $\lambda = (20, 18, 17, 13, 11, 9, 8, 7)$.

LEMMA 2.2. Equivalent triples produce the same signed permutation and the same strict partition.

Proof. Given a triple τ , suppose equality holds for some i in (*), so $p_i - p_{i+1} + q_i - q_{i+1} = k_{i+1} - k_i$. Let τ' be the triple obtained from τ by deleting (k_i, p_i, q_i) . So the i th component of τ' is $(k_{i+1}, p_{i+1}, q_{i+1})$. We will compare the algorithms for forming $w(\tau)$ and $w(\tau')$, step by step.

For $j < i$, Steps (j) are evidently the same. We will show that Steps (i) and (i+1) for $w(\tau)$ have the same effect as Step (i) for $w(\tau')$.

First consider which entries are to be placed. For $w(\tau)$, we take the $k_i - k_{i-1}$ largest unused entries less than or equal to $\overline{q_i}$, and then the $k_{i+1} - k_i$ largest unused entries

⁽⁴⁾It is useful to break each step in the construction into sub-steps. In Step (i), first pick out the $k_i - k_{i-1}$ largest unused entries less than or equal to $\overline{q_i}$ and place them in a “bin”. Second, go to position p_i : If this position is available, take the smallest entry from the bin, place it here, and move to position $p_i + 1$; if position p_i is unavailable, just move to position $p_i + 1$. Repeat the second sub-step, starting at position $p_i + 1$. Finish when the bin is empty.

less than or equal to $\overline{q_{i+1}}$. Taken together, these are same as those to be placed for $w(\tau')$, namely the $k_{i+1} - k_{i-1}$ largest unused entries which are less than or equal to $\overline{q_{i+1}}$. Let us collect these entries in a bin, sorted in increasing order. Schematically, the bin looks as shown below.

$$\underbrace{* \cdots *}_{p_i - p_{i+1}} \quad \underbrace{* \cdots *}_{k_i - k_{i-1}} \quad \underbrace{* \cdots *}_{q_i - q_{i+1}}$$

If $q_{i+1} < q_i$, then the entries $\overline{q_i + 1}, \dots, \overline{q_{i+1}}$ are the $q_i - q_{i+1}$ largest entries in the bin. This means that the middle group of $k_i - k_{i-1}$ entries are the largest ones which are less than or equal to $\overline{q_i}$, that is, the ones to be placed in Step (i) for $w(\tau)$.

Now we turn to the placement of these entries. For $w(\tau)$, in Step (i) we pick out the $k_i - k_{i-1}$ entries from the middle group in the above diagram of the bin, go to position p_i , and place these entries in the next available spots. Then, in Step (i+1), we go to position p_{i+1} and place the remaining entries in the next available spots. The result is the same as starting at position p_{i+1} and placing all the entries at once, which is the recipe for Step (i) of $w(\tau')$.

For $j > i$, Step (j+1) for $w(\tau)$ is the same as Step (j) for $w(\tau')$, and it follows that the signed permutations are identical. The claim about strict partitions is easily verified from the definition. \square

In the rest of the article, we study some of the properties of this construction. Since equivalent triples produce the same signed permutation and the same strict partition, we will generally assume triples are essential.

LEMMA 2.3. *Let $w = w(\tau)$, for an essential triple $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$. The descents of w are at the positions $p_i - 1$. In fact, for each i , we have $w(p_i - 1) > \overline{q_i} \geq w(p_i)$, and there are no other descents.*

Proof. In Step (1), no descents are created, unless $p_1 = 1$, in which case the permutation has a single descent at 0.

Now for $1 < i \leq s$, consider the situation before Step (i) in constructing $w(\tau)$. Assume inductively that for $j < i$, there is a descent at position $p_j - 1$ whenever this position has been filled, satisfying $w(p_j - 1) > \overline{q_j} \geq w(p_j)$, and there are no other descents.

In carrying out Step (i), we place negative entries in consecutive vacant positions, from left to right, starting at position p_i (or the next vacant position to the right, if $p_i = p_{i-1}$). We consider “sub-steps” of Step (i), where we are placing an entry at position $p \geq p_i$, and distinguish three cases.

First, suppose we are at position p , with $p < p_{i-1} - 1$. In this case, the previous entry placed in Step (i) (if any) was placed at position $p - 1$, so we did not create a descent at $p - 1$. Position $p + 1$ is still vacant, so no new descents are created.

To illustrate, take $\tau = (1\ 3\ 4\ 5\ 8, 9\ 9\ 6\ 4\ 3, 12\ 9\ 8\ 8\ 5)$, as in the example above. In Step (3), we place $\overline{8}$ without creating a descent:

$$w = \dots \overline{8} \dots \overline{12}\ \overline{10}\ \overline{9}.$$

Next, suppose we are at position $p = p_{i-1} - 1$. This means $p_{i-1} - p_i \leq k_i - k_{i-1}$, so let $\beta = (k_i - k_{i-1}) - (p_{i-1} - p_i)$ be the number of entries remaining to be placed in Step (i), after placing the current one. The inequality (*) in the triple condition implies $q_i + \beta < q_{i-1}$, which in turn means that the entries used in previous steps are all strictly less than $\overline{q_i + \beta}$; therefore the $\beta + 1$ entries $\overline{q_i + \beta}, \overline{q_i + \beta - 1}, \dots, \overline{q_i}$ are all available to be placed. It follows that $w(p_{i-1} - 1) = \overline{q_i + \beta}$, which creates a descent satisfying the claim of the lemma. Again, we did not create a descent at position $p - 1$, for the same reason as in the previous case.

Continuing the running example, the first entry placed in Step (5) is the $\bar{7}$, creating a descent at position 3:

$$w = \cdot \cdot \bar{7} \bar{11} \cdot \bar{8} \cdot \cdot \bar{12} \bar{10} \bar{9}.$$

(In this case, we had $q_5 = 5$ and $\beta = 2$.)

Finally, suppose we are at position $p = p_{i-1}$. Set $\beta = (k_i - k_{i-1}) - (p_{i-1} - p_i) - 1$, so $\beta \geq 0$; this is the number of entries to be placed after the current one. Now the essential triple condition implies $q_i + \beta + 1 < q_{i-1}$. The entries that will be placed are therefore $-q_i - \beta, -q_i - \beta + 1, \dots, -q_i$. These are all greater than the ones that already occupy positions to the right of p_{i-1} . Whenever such an entry is placed in a vacant position to the right of a filled position, then, it does not create a descent at that filled position. Whenever it is placed to the left of a filled position, it does create a descent at the position it is placed. The latter can only happen at positions $p_j - 1$, for $j < i - 1$.

In Step (5) of our example, after placing the $\bar{7}$ in position $p_5 = p_4 - 1$, we come to position p_4 . The entries to be placed are $\bar{6}, \bar{5}$. In placing the $\bar{6}$ in the next vacant position, position 5, we do not create a descent at the filled position to its left, but we do create a descent at position 5, since position 6 is already filled:

$$w = \cdot \cdot \bar{7} \bar{11} \bar{6} \bar{8} \bar{5} \cdot \bar{12} \bar{10} \bar{9}.$$

In placing the $\bar{5}$, there is no descent created, since the position to its right is vacant.

At Step (s+1), the same reasoning as in the last case considered shows that descents are created at those $p_j - 1$ which are still vacant, and nowhere else. \square

Given a triple $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$, the *dual triple* is $\tau^* = (\mathbf{k}, \mathbf{q}, \mathbf{p})$.

LEMMA 2.4. We have $w(\tau^*) = w(\tau)^{-1}$.

Proof. Suppose $w \in W_n$, and consider $\iota(w)$ as a bijective map from the set $\{\bar{n}, \dots, 0, \dots, n\}$ to itself. To determine $\iota(w)$, it is enough to know its values on any set of integers whose absolute values are $\{1, \dots, n\}$. The construction of w translates as follows. Let $a(1) = \{p_1, p_1 + 1, \dots, p_1 + k_1 - 1\}$, let $a(2)$ be the set of $k_2 - k_1$ consecutive integers in $\{1, \dots, n\} \setminus a(1)$ starting from the p_2 th element, and define sets $a(3), \dots, a(s)$ similarly. Let $a(s+1)$ be the complement of $a(1), \dots, a(s)$, so the sets $a(1), \dots, a(s+1)$ partition $\{1, \dots, n\}$. Define $b(1), \dots, b(s+1)$ in the same way, using \mathbf{q} in place of \mathbf{p} . Also write $\bar{a}(i)$ and $\bar{b}(i)$ for the corresponding sets of negative integers. Now $\iota(w)$ maps $\bar{a}(i)$ to $\bar{b}(i)$ for $1 \leq i \leq s$, and it maps $\bar{a}(s+1)$ to $b(s+1)$. It follows that $\iota(w)^{-1}$ maps $\bar{b}(i)$ to $\bar{a}(i)$, or equivalently, $b(i)$ to $\bar{a}(i)$, and maps $b(s+1)$ to $\bar{a}(s+1)$. In other words, the inverse is obtained by switching the roles of a and b , so the lemma is proved. \square

LEMMA 2.5.

- (1) Let $w = w(\tau)$ be a vexillary signed permutation, for an essential triple $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$. Then the SE corners of the diagram of $\iota(w)$ are the boxes $(q_i - 1, \bar{p}_i)$, together with their reflections $(\bar{q}_i, p_i - 1)$. (In particular, no box (a, b) with $a, b < 0$ occurs as a SE corner.)
- (2) k_i is the number of dots strictly south and weakly west of the i th SE corner in the diagram (in position $(q_i - 1, \bar{p}_i)$).

Proof. By Lemma 1.1, it suffices to show that the first s corners of the diagram of $\iota(w)$ (ordered SW to NE) are as claimed. For $p > 0$, a signed permutation w has a descent at position $p - 1$ iff $\iota(w)$ has descents at positions $p - 1$ and \bar{p} . By Lemma 2.3, the descents of $\iota(w)$ are at positions $p_i - 1$ and \bar{p}_i , and the inequalities (2)

$$\iota(w)(\bar{p}_i) > q_i - 1 \geq \iota(w)(\overline{p_i - 1})$$

follow from

$$w(p_i - 1) \geq q_i - 1 > w(p_i),$$

using Lemma 2.3 again.

A similar argument establishes the inequalities (2'):

$$\iota(w)^{-1}(q_i - 1) > \bar{p}_i \geq \iota(w)^{-1}(q_i)$$

(Swap \mathbf{p} and \mathbf{q} , and apply Lemma 2.3 to $w(\mathbf{k}, \mathbf{q}, \mathbf{p})$, using Lemma 2.4 to see $w(\mathbf{k}, \mathbf{q}, \mathbf{p}) = w^{-1}$.)

Part (2) is easy from the construction: At the end of Step (i), all the entries to the right of p_i are at most \bar{q}_i , and there are k_i of them. After Step (i), any entry placed to the right of p_i is greater than \bar{q}_i . \square

THEOREM 2.6. *For $w \in W_n$, the following are equivalent:*

- (i) *The signed permutation w is vexillary.*
- (ii) *The essential positions (p_i, q_i) of w can be ordered so that $p_1 \geq \dots \geq p_s > 0$ and $q_1 \geq \dots \geq q_s > 0$.*
- (iii) *The permutation $\iota(w)$ is vexillary (as an element of S_{2n+1}).*

Proof. We will show (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i). The implication (i) \Rightarrow (ii) is immediate from Lemma 2.5, and (ii) \Rightarrow (iii) follows from Lemma 1.1. It remains to show (iii) \Rightarrow (i).

Suppose $\iota(w)$ is vexillary, and recall that this is equivalent to requiring that the SE corners of its diagram proceed from southwest to northeast [11, Remark 9.17]. Take those SE corners (a, b) such that $b < 0$; Lemma 1.1 implies $a \geq 0$. Let $s = \lceil \frac{\#\text{Ess}(\iota(w))}{2} \rceil$ be the number of such boxes. Reading from SW to NE, label them (a_i, b_i) , with $1 \leq i \leq s$. Set $p_i = -b_i$ and $q_i = a_i + 1$, and let k_i be the number of dots strictly south and weakly west of (a_i, b_i) .

We claim that $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$ is an essential triple. Indeed, looking at the extended diagram \tilde{D}_w , the fact that the boxes $(q_i - 1, \bar{p}_i)$ are SE corners implies the inequalities (*). It follows that w is equal to $w(\tau)$, since a permutation is determined by its essential set [11, Lemma 3.10(b)], [1, Theorem 2.3]. \square

REMARK 2.7. Our definition of vexillary signed permutation does not refer to Lie type, and in our setting there is no difference between types B, C, and D. (As noted in the introduction, this stands in contrast to Billey–Lam vexillary elements: ours coincide with their type B elements.) For applications to degeneracy loci, no further conditions are needed, even in type D, instead noting that there are two families of maximal isotropic subspaces for type D loci. On the other hand, the vexillary signed permutations lying in the Weyl group of type D, the index-2 subgroup of W_n having an even number of negative signs, are just those corresponding to triples with k_s even.

3. PATTERN AVOIDANCE

Given a permutation π in S_m , a permutation v contains the pattern π if, when written in one-line notation, there is an m -element subsequence of v which is in the same relative order as π ; otherwise v avoids π . There is a similar notion for signed permutations and signed patterns. Given a signed pattern $\pi = \pi(1) \pi(2) \dots \pi(m)$ in W_m , a signed permutation w contains π if there is a subsequence $w(i_1) \dots w(i_m)$ such that the signs of $w(i_j)$ and $\pi(j)$ are the same for all j , and also the absolute values of the subsequence are in the same relative order as the absolute values of π . Otherwise w avoids π . (See, e.g. [6, Definition 6].)

For example, $\bar{5} 1 3 \bar{2} 4$ contains the pattern $[\bar{3} 2 \bar{1}]$, as the subsequence $\bar{5} 3 \bar{2}$, but $\bar{5} 1 2 \bar{3} \bar{4}$ avoids $[\bar{3} 2 \bar{1}]$.

PROPOSITION 3.1. A signed permutation w is vexillary if and only if w avoids the signed patterns $[2\ 1]$, $[\overline{3}\ 2\ \overline{1}]$, $[\overline{4}\ \overline{1}\ \overline{2}\ 3]$, $[\overline{4}\ 1\ \overline{2}\ 3]$, $[\overline{3}\ \overline{4}\ \overline{1}\ \overline{2}]$, $[\overline{3}\ \overline{4}\ 1\ \overline{2}]$, $[2\ \overline{3}\ 4\ \overline{1}]$, $[2\ \overline{3}\ 4\ \overline{1}]$, and $[3\ \overline{4}\ \overline{1}\ \overline{2}]$.

Proof. We will use the characterization of Theorem 2.6, and show that a signed permutation w avoids these nine patterns if and only if $\iota(w)$ is vexillary in S_{2n+1} . Recall that a permutation is vexillary if and only if it avoids the pattern $[2\ 1\ 4\ 3]$. When embedded by ι , the nine signed patterns listed in (ii) all contain $[2\ 1\ 4\ 3]$, so if $\iota(w)$ avoids $[2\ 1\ 4\ 3]$, then w must avoid these signed patterns; this proves the “ \Rightarrow ” direction.

For the “ \Leftarrow ” implication, we claim that if $\iota(w)$ contains the pattern $[2\ 1\ 4\ 3]$, then w contains one of the nine signed patterns listed in the Proposition. To check this, we break the ways $[2\ 1\ 4\ 3]$ can appear in $\iota(w)$ into cases, and observe that in each case, one of the listed patterns appears.

For keeping track of the cases, we introduce some temporary notation. An instance of the pattern $[2\ 1\ 4\ 3]$ is witnessed by $\iota(w)(b) < \iota(w)(a) < \iota(w)(d) < \iota(w)(c)$ for some $a < b < c < d$. We record which of a, b, c, d are negative by placing a vertical bar in the word $[2\ 1\ 4\ 3]$, and which of the corresponding values of $\iota(w)$ are negative by placing underlines. For example, if $w = 2\ \overline{3}\ \overline{5}\ 4\ \overline{1}$, so

$$\iota(w) = 1\ \overline{4}\ 5\ 3\ \overline{2}\ 0\ 2\ \overline{3}\ \overline{5}\ 4\ \overline{1},$$

then an instance of $[2\ 1\ 4\ 3]$ occurs as $[2|\underline{1}\ 4\ \underline{3}]$ in the subsequence $\overline{4}\ \overline{5}\ 4\ \overline{1}$ (among others), and another occurs as $[|\underline{2}\ \underline{1}\ 4\ \underline{3}]$ in the subsequence $\overline{3}\ \overline{5}\ 4\ \overline{1}$.

We now check all cases, organized by position of the vertical bar and underlines.

Any occurrence of $[|\underline{2}\ 1\ 4\ \underline{3}]$, $[|\underline{2}\ \underline{1}\ 4\ \underline{3}]$, or $[|\underline{2}\ \underline{1}\ 4\ \underline{3}]$ contains the signed pattern $[2\ 1]$. Suppose $\alpha < \beta < \gamma$, so an occurrence of $[|\underline{2}\ \underline{1}\ 4\ \underline{3}]$ is a subsequence $\overline{\beta}\ \overline{\gamma}\ x\ \overline{\alpha}$. This is the signed pattern $[\overline{2}\ \overline{3}\ 4\ \overline{1}]$ if $x > \gamma$, it is $[\overline{3}\ \overline{4}\ 1\ \overline{2}]$ if $x < \alpha$, and it contains $[\overline{3}\ 2\ \overline{1}]$ if $\alpha < x < \gamma$. Finally, an occurrence of $[|\underline{2}\ \underline{1}\ 4\ \underline{3}]$ is the signed pattern $[\overline{3}\ \overline{4}\ \overline{1}\ \overline{2}]$.

The cases $[2|\underline{1}\ 4\ \underline{3}]$, $[2|\underline{1}\ 4\ \underline{3}]$, and $[2|\underline{1}\ 4\ \underline{3}]$ all contain the signed pattern $[2\ 1]$, as before. With $a < b < c$, the case $[2|\underline{1}\ 4\ \underline{3}]$ depends on the position of the “2”. A subsequence $\beta\ \overline{\gamma}\ x\ \overline{\alpha}$ is the signed pattern $[2\ \overline{3}\ 4\ \overline{1}]$ if $x > \gamma$; it is $[3\ \overline{4}\ 1\ \overline{2}]$ if $x < \alpha$, and it contains $[\overline{3}\ 2\ \overline{1}]$ if $\alpha < x < \gamma$. The subsequences $\overline{\gamma}\ \beta\ x\ \overline{\alpha}$ and $\overline{\gamma}\ x\ \beta\ \overline{\alpha}$ contain $[\overline{3}\ 2\ \overline{1}]$. The subsequence $\overline{\gamma}\ x\ \overline{\alpha}\ \beta$ is the signed pattern $[\overline{4}\ 1\ \overline{2}\ 3]$ if $x < \alpha$, it contains $[\overline{3}\ 2\ \overline{1}]$ if $\alpha < x < \beta$, and it contains $[2\ 1]$ if $x > \beta$. Similarly, depending on the position of the “2”, the case $[2|\underline{1}\ 4\ \underline{3}]$ contains $[3\ \overline{4}\ \overline{1}\ \overline{2}]$, $[\overline{3}\ 2\ \overline{1}]$, or $[\overline{4}\ \overline{1}\ \overline{2}\ 3]$.

All five cases $[2\ 1|\underline{4}\ \underline{3}]$, $[2\ \underline{1}|\underline{4}\ \underline{3}]$, $[\underline{2}\ \underline{1}|\underline{4}\ \underline{3}]$, $[\underline{2}\ \underline{1}|\underline{4}\ \underline{3}]$, and $[\underline{2}\ \underline{1}|\underline{4}\ \underline{3}]$ contain the signed pattern $[2\ 1]$.

The remaining cases are symmetric to the ones checked above. □

REMARK 3.2. An identical analysis shows that w avoids the same nine signed patterns if and only if $\iota'(w) \in S_{2n}$ avoids $[2\ 1\ 4\ 3]$; therefore w is vexillary if and only if $\iota'(w)$ is vexillary.

The pattern avoidance criterion for ordinary permutations allowed J. West to enumerate the vexillary permutations in S_n [19]. His proof uses a bijection between permutations avoiding $[2\ 1\ 4\ 3]$ (vexillary permutations) and those that avoid $[4\ 3\ 2\ 1]$. One might expect a similar enumeration of vexillary signed permutations. In fact, for the even embedding $\iota': W_n \hookrightarrow S_{2n}$ (which omits “ $w(0) = 0$ ”), Egge showed that the number of $w \in W_n$ such that $\iota'(w)$ avoids $[4\ 3\ 2\ 1]$ is equal to

$$V_n := \sum_{k=0}^n \binom{n}{k}^2 C_k,$$

where $C_k = \frac{1}{k+1} \binom{2k}{k}$ is the k th Catalan number [8]. Together with computer verification up to $n = 7$, this suggests the conjecture that V_n is also the number of vexillary

signed permutations in W_n . West's bijection does not preserve the subgroup of signed permutations, though, so the proof is not immediately clear⁽⁵⁾.

4. LABELLED YOUNG DIAGRAMS

There is an alternative way to encode vexillary signed permutations, useful for determining when right-multiplication by a simple reflection takes one vexillary signed permutation to another. Identifying a strict partition λ with its shifted Young diagram, a *labelled Young diagram* of shape λ is an assignment of integers to the SE corners, weakly decreasing from top to bottom, such that the labels m_1, \dots, m_s in rows k_1, \dots, k_s satisfy $0 \leq m_i < \lambda_{k_i}$ and $m_i - m_{i+1} \leq \lambda_{k_i} - \lambda_{k_{i+1}}$. (Note that $\lambda_{k_i} - \lambda_{k_{i+1}} + 1$ is the number of boxes in the rim-hook connecting the corners labelled m_i and m_{i+1} .)

Labelled Young diagrams are in bijection with essential triples: given $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$, the shifted Young diagram for $\lambda(\tau)$ has SE corners in rows k_1, \dots, k_s , and one forms a labelled Young diagram of this shape by placing integers $m_i = p_i - 1$ in these corners. Examples are shown in Figure 3.




FIGURE 3. Labelled Young diagrams associated to $\tau = (2\ 3, 2\ 2, 3\ 1)$ and $\tau^* = (2\ 3, 3\ 1, 2\ 2)$. The corresponding vexillary signed permutations are $2\ \bar{4}\ \bar{3}\ \bar{1}$ and $\bar{4}\ 1\ \bar{3}\ \bar{2}$, respectively.

Let $w = w(\tau)$, and let Y be the corresponding labelled Young diagram of shape $\lambda(\tau)$. By Lemma 2.3, the descents of w are the corner labels of Y , i.e. $\ell(ws_m) = \ell(w) - 1$ if and only if m is a label.

A corner label m in Y is *removable* if

- it appears in Y exactly once, and
- when $m = 0$, the row in which it appears contains a single box.

Given a removable label m , one can remove the box containing it to form a new labelled Young diagram $Y \setminus m$, whose corners are labelled according to the following four rules.

- (i) If a corner of $Y \setminus m$ is also a corner of Y , its label is the same.
- (ii) If removing m produces a new corner one box to the left, label that corner $m - 1$.
- (iii) If removing m produces a new corner one box above, label that corner $m + 1$.
- (iv) If removing m produces two new corners, label the one to the left $m - 1$ and the one above $m + 1$.

These are the only possibilities for removing a corner from the Young diagram. Examples are shown in Figure 4.

If w is the vexillary signed permutation corresponding to a labelled Young diagram Y , then ws_m is vexillary when m is removable: the reader may verify that $Y \setminus m$ is its corresponding labelled Young diagram. The next theorem says the converse also holds.

⁽⁵⁾Kaarel Haenni has announced a proof of this conjecture.




FIGURE 4. Removing corner labels.

THEOREM 4.1. *Let Y be the labelled Young diagram corresponding to a vexillary signed permutation w . Then ws_m is vexillary of length $\ell(w) - 1$ if and only if m is a removable label of Y .*

Proof. Let $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$ be the corresponding essential triple. Using the pattern avoidance criterion for vexillarity, we will show that if a label m is not removable, then ws_m is not vexillary.

First suppose m occurs more than once as a label in Y , or equivalently, the index $p = m + 1$ is repeated in the sequence \mathbf{p} . From the construction of $w(\tau)$, the repetition of p means there is a gap in the increasing sequence of negative integers starting at position p in w . In other words, there is a “partial pattern” $[\bar{3} \bar{1}]$ in w , with the $\bar{3}$ occurring at position p .

If $m = 0$, then the gap must be filled by a positive integer occurring to the right. That is, the pattern $[\bar{3} \bar{1} 2]$ occurs in w , with the “ $\bar{3}$ ” in position 1. We find that ws_0 contains the pattern $[\bar{3} \bar{1} 2]$, so in particular it contains $[2 \ 1]$ and is not vexillary.

If $m > 0$, and the entry of w in position $m = p - 1$ is positive, then one of the following patterns appears:

- $[1 \ \bar{4} \ \bar{2} \ 3]$ (with “1” at position $p - 1$ and “ $\bar{4}$ ” at position p),
- $[\bar{3} \ 1 \ \bar{4} \ \bar{2}]$ (with “1” at position $p - 1$ and “ $\bar{4}$ ” at position p),
- $[2 \ \bar{3} \ \bar{1}]$ (with “2” at position $p - 1$ and “ $\bar{3}$ ” at position p),
- $[2 \ 4 \ \bar{3} \ \bar{1}]$ (with “4” at position $p - 1$ and “ $\bar{3}$ ” at position p), or
- $[\bar{2} \ 4 \ \bar{3} \ \bar{1}]$ (with “4” at position $p - 1$ and “ $\bar{3}$ ” at position p).

If the entry in position $p - 1$ is negative, it must form a partial pattern $[\bar{1} \ \bar{4} \ \bar{2}]$, with the “ $\bar{1}$ ” and “ $\bar{4}$ ” in positions $p - 1$ and p . The gap can be filled to yield one of three patterns:

$$[\bar{1} \ \bar{4} \ \bar{2} \ 3], \quad [\bar{3} \ \bar{1} \ \bar{4} \ \bar{2}], \quad \text{or} \quad [3 \ \bar{1} \ \bar{4} \ \bar{2}].$$

In each case, right-multiplication by s_m yields a forbidden pattern, verifying that ws_m is not vexillary.

Finally, suppose $m = 0$ occurs as a label in a row with more than one box. We may assume the label is not repeated, so this must be the last row of Y , i.e. $p_s = 1$ and

$q_s > 1$. From the construction, this means the positive integer 1 occurs to the right of position 1 in w . Applying s_0 therefore results in the forbidden pattern [2 1]. \square

By reversing the rules for removing a corner, one obtains rules for adding a box to a labelled Young diagram. Suppose Y has labels m_i for corners in rows k_i , and fix an index j such that $m_j - m_{j+1} > 1$. (To include extreme cases $j = 0$ and $j = s$, we allow $0 \leq j \leq s$ and use the conventions $k_0 = 0$, $m_0 = +\infty$, $\lambda_0 = +\infty$, $k_{s+1} = k_s + 1$, $m_{s+1} = -1$, and $\lambda_{k_{s+1}} = 0$.) A new labelled Young diagram $Y \cup m$ is defined as follows, for certain m to be specified.

- (i) If $k_{j+1} - k_j > 1$ and $\lambda_{k_j} - \lambda_{k_{j+1}} > k_{j+1} - k_j + 1$, then $Y \cup m$ is defined for any m satisfying

$$\begin{aligned} m_j &> m > m_{j+1}, \\ m - m_{j+1} &\leq k_{j+1} - k_j, \quad \text{and} \\ m_j - m &\leq \lambda_{k_j} - \lambda_{k_{j+1}} + k_j - k_{j+1}, \end{aligned}$$

by creating a new corner labelled m in row $k_j + 1$ and leaving all other labels unchanged.

- (ii) If $k_{j+1} - k_j = 1$ and $\lambda_{k_j} - \lambda_{k_{j+1}} > 2$, then $Y \cup m$ is defined for $m = m_{j+1} + 1$, by placing a new box labelled m in row $k_j + 1$ and erasing the old label m_{j+1} .
- (iii) If $k_{j+1} - k_j > 1$ and $\lambda_{k_j} - \lambda_{k_{j+1}} = k_{j+1} - k_j + 1$, then $Y \cup m$ is defined for $m = m_j - 1$, by placing a new box labelled m in row $k_j + 1$ and erasing the old label m_j .
- (iv) If $k_{j+1} - k_j = 1$ and $\lambda_{k_j} - \lambda_{k_{j+1}} = 2$, then $Y \cup m$ is defined for $m = m_{j+1} + 1 = m_j - 1$, by placing a new box labelled m in row $k_j + 1$ and erasing the old labels m_j and m_{j+1} .

An integer m is *insertable* in a labelled Young diagram Y if there is a j such that $m_j > m > m_{j+1}$ and one of the above four cases holds.

COROLLARY 4.2. *Suppose w is vexillary, with labelled Young diagram Y . Then ws_m is vexillary of length $\ell(w) + 1$ if and only if m is insertable in Y , in which case its labelled Young diagram is $Y \cup m$.*

For a labelled Young diagram Y with labels m_i in rows k_i , set $l_i = \lambda_{k_i} - m_i - 1$, and define $n(Y) = \max\{m_i + k_i, l_i + k_i\}_{1 \leq i \leq s}$.

COROLLARY 4.3. *Let Y be a labelled Young diagram with corresponding vexillary signed permutation w . If $n \geq n(Y)$, then there is a sequence of vexillary signed permutations $w = w_{(0)}, w_{(1)}, \dots, w_{(r)} = w_{\circ}^{(n)}$, such that for each i there is an m with $w_{(i+1)} = w_{(i)}s_m$ and $\ell(w_{(i+1)}) = \ell(w_{(i)}) + 1$.*

Proof. Unless Y has shape $\lambda = (2n - 1, 2n - 3, \dots, 1)$ with labels $n - 1, n - 2, \dots, 0$ (corresponding to the vexillary signed permutation $w_{\circ}^{(n)}$), there is an insertable m with $0 \leq m \leq n - 1$. Indeed, suppose $n \geq n(Y)$, but no such insertable m exists. Then rules (i) and (ii) imply $m_1 = n - 1$ must appear in the first row; all the rules together then show $m_2 = n - 2$ appears in the second row; and continuing this way, we see 0 must appear in the n th row. A diagram with these labels has $n(Y) \leq n$ only if it is the one corresponding to $w_{\circ}^{(n)}$ (in which case $n(Y) = n$). \square

The inequalities on m_i imply that the labels can be replaced by $l_i = \lambda_{k_i} - m_i - 1$ to form a *dual labelled Young diagram* Y^* . If Y corresponds to the triple τ , then Y^* corresponds to the dual triple τ^* , so the corresponding vexillary signed permutations are inverses, by Lemma 2.4. The analogous results for left-multiplication by s_m follow from these observations.

COROLLARY 4.4. *Let Y be a labelled Young diagram corresponding to a vexillary signed permutation w . Then $s_m w$ is vexillary of length $\ell(w) - 1$ if and only if m is a removable label of Y^* . Similarly, $s_m w$ is vexillary of length $\ell(w) + 1$ if and only if m is insertable in Y^* .*

If $n \geq n(Y)$, then there is a sequence of vexillary signed permutations $w = w_{(0)}, w_{(1)}, \dots, w_{(r)} = w_{\circ}^{(n)}$, such that for each i there is an m with $w_{(i+1)} = s_m w_{(i)}$ and $\ell(w_{(i+1)}) = \ell(w_{(i)}) + 1$.

Inserting a label is a more flexible operation than removing a label. Starting from a given vexillary permutation w , there need not be a descending chain of vexillary permutations $w_{(i)}$ ending in $w_{(r)} = \text{id}$, with $w_{(i+1)} = w_{(i)} s_m$ and $\ell(w_{(i+1)}) = \ell(w_{(i)}) - 1$. For instance, taking $\tau = (2\ 3, 2\ 2, 3\ 1)$ as in Figure 3, the only m such that $\ell(ws_m) = \ell(w) - 1$ is $m = 1$, but this is not a removable label.

However, at least one of Y or Y^* always has a removable label. Letting τ be the corresponding triple, the next statement is proved by induction on the length of $w(\tau)$, removing a label from either Y or Y^* .

COROLLARY 4.5. *For $w = w(\tau)$, we have $\ell(w) = |\lambda(\tau)|$.*

(This can also be proved directly from the construction of $w(\tau)$ by counting inversions.)

5. TRANSITIONS

A signed permutation w is *maximal grassmannian* if its only descent is at 0. A maximal grassmannian signed permutation corresponds to a strict partition λ , by recording the absolute values of the barred entries; for example, $w = 4\ 2\ \bar{1}\ 3$ corresponds to $\lambda = (4, 2, 1)$. It is straightforward to check that maximal grassmannian signed permutations are vexillary: its labelled Young diagram has shape λ and all corners labelled 0. For a strict partition λ , we will write w_λ for the corresponding maximal grassmannian signed permutation.

Transitions provide a way of reducing arbitrary signed permutations to maximal grassmannian ones. They were used by Billey to study Schubert polynomials and Stanley symmetric functions [4].

For $i < j$, let t_{ij} be the transposition exchanging positions i and j , and for $i \leq j$ let s_{ij} exchange i and \bar{j} . (Thus t_{ij} is the reflection in the hyperplane defined by $e_i - e_j$, and s_{ij} is the reflection in the hyperplane defined by $e_i + e_j$.) For any signed permutation w , let m be the last descent, and let j be the largest index greater than m such that $w(m) > w(j)$. A *transition* of w is a signed permutation w^- , of the same length as w , such that $w^- = wt_{mj}t_{im}$ for some $i < m$ or $w^- = wt_{mj}s_{im}$ for any i .

Definitions and properties of the type B Stanley symmetric function H_w may be found in [5] or [10]. Here we need two properties, from [4]. First, for a strict partition λ , we have

$$(5) \quad H_{w_\lambda} = P_\lambda,$$

where the latter is the Schur P -function. Second, for any w , with m and j defined as above, there is a recursive formula

$$(6) \quad H_w = (2H_{wt_{mj}s_{mm}} +) \sum_{w^- \neq wt_{mj}s_{mm}} H_{w^-},$$

the sum over transitions of w , with the first term appearing when $\ell(wt_{mj}s_{mm}) = \ell(w)$ (i.e. when this is also a transition).

LEMMA 5.1. *Let Y be a labelled Young diagram, and assume the largest label m is greater than 0. Suppose $m = m_1 = \cdots = m_r > m_{r+1}$, and let Y^- be the result of replacing the r th corner label m_r by $m - 1$. Let w and w^- be the corresponding vexillary signed permutations. Then w^- is the unique transition of w .*

Proof. Let $\tau = (\mathbf{k}, \mathbf{p}, \mathbf{q})$ be the triple corresponding to Y . From the construction, $m = p_1 - 1 = \cdots = p_r - 1$, and $j = m + k_r$. Note that $w(j) = \bar{q}_r$. The transposition t_{mj} swaps $w(m)$ and $w(j)$; since the sequence $w(p_1), \dots, w(j)$ is increasing, $\ell(wt_{mj}) = \ell(w) - 1$.

If there is an entry to the left of position m that is less than \bar{q}_r , let $i < m$ be the largest such index. Then $w^- = wt_{mj}t_{im}$ is a transition of w , and its labelled Young diagram is Y^- . Furthermore, for any $i < i' < m$, right-multiplication by $t_{i'm}$ decreases the length of wt_{mj} , and for any $0 < i' < i$, right-multiplication by $t_{i'm}$ increases the length of wt_{mj} by at least 2; similarly, $\ell(wt_{mj}s_{i'm}) \neq \ell(w)$ for any i' . So there is no other transition.

If there is no entry to the left of m less than \bar{q}_r , let $i > m$ be the smallest index such that $w(i) > q_r$. Then $w^- = wt_{mj}s_{mi}$ is a transition of w , with labelled Young diagram Y^- . As in the previous case, one checks that this is the only transition. \square

Combining Equations (5) and (6) with Lemma 5.1 yields a formula for the Stanley symmetric function of a vexillary signed permutation.

COROLLARY 5.2. *The Stanley symmetric function $H_{w(\tau)}$ is equal to the Schur P -function $P_{\lambda(\tau)}$.*

It follows that for a vexillary signed permutation $w = w(\tau)$, the partition $\lambda^B(w)$ defined in [6, § 5] is equal to our $\lambda(\tau)$.

Acknowledgements. We are grateful to S. Billey for many helpful suggestions and comments, and to a referee for suggestions for improving the manuscript.

REFERENCES

- [1] David Anderson, *Diagrams and essential sets for signed permutations*, Electron. J. Combin. **25** (2018), no. 3, 3.46 (23 pages).
- [2] David Anderson and William Fulton, *Degeneracy loci, Pfaffians, and vexillary signed permutations in types B, C, and D*, <https://arxiv.org/abs/1210.2066>, 2012.
- [3] ———, *Chern class formulas for classical-type degeneracy loci*, Compos. Math. **154** (2018), no. 8, 1746–1774.
- [4] Sara Billey, *Transition equations for isotropic flag manifolds*, Discrete Math. **193** (1998), no. 1–3, 69–84, Selected papers in honor of Adriano Garsia (Taormina, 1994).
- [5] Sara Billey and Mark Haiman, *Schubert polynomials for the classical groups*, J. Amer. Math. Soc. **8** (1995), no. 2, 443–482.
- [6] Sara Billey and Tao Kai Lam, *Vexillary elements in the hyperoctahedral group*, J. Algebraic Combin. **8** (1998), no. 2, 139–152.
- [7] Anders Björner and Francesco Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, 2005.
- [8] Eric S. Egge, *Enumerating rc -invariant permutations with no long decreasing subsequences*, Ann. Comb. **14** (2010), no. 1, 85–101.
- [9] Kimmo Eriksson and Svante Linusson, *Combinatorics of Fulton’s essential set*, Duke Math. J. **85** (1996), no. 1, 61–76.
- [10] Sergey Fomin and Anatol N. Kirillov, *Combinatorial B_n -analogues of Schubert polynomials*, Trans. Amer. Math. Soc. **348** (1996), no. 9, 3591–3620.
- [11] William Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420.
- [12] William Fulton and Piotr Pragacz, *Schubert varieties and degeneracy loci*, Lecture Notes in Mathematics, vol. 1689, Springer-Verlag, Berlin, 1998, Appendix J by the authors in collaboration with I. Ciocan-Fontanine.

- [13] Takeshi Ikeda, Leonardo C. Mihalcea, and Hiroshi Naruse, *Double Schubert polynomials for the classical groups*, Adv. Math. **226** (2011), no. 1, 840–886.
- [14] Maxim Kazarian, *On Lagrange and symmetric degeneracy loci*, Isaac Newton Institute for Mathematical Sciences Preprint Series, 2000.
- [15] Anatol N. Kirillov, *Notes on Schubert, Grothendieck and key polynomials*, SIGMA Symmetry Integrability Geom. Methods Appl. **12** (2016), Paper No. 034 (56 pages).
- [16] Jordan Lambert, *Theta-vexillary signed permutations*, Electron. J. Combin. **25** (2018), no. 4, Paper 4.53 (30 pages).
- [17] Alain Lascoux and Marcel-Paul Schützenberger, *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), no. 13, 447–450.
- [18] ———, *Schubert polynomials and the Littlewood–Richardson rule*, Lett. Math. Phys. **10** (1985), no. 2-3, 111–124.
- [19] Julian West, *Generating trees and the Catalan and Schröder numbers*, Discrete Math. **146** (1995), no. 1-3, 247–262.

DAVID ANDERSON, Department of Mathematics, The Ohio State University, Columbus, Ohio 43210,
USA
E-mail : `anderson.2804@math.osu.edu`

WILLIAM FULTON, Department of Mathematics, University of Michigan, Ann Arbor, Michigan
48109, USA
E-mail : `wfulton@umich.edu`