

ALGEBRAIC COMBINATORICS

André de Moura, Elijah Gunther, Sam Payne, Jason Schuchardt & Alan Stapledon

Triangulations of simplices with vanishing local h -polynomial

Volume 3, issue 6 (2020), p. 1417-1430.

http://alco.centre-mersenne.org/item/ALCO_2020__3_6_1417_0


© The journal and the authors, 2020.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is member of the
Centre Mersenne for Open Scientific Publishing
www.centre-mersenne.org



Triangulations of simplices with vanishing local h -polynomial

André de Moura, Elijah Gunther, Sam Payne, Jason Schuchardt & Alan Stapledon

ABSTRACT Motivated by connections to intersection homology of toric morphisms, the motivic monodromy conjecture, and a question of Stanley, we study the structure of geometric triangulations of simplices whose local h -polynomial vanishes. As a first step, we identify a class of refinements that preserve the local h -polynomial. In dimensions 2 and 3, we show that all geometric triangulations with vanishing local h -polynomial are obtained from one or two simple examples by a sequence of such refinements. In higher dimensions, we prove some partial results and give further examples.

1. INTRODUCTION

Let Γ be a triangulation of a simplex Δ of dimension $d-1$. The h -polynomial $h(\Gamma; x) = h_0 + h_1x + \cdots + h_dx^d$ is a common and convenient way of encoding the number of faces of Γ in each dimension. It is characterized by the equation

$$\sum_{i=0}^d h_i(x+1)^{d-i} = \sum_{i=0}^d f_{i-1}x^{d-i},$$

where $f_{-1} = 1$ and f_i is the number of i -dimensional faces of Γ , for $i \geq 0$. The coefficients h_i are non-negative integers. One powerful tool for studying $h(\Gamma; x)$ is the local h -polynomial $\ell(\Gamma; x) = \ell_0 + \ell_1x + \cdots + \ell_dx^d$, introduced by Stanley in his seminal paper [7]. It is characterized via Möbius inversion by the equation

$$(1) \quad h(\Gamma; x) = \sum_{F \leq \Delta} \ell(\Gamma_F; x),$$

where Γ_F denotes the restriction of the triangulation Γ to a face F (which may be empty or all of Δ), together with the condition $\ell(\emptyset; x) = 1$.

The local h -polynomial has remarkable properties. In particular, the coefficients ℓ_i are nonnegative and satisfy $\ell_i = \ell_{d-i}$. Moreover, if the subdivision is regular, then these coefficients are unimodal. Among other applications, Stanley used local h -polynomials to prove that h -polynomials increase coefficientwise under refinement.

As discussed in Section 2, the local h -polynomial also behaves predictably with respect to basic operations on subdivisions. It is additive for refinements that nontrivially subdivide only one facet, multiplicative for joins, and vanishes on the trivial subdivision. In particular, if Γ' is a refinement of Γ that nontrivially subdivides only


Manuscript received 9th March 2020, revised 19th August 2020, accepted 20th August 2020.

KEYWORDS. local h -polynomials, triangulations of simplices, geometric triangulations.

ACKNOWLEDGEMENTS. This work was partially supported by NSF grant DMS-1702428.

one facet, and that subdivision is the cone over a subdivision of a codimension 1 face, then $\ell(\Gamma'; x) = \ell(\Gamma; x)$. We call such subdivisions *conical facet refinements*.

Our aim is to study geometric triangulations with vanishing local h -polynomial. One might hope that all such subdivisions are obtained from the trivial subdivision by a sequence of conical facet refinements. However, not all triangulations Γ with vanishing local h -polynomial can be obtained from the trivial subdivision in this way. One notable example in dimension 2 is the following subdivision, which we call the *triforce*⁽¹⁾.



All of our theorems are for *geometric triangulations*, and all triangulations in this paper are assumed to be geometric.

THEOREM 1.1. *In dimension 2, any triangulation with vanishing local h -polynomial is obtained from either the trivial subdivision or the triforce by a sequence of conical facet refinements.*

The two cases in the theorem are distinguished by the *internal edge graph*, i.e. the union of the edges that meet the interior of the subdivided simplex, which also figures prominently in the proof. For an iterated conical facet refinement of the triforce subdivision, the internal edge graph has Euler characteristic zero and contains no vertices of the original triangle. For an iterated conical facet refinement of the trivial subdivision in dimension 2, the internal edge graph is a tree that contains exactly one of the vertices of the original triangle.

In dimension three, the structural classification is even simpler.

THEOREM 1.2. *In dimension 3, any triangulation with vanishing local h -polynomial is obtained from the trivial subdivision by a sequence of conical facet refinements.*

The proof again relies on an analysis of the internal edge graph, which in dimension three is a union of trees, each of which contains exactly one vertex supported on a face of codimension at least 2 in the original simplex.

This structure of the internal edge graph is similar in higher dimensions. Note that $\ell_0 = 0$ for any triangulation of a nonempty simplex.

THEOREM 1.3. *Let Γ be a triangulation of a simplex of dimension at least 3 such that $\ell_1 = \ell_2 = 0$. Then the internal edge graph of Γ is a union of trees each of which contains exactly one vertex supported on a face of codimension at least 2.*

However, the pattern of obtaining all triangulations whose local h -polynomials vanish from a finite collection of examples by iterated conical facet refinements does not continue in higher dimensions. See Section 5.

Our investigation into the structure of triangulations with vanishing local h -polynomial is motivated by recent connections to algebraic and arithmetic geometry. Local h -polynomials appear prominently in formulas for dimensions of homology groups of intersection complexes for toric morphisms [3] and multiplicities of eigenvalues of monodromy [6, 8]. In Igusa’s p -adic monodromy conjecture [5], and the motivic generalization of Denef and Loeser [4], the essential question is understanding whether or not these multiplicities vanish. This problem is also natural and interesting from a

⁽¹⁾This name reflects the subdivision’s realization in a sacred golden relic that is the ultimate source of power in the action-adventure video game series *The Legend of Zelda*.

purely combinatorial viewpoint; Stanley specifically asked for a nice characterization of such triangulations in his original paper [7, Problem 4.13].

2. PRELIMINARIES

We consider only geometric subdivisions Γ of a $(d - 1)$ -simplex Δ , except where explicitly stated otherwise. In particular, all of the triangulations that we consider are realized by subdividing a linearly embedded simplex into subsimplices. For further details and background on local h -polynomials, we refer the reader to the recent survey article of Athanasiadis [1], as well as [7].

2.1. FORMULAS FOR THE LOCAL h -POLYNOMIAL. We recall two useful formulas for the local h -polynomial $\ell(\Gamma; x)$. First, by applying Möbius inversion to (1), we can express $\ell(\Gamma; x)$ as an alternating sum of h -polynomials:

$$(2) \quad \ell(\Gamma; x) = \sum_{F \leq \Delta} (-1)^{\text{codim}(F)} h(\Gamma_F; x).$$

Here, the codimension is $\text{codim}(F) = d - 1 - \dim(F)$, since $\dim(\Delta) = d - 1$.

Let $\sigma : \Gamma \rightarrow \Delta$ be the map taking a face $G \in \Gamma$ to the smallest face $F \leq \Delta$ that contains it. One says that G is *carried by* $\sigma(G)$, and the *excess* of G is

$$e(G) := \dim(\sigma(G)) - \dim(G).$$

As in [7, Prop. 2.2], we can then express the local h -polynomial as

$$(3) \quad \ell(\Gamma; x) = \sum_{G \in \Gamma} (-1)^{\text{codim}(G)} x^{d-e(G)} (x - 1)^{e(G)}.$$

DEFINITION 2.1. We say that $G \in \Gamma$ is an interior face if $\sigma(G) = \Delta$.

2.2. ELEMENTARY PROPERTIES OF THE LOCAL h -POLYNOMIAL. The following basic properties of local h -polynomials were mentioned in the introduction and will be used in our main arguments.

Given triangulations Γ and Γ' of simplices Δ and Δ' , respectively, the join $\Gamma * \Gamma'$ is naturally a triangulation of the simplex $\Delta * \Delta'$. It is well-known that h -polynomials are multiplicative for joins, i.e.

$$h(\Gamma * \Gamma'; x) = h(\Gamma; x) \cdot h(\Gamma'; x).$$

It follows that the local h -polynomial has the analogous property [2, Lemma 2.2], i.e.

$$(4) \quad \ell(\Gamma * \Gamma'; x) = \ell(\Gamma; x) \cdot \ell(\Gamma'; x).$$

A triangulation Γ' of Δ is a *refinement* of Γ if every $G \in \Gamma$ is a union of faces of Γ' . We then write Γ'_G for the restriction of Γ' to G . We are particularly interested in the restriction of Γ' to maximal faces, or *facets*.

DEFINITION 2.2. Let Γ' be a refinement of Γ , and let G be a facet of Γ . We say that Γ' is a *facet refinement* of Γ along G if, for any facet $H \neq G$, the restriction Γ'_H is trivial.




FIGURE 1. A facet refinement of the triforme along its lower left facet.

Local h -polynomials behave additively with respect to facet refinements.

PROPOSITION 2.3. *Let Γ' be a facet refinement of Γ along G , and let Γ'_G denote the triangulation of G induced by restricting Γ' . Then*

$$\ell(\Gamma'; x) = \ell(\Gamma; x) + \ell(\Gamma'_G; x).$$

This follows from [6, Corollary 4.7], a much more general result about local h -polynomials for compositions of strong formal subdivisions, applied to $\Gamma' \rightarrow \Gamma \rightarrow \Delta$. For the reader's convenience, we include a direct proof in our setting.

Proof. The assertion is vacuously true in dimension zero. We proceed by induction on dimension. First, we observe that

$$(5) \quad h(\Gamma'; x) = h(\Gamma; x) + h(\Gamma'_G; x) - h(G; x).$$

This follows from the formula $\sum_{i=0}^d f_i(x-1)^{d-i} = \sum_{i=0}^d h_i x^{d-i}$, by inclusion-exclusion.

Next, write each h -polynomial in (5) as a sum of local h -polynomials, using (1). The contributions of the empty faces cancel, and each nonempty face of G contributes 0, since it is trivially subdivided. Therefore, we have

$$(6) \quad \sum_{\{\emptyset\} \neq H \leq \Delta} \ell(\Gamma'_H; x) = \sum_{\{\emptyset\} \neq H \leq \Delta} \ell(\Gamma_H; x) + \sum_{\{\emptyset\} \neq H' \leq G} \ell(\Gamma'_{H'}; x).$$

For each nonempty face $H \leq \Delta$, either $\Gamma'_H = \Gamma_H$ or Γ'_H is a facet refinement of Γ_H along $G \cap H$. If H is a proper face of Δ then, by induction on dimension, we may assume $\ell(\Gamma'_H; x) = \ell(\Gamma_H; x) + \ell(\Gamma'_{G \cap H}; x)$. Similarly, for any nonempty face $H' \leq G$, either $\Gamma'_{H'}$ is the trivial subdivision, or H' is carried by a face $H \leq \Delta$ such that $H' = G \cap H$ and Γ'_H is a facet refinement of Γ_H along H' . Thus, the contributions of proper faces in (6) cancel, and we conclude that $\ell(\Gamma'; x) = \ell(\Gamma; x) + \ell(\Gamma'_G; x)$. \square

DEFINITION 2.4. *A facet refinement Γ' of Γ along G is a conical facet refinement if Γ'_G is the cone over Γ'_H for some codimension 1 face $H < G$.*

COROLLARY 2.5. *Let Γ' be a facet refinement of Γ along G , and suppose that Γ'_G is the join of two triangulations of faces of G , one of which has vanishing local h -polynomial. Then $\ell(\Gamma'; x) = \ell(\Gamma; x)$. In particular, if Γ' is a conical facet refinement of Γ , then $\ell(\Gamma'; x) = \ell(\Gamma; x)$.*

Proof. This follows immediately from (4) and Proposition 2.3. \square




FIGURE 2. A triangulation obtained from the triform by a series of conical facet refinements.

3. THE INTERNAL EDGE GRAPH

The proofs of the main results of this paper all involve the *internal edge graph*, formed by the edges of a triangulation Γ that meet the interior of the simplex Δ . In this section, we study the properties of this graph when $d \geq 3$, and $\ell_1 = \ell_2 = 0$.

Let f_i^j denote the number of i -simplices in Γ that are carried by j -faces of Δ . From equation (3), we see that $\ell_0 = 0$, $\ell_1 = f_0^{d-1}$, and

$$(7) \quad \ell_2 = f_1^{d-1} - f_0^{d-2} - (d-1)f_0^{d-1}.$$

In particular, if $\ell_1 = 0$ then $\ell_2 = f_1^{d-1} - f_0^{d-2}$.

Stanley proved non-negativity of all ℓ_i , using methods from commutative algebra. It follows that if there are no interior vertices (i.e. if $\ell_1 = 0$) then $f_1^{d-1} \geq f_0^{d-2}$. Note that every vertex carried by a $(d-2)$ -face is contained in an interior edge, so this inequality is a statement about the internal edge graph. We now give a combinatorial proof of this inequality in a stronger form, showing in particular that it holds separately on each connected component of the internal edge graph.

DEFINITION 3.1. *The internal edge graph of a subdivision Γ of Δ is the graph $\Sigma(\Gamma)$ whose edges are the edges of Γ carried by the improper face Δ , and whose vertices are the vertices incident to those edges.*

When Γ is clear from context, we will write Σ rather than $\Sigma(\Gamma)$.

PROPOSITION 3.2. *Let Γ be a triangulation of a $(d-1)$ -simplex Δ with $d \geq 3$ and $\ell_1 = 0$. Then each connected component C of $\Sigma(\Gamma)$ contains either a vertex of excess less than $d-2$ or a simple 3-cycle.*

Furthermore, if $d \geq 4$, and C has no vertices of excess less than $d-2$ then C contains at least two distinct simple 3-cycles.

Proof. We may assume that Δ is the standard $(d-1)$ -simplex in \mathbb{R}^d , i.e. the convex hull of the standard basis vectors e_1, \dots, e_d . Let $F_i = \text{Conv}\{e_1, \dots, \widehat{e_i}, \dots, e_d\}$, and let $\pi_i : \mathbb{R}^d \rightarrow \mathbb{R}$ be the projection onto the i^{th} coordinate axis.

Suppose C is a connected component of Σ all of whose vertices have excess $d-2$. We now show that C contains a simple 3-cycle.

Consider the two codimension 1 faces carrying the endpoints of an edge of C . Without loss of generality, we may assume these to be F_1 and F_2 . Let L be the linear functional $\pi_1 + \pi_2$. Let $e = \text{Conv}\{v_1, v_2\}$ be an edge of C such that $v_1 \in F_1$, and $v_2 \in F_2$, such that $L(v_1 + v_2)$ is maximal among all such edges. For $(i, j) \in \{(1, 2), (2, 1)\}$, let $P_i : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be the projection to the linear hyperplane spanned by F_i along $v_j - v_i$. More explicitly, for all $w \in \mathbb{R}^d$,

$$P_i(w) = w - \frac{\pi_i(w)}{\pi_i(v_j)}(v_j - v_i).$$

Substituting in the definitions above, we observe that

$$\begin{aligned} \pi_i(v_j)L(P_i(w) - v_i) &= \pi_i(v_j)\pi_j(P_i(w)) - \pi_i(v_j)\pi_j(v_i) \\ &= \pi_i(v_j)\pi_j(w) + \pi_j(v_i)\pi_i(w) - \pi_i(v_j)\pi_j(v_i). \end{aligned}$$

This is invariant under switching i and j , and hence

$$(8) \quad \pi_1(v_2)L(P_1(w) - v_1) = \pi_2(v_1)L(P_2(w) - v_2).$$

Applying P_1 to the closed star $\text{Cl}(\text{Star}(e))$, we see that e projects to v_1 , and 2-faces project to segments ending at v_1 . Since e is an interior edge, v_1 is in the relative interior of $P_1(\text{Cl}(\text{Star}(e)))$. Therefore, for any hyperplane H in the image of P_1 that contains v_1 , there is a vertex of the link of e that projects to either side of that hyperplane. In

particular, there is a vertex w in the link of e such that $L(P_1(w) - v_1) > 0$. Recalling that all vertices of C have excess $d - 2$, we note that $\pi_1(v_2) > 0$ and $\pi_2(v_1) > 0$, otherwise they would be contained in $F_1 \cap F_2$. Therefore, by (8), $L(P_2(w) - v_2) > 0$.

We claim that w is not carried by F_1 or F_2 . Indeed, if w is carried by F_1 , we would have $P_1(w) = w$ and $L(v_2 + w) > L(v_2 + v_1)$, contradicting our maximality assumption on e . Thus $w \notin F_1$. An identical argument shows that w is not carried by F_2 . This proves the claim. Then without loss of generality, we may assume that w is carried by F_3 . The three interior edges of C connecting v_1 , v_2 , and w form a simple 3-cycle. If $d = 3$, then we are done.

Suppose $d \geq 4$. We must show that C contains another simple 3-cycle. Since w is in $\text{lk } e$, Γ has an interior 2-face $G = \text{Conv}\{v_1, v_2, w\}$. Let p denote the projection along G onto the codimension 2 linear subspace spanned by $F_1 \cap F_2$. Then $p(G)$ is in the relative interior of $p(\text{Cl}(\text{Star}(G)))$, and hence there exists a vertex $w' \notin F_1 \cap F_2 \cap F_3$ in the link of G . Without loss of generality, $w' \notin F_1$. Then $\text{Conv}\{v_1, w'\}$ is an interior edge in C , and hence w' is carried by a codimension 1 face of Δ . Since $w' \notin F_2 \cap F_3$, we may assume, without loss of generality, that $w' \notin F_2$. Then the edges of G and $\text{Conv}\{v_1, v_2, w'\}$ form two distinct simple 3-cycles in C . \square

Let $f_i^j(C)$ denote the number of i -faces of C that are carried by j -faces of Δ .

COROLLARY 3.3. *Suppose $d \geq 3$ and $\ell_1 = 0$. Then each connected component C of Σ has*

$$f_0^{d-2}(C) \leq f_1^{d-1}(C).$$

Proof. Recall that the Euler characteristic of a connected graph C is

$$\chi(C) = \#\{\text{vertices of } C\} - \#\{\text{edges of } C\}.$$

This is also equal to $1 - h^1(C)$, where h^1 denotes the first Betti number. Hence $\chi(C) \leq 1$, with equality if and only if C is a tree. Moreover, $\chi(C) = 0$ if and only if C contains a unique simple cycle.

Let C be a connected component of the internal edge graph. Then

$$f_0^{d-2}(C) - f_1^{d-1}(C) = \chi(C) - f_0^{<d-2}(C).$$

By Proposition 3.2, either $\chi(C) = 1$ and $f_0^{<d-2}$ is positive, or $\chi(C) \leq 0$. \square

In Section 4, we will repeatedly use the following structural description of the internal edge graph.

PROPOSITION 3.4. *Suppose $d \geq 3$, $\ell_1 = \ell_2 = 0$, and C is a connected component of Σ . Then either*

- (1) C is a tree with a unique vertex of excess less than $d - 2$, or
- (2) $d = 3$, every vertex of C has excess $d - 2$, and C has a unique simple 3-cycle.

Proof. Since $\ell_1 = 0$, we can write ℓ_2 as a sum over connected components of Σ

$$\begin{aligned} \ell_2 &= \sum_C f_1^{d-1}(C) - f_0^{d-2}(C) \\ &= \sum_C f_0^{<d-2}(C) - \chi(C). \end{aligned}$$

By Proposition 3.2 and Corollary 3.3, each summand is nonnegative, and the summand corresponding to C is zero if and only if either C is a tree with a unique vertex of excess less than $d - 2$, or $d = 3$, every vertex of C has excess $d - 2$, and C contains a unique simple 3-cycle. \square

Note that Theorem 1.3 follows immediately from Proposition 3.4.

REMARK 3.5. Note that the internal edge graph of any triangulation of a 2-simplex is connected. To see this, consider a polyhedral subdivision of a triangle with disconnected internal edge graph. It must contain a polygon that meets two different components of the internal edge graph, and such a polygon has at least four edges: at least two segments of the boundary, and at least one edge from each component of the internal edge graph that it meets. In particular, it is not a triangulation.

DEFINITION 3.6. Let F be a facet in a triangulation of the simplex Δ . We say that F is a pyramid if there is some proper face of Δ that contains every vertex of F except one.

REMARK 3.7. Understanding when a facet of a triangulation is a pyramid is helpful for potential applications, e.g. to the monodromy conjecture for nondegenerate hypersurfaces, as mentioned in the introduction. Note that, if Γ' is a conical facet refinement of Γ along G , then both G and every facet of Γ'_G is such a pyramid. Hence, it follows from Theorem 1.2 that, if $d = 4$ and $\ell(\Gamma; x) = 0$, then every facet is a pyramid. Similarly, it follows from Theorem 1.1 that, if $d = 3$ and $\ell(\Gamma; x) = 0$, then at most one facet is not a pyramid.

The following proposition is not used in the remainder of the paper, but illustrates how the structure of the internal edge graph gives useful statements about which facets of a triangulation with vanishing local h -polynomial are pyramids.

PROPOSITION 3.8. Let Γ be a triangulation of a $(d - 1)$ -simplex Δ with $d \geq 4$. If $\ell_1 = \ell_2 = 0$, then every facet G of Γ that contains an interior edge is a pyramid.

Proof. Let Γ be a facet of a triangulation of a $(d - 1)$ -simplex Δ with $d \geq 4$, such that $\ell_1 = \ell_2 = 0$. Let G be a facet of Γ that contains an interior edge. We assume that G is not a pyramid, and will derive a contradiction. Let $e_1 = \text{Conv}\{v_1, v_2\}$ be an interior edge in G . Let C be the connected component of the interior edge graph $\Sigma(\Gamma)$ containing e_1 .

By Proposition 3.4, either v_1 or v_2 is carried by a codimension 1 face. Without loss of generality, v_1 is carried by a codimension 1 face F_1 . Note that $v_2 \notin F_1$. Since G is not a pyramid, it has a vertex $w_2 \notin F_1$ distinct from v_2 . Then $e_2 = \text{Conv}\{v_1, w_2\}$ lies in C . By Proposition 3.4, either v_2 or w_2 is carried by a codimension 1 face. Without loss of generality, v_2 is carried by a codimension 1 face F_2 . If $w_2 \notin F_2$, then $\text{Conv}\{v_2, w_2\}$ is an interior edge and C contains a cycle, contradicting Proposition 3.4. Hence, $w_2 \in F_2$. Since G is not a pyramid, it has a vertex $w_1 \notin F_2$ distinct from v_1 . Then $e_3 = \text{Conv}\{w_1, v_2\}$ lies in C . If $w_1 \notin F_1$, then $\text{Conv}\{v_1, w_1\}$ is an interior edge and C contains a cycle, contradicting Proposition 3.4. Hence $w_1 \in F_1$. By Proposition 3.4, either w_1 or w_2 is carried by a codimension 1 face, meaning that they cannot both be contained in some third codimension 1 face F_3 . Therefore, $e_4 = \text{Conv}\{w_1, w_2\}$ lies in C , and e_1, e_3, e_4, e_2 forms a cycle in C , contradicting Proposition 3.4. \square

4. DIMENSIONS 2 AND 3

In this section we prove our main structural results, that all subdivisions with vanishing local h can be obtained by a sequence of conical facet refinements from the trivial subdivision and the triforce subdivision in dimension 2, and from the trivial subdivision in dimension 3. For both results our proof is by induction on the number of vertices or interior edges in the subdivision. In the induction step, we identify a subcomplex that arises from a conical subdivision of a facet in a coarser subdivision.

Recall that the support of a subcomplex $\Gamma' \subset \Gamma$ is the union of the faces in Γ' . It is a closed subset of Δ . The relative boundary of any closed subset $F \subset \Delta$ is the intersection of F with the closure of its complement $\Delta \setminus F$. Note, in particular, that if F is the support of a subcomplex $\Gamma' \subset \Gamma$, then the relative boundary of F is the

support of a subcomplex of Γ' . When no confusion seems possible, we will refer to this subcomplex as the relative boundary of Γ' .

LEMMA 4.1. *Let Γ be a triangulation of a simplex Δ and let $\Gamma' \subseteq \Gamma$ be a subcomplex whose support is a simplex F of dimension $d - 1$. Suppose that Γ' induces the trivial subdivision on the relative boundary of F in Δ . Then $\ell(\Gamma; x) - \ell(\Gamma'; x)$ has nonnegative coefficients. In particular, if $\ell(\Gamma; x) = 0$, then $\ell(\Gamma'; x) = 0$.*

Proof. Since Γ' does not nontrivially subdivide the relative boundary of F in Δ , we get a subdivision Γ'' of Δ by replacing Γ' with F itself. Note that Γ is then a facet refinement of Γ'' along F , so by Proposition 2.3, we have $\ell(\Gamma; x) = \ell(\Gamma''; x) + \ell(\Gamma'; x)$, and the lemma follows. \square

REMARK 4.2. Note that, in the proof of the lemma, Γ is a conical facet refinement of Γ'' (resp. obtained from Γ'' by a sequence of conical facet refinements) if and only if Γ' is a conical facet refinement of F (resp. obtained from the trivial subdivision of F by a sequence of conical facet refinements).

4.1. DIMENSION 2. We now apply the lemma to prove Theorem 1.1, which says that any triangulation of a 2-dimensional simplex with vanishing local h -polynomial is obtained from either the trivial subdivision or the triforce by a sequence of conical facet refinements.

Proof of Theorem 1.1. If Γ has no interior edges, then it is the trivial subdivision; this case of the theorem is obvious. We proceed by induction on the number of interior edges, and consider two cases, according to the possible structures of the internal edge graph given by Proposition 3.4. (Recall also that the internal edge graph is connected, by Remark 3.5.)

CASE 1: $\Sigma(\Gamma)$ is a tree that contains exactly one vertex of excess zero.

We claim that Γ is obtained from the trivial subdivision of Δ by a sequence of conical facet refinements. To see this, note that any interior edge that contains a vertex of excess zero divides Γ into two triangular subcomplexes whose relative boundary is not subdivided, as shown.



These two subcomplexes have vanishing local h -polynomials, by Lemma 4.1. By induction, each is obtained from the trivial subdivision by a sequence of conical facet refinements. Hence Γ is obtained from the subdivision into two triangles by a sequence of conical facet refinements. The subdivision into two triangles is itself a conical facet refinement of the trivial subdivision, and we conclude that Γ is obtained from the trivial subdivision by a sequence of conical facet refinements, as claimed.

CASE 2: $\Sigma(\Gamma)$ contains a simple 3-cycle and all of its vertices have excess 1.

We claim that Γ is obtained from the triforce by a sequence of conical facet refinements. To see this, note that the three vertices of the simple 3-cycle must be carried by the three sides of Δ . This splits Γ into four triangular subcomplexes, each with unsubdivided relative boundary, i.e. Γ is a refinement of the triforce, and the induced subdivision on the relative boundary of each of the four triangles is trivial. Hence, by Lemma 4.1, each of the four induced subdivisions has vanishing local h -polynomial. In fact, the induced subdivision of the interior triangle must be trivial (because Γ has no interior vertices). In each of the other three triangles, the induced subdivision cannot contain a cycle of interior edges, and hence the induced subdivisions are obtained

from the trivial subdivision by a sequence of conical facet refinements. We conclude that Γ is obtained from the trifold by a sequence of conical facet refinements, as claimed. \square

4.2. DIMENSION 3. We now apply similar inductive arguments to prove Theorem 1.2, which says that any subdivision Γ of a 3-dimensional simplex Δ with vanishing local h -polynomial is obtained from the trivial subdivision by a sequence of conical facet refinements.

Proof of Theorem 1.2. If Γ has only 4 vertices, then it is the trivial subdivision and the conclusion is obvious. We proceed by induction on the number of vertices. By Proposition 3.4, each connected component of the internal edge graph $\Sigma(\Gamma)$ is a tree with exactly one vertex of excess less than 2.

CASE 1: $\Sigma(\Gamma)$ is empty.

Let F be a 2-face of Δ . We claim that $\ell(\Gamma_F; x) = 0$. To see this, note that any vertex carried by F must be contained in an interior edge. Hence Γ_F has no interior vertices, and $\ell_1(\Gamma_F; x) = 0$. Now Γ_F is a triangulation of the 2-simplex, so the symmetry of the local h -polynomial implies that $\ell_i(\Gamma_G; x) = \ell_{3-i}(\Gamma_F; x)$. Hence $\ell_2(\Gamma_F; x) = \ell_1(\Gamma_F; x) = 0$, and we conclude that $\ell(\Gamma_F; x) = 0$, as claimed.

We proceed to consider three subcases, according to the internal edge graphs of the restrictions of Γ to the 2-faces of Δ .

SUBCASE 1.1: All 2-faces have empty internal edge graph.

Any interior 2-face of Γ contains an edge that is carried by Γ_F for some 2-face $F \leq \Delta$. So Γ has no interior 2-faces, and hence is the trivial subdivision.

SUBCASE 1.2: Some 2-face has a 3-cycle in its internal edge graph.

Let F_1 be a 2-face of Δ such that Γ_{F_1} contains a 3-cycle of interior edges. The vertices of this 3-cycle, v_2, v_3 , and v_4 , are carried by distinct edges of F_1 . Label the other 2-faces of Δ as F_2, F_3, F_4 so that v_i is carried by $F_1 \cap F_i$.

The 2-face $\text{Conv}\{v_2, v_3, v_4\}$ is contained in some 3-face $T = \text{Conv}\{v_2, v_3, v_4, x\}$ of Γ . Since $\Sigma(\Gamma)$ is empty, the edges $\text{Conv}\{x, v_i\}$ must be contained in the boundary of Δ . Hence x must lie in the intersection $F_2 \cap F_3 \cap F_4$, which is the vertex of Δ opposite F_1 . Then the 2-faces $\text{Conv}\{v_1, v_2, x\}$, $\text{Conv}\{v_2, v_3, x\}$ and $\text{Conv}\{v_1, v_3, x\}$ cut Δ into four tetrahedral regions (the cone over a trifold subdivision of F_1). By Lemma 4.1, the restriction of Γ to each of these regions has vanishing local h -polynomial. By induction, the induced subdivision of each of these regions is obtained from the trivial subdivision by a sequence of conical facet refinements, and we are done.

SUBCASE 1.3: No 2-face has a 3-cycle, and some 2-face has an interior edge.

Let F be a 2-face of Δ such that Γ_F contains an interior edge. By Proposition 3.4, there is an edge e carried by F with a vertex of excess 0. Label the vertices of Δ as A, B, C , and D , so that $F = \text{Conv}\{B, C, D\}$ and e contains D . Let $E \in \text{Conv}\{B, C\}$ be the second vertex of e .

Let G be an interior 2-face of Γ containing e . Then $G = \text{Conv}\{e, v\}$ for some vertex v of Γ . This vertex v must lie on $\text{Conv}\{A, B\}$ or $\text{Conv}\{A, C\}$, so that the edges $\text{Conv}\{E, v\}$ and $\text{Conv}\{D, v\}$ are not interior. Furthermore, $v \neq B, C$, since G is interior. Then either $v = A$, or, without loss of generality, we may assume $\sigma(v) = \text{Conv}\{A, C\}$. These possibilities are illustrated in Figure 3.

If $v = A$ then G , marked in grey in Figure 3a, divides Δ into two distinct tetrahedra, that satisfy the conditions of the Lemma 4.1, and by the usual induction argument we are done.

We may therefore assume $\sigma(v) = \text{Conv}\{A, C\}$ as in Figure 3b. In this case, G cuts off a tetrahedron $\text{Conv}\{v, C, D, E\}$, which satisfies the conditions of Lemma 4.1. By




FIGURE 3. Possible locations for v in Subcase 1.3

induction, it suffices to consider the case where this tetrahedron is trivially subdivided. Also since G is an interior 2-face, it must be contained in another tetrahedron $\text{Conv}\{v, D, E, w\}$ of Γ . We must have $w \in \text{Conv}\{v, A\} \cup \text{Conv}\{A, B\} \cup \text{Conv}\{E, B\} \setminus \{v, E\}$, in order to prevent the edges $\text{Conv}\{w, v\}$ and $\text{Conv}\{w, E\}$ from being interior, and so that w is on the other side of G from C .

Now if $\sigma(w) = \text{Conv}\{A, B\}$, then $\Gamma_{\text{Conv}\{A, B, C\}}$ will contain a cycle of interior edges, with vertices v, E, w . But we assumed that no 2-face has a cycle in its internal edge graph. Thus, $w \in \text{Conv}\{v, A\} \setminus \{v\}$ or $w \in \text{Conv}\{E, B\} \setminus \{E\}$. Without loss of generality, we may assume $w \in \text{Conv}\{E, B\} \setminus \{E\}$. Then regardless of whether $w = B$ or is instead in the interior of the segment, the tetrahedron $\text{Conv}\{v, D, w, C\}$ is one to which Lemma 4.1 applies. Moreover, $\Gamma_{\text{Conv}\{v, D, w, C\}}$ is nontrivial, because G lies in the interior. Hence $\Gamma_{\text{Conv}\{v, D, w, C\}}$ is obtained from the trivial subdivision by a series of conical facet refinements, and the conclusion follows by induction.

CASE 2: $\Sigma(\Gamma)$ is nonempty.

By Proposition 3.4, we know that each connected component of $\Sigma(\Gamma)$ is a tree rooted at a unique vertex of excess less than 2. We consider one of these components C and let v_1 be one of its leaves at maximal distance from the root. Let w be the unique vertex adjacent to v_1 in C , and let F be the 2-face carrying v_1 . Let v_2, \dots, v_r be the other leaves that are adjacent to w and carried by F , as in Figure 4.




FIGURE 4. Structure of C .

Let X be the subcomplex of Γ_F that is the union of the closed stars of v_1, \dots, v_r .

CLAIM: $X \subseteq \text{Cl}(\text{Star}(w))$.

In other words, we claim that Γ contains, as a subcomplex, the cone over X with vertex w . To prove this, it suffices to show that if G is a k -face of Γ_F containing some

vertex v_i , then G is contained in a face of $\text{Star}(w)$. Any such G must be contained in an interior $(k+1)$ -face of Γ . Let w' be a vertex such that $\text{Conv}\{G, w'\}$ is an interior $(k+1)$ -face. Note that $\text{Conv}\{v_i, w'\}$ is an interior edge. Since v_i is a leaf of the internal edge graph, we have $w' = w$. Hence $\text{Conv}\{G, w\}$ is a face of $\text{Star}(w)$ that contains G . This proves the claim.

SUBCASE 2.1: *There is a vertex of X carried by F that is not in $\{v_1, \dots, v_r\}$.* Let v be a vertex of X that is carried by F and not contained in $\{v_1, \dots, v_r\}$. Since $v \in X$, there is an edge $\text{Conv}\{v, v_i\}$ in X for some index i . By the claim proved above, $\text{Conv}\{v, v_i, w\}$ is a face of Γ , and $\text{Conv}\{v, w\}$ is an edge of C . Since $v \notin \{v_1, \dots, v_r\}$, it must be the parent of w .

Consider the link $\text{lk}_{\Gamma_F}(v)$. This is a cycle of edges in Γ_F that contains v_i . If every edge in $\text{lk}_{\Gamma_F}(v)$ contains some $v_j \in \{v_1, \dots, v_r\}$, then $\text{Star}(v)$ is contained in $\text{Cl}(\text{Star}(w))$ and hence $\text{Conv}\{v, w\}$ is the unique interior edge containing v . This contradicts the fact that v is the parent of w . Hence $\text{lk}_{\Gamma_F}(v)$ contains multiple vertices not in $\{v_1, \dots, v_r\}$.

Choose vertices w_1 and w_2 in $\text{lk}_{\Gamma_F}(v)$ that are not in $\{v_1, \dots, v_r\}$ so that every vertex in the path between w_1 and w_2 that contains v_i is in $\{v_1, \dots, v_r\}$. Note that, since w_1 and w_2 are adjacent to w and not in $\{v_1, \dots, v_r\}$, and since v is the parent of w in C , the edges $\text{Conv}\{w, w_i\}$ must not be interior. Similarly, w cannot be the root of C . Hence w is carried by a 2-face that contains both w_1 and w_2 .

It follows that w_1 and w_2 lie on an edge of F . Thus the subcomplex with support $\text{Conv}\{v, w_1, w_2, w\}$ satisfies the conditions of Lemma 4.1.

SUBCASE 2.1.1: *$\text{Conv}\{v, w_1, w_2, w\}$ is trivially subdivided.*

We claim that this is impossible. Indeed, if $\text{Conv}\{v, w_1, w_2, w\}$ is trivially subdivided, then $\text{lk}_{\Gamma_F}(v)$ consists of the edge $\text{Conv}\{w_1, w_2\}$ together with a sequence of edges connecting w_1 to w_2 in which all other vertices are contained in $\{v_1, \dots, v_r\}$. Then every facet containing v is a pyramid with base on F and vertex at w . Hence, v is a leaf of C . This is impossible, since we have already shown that v is the parent of w . This proves the claim.

SUBCASE 2.1.2: *$\text{Conv}\{v, w_1, w_2, w\}$ is nontrivially subdivided.*

In this case, by the induction hypothesis, $\Gamma_{\text{Conv}\{v, w_1, w_2, w\}}$ is obtained from the trivial subdivision by a series of conical facet refinements, and the result follows by induction.

SUBCASE 2.2: *Every vertex of X carried by F is in $\{v_1, \dots, v_r\}$.*

In this case, every edge in the boundary of X connects two vertices that are in the boundary of F . Then the closure X' of any connected component of the relative interior of X is a subcomplex with support a polygon whose relative boundary in F is trivially subdivided. If the support of X' is a triangle, then we can apply Lemma 4.1 for the subcomplex of Γ given by the cone over X' with vertex w , and be done by induction.

If X' is not a triangle, then we show that X' is contained in a larger subcomplex $Y \subset \Gamma_F$ such that

- (1) The relative boundary of Y in F is trivially subdivided.
- (2) The cone over Y with vertex w is a subcomplex of Γ .
- (3) The support of Y is a triangle.

Once we find such a Y , then the conclusion of the theorem follows by Lemma 4.1 and induction on the number of vertices.

In order to find Y as above, we begin with $Y_0 = X'$, and construct an increasing sequence of subcomplexes

$$Y_0 \subset Y_1 \subset \dots \subset Y_n = Y$$

such that each Y_i has properties (1) and (2), and the support of Y is a triangle.

We describe the construction of Y_{i+1} from Y_i in terms of corner cutting edges.

DEFINITION 4.3. A boundary edge e of Y_i is corner cutting if its vertices are carried by edges of F , and Y_i lies on the side of e opposite from the corner that it cuts off.

In our construction, if Y_i is not a triangle then Y_{i+1} has strictly fewer corner cutting edges than Y_i . Since Y_0 has at most 3 corner cutting edges, the procedure will terminate for some $n \leq 3$.




FIGURE 5. A corner cutting edge e with vertices w_1 and w_2

SUBCASE 2.2.1: Y_i has no corner cutting edges.

In this case, we claim that the support of Y_i is triangular. First, if Y_i has no boundary edges, then its support is F . Suppose Y_i has some boundary edge e . If both vertices of e are carried by edges, then Y_i is on the same side as the vertex of F cut off by e . Any other boundary edge of Y_i would have to be corner cutting. Hence, the support of Y_i is the triangle cut off by e , and we are done.

Otherwise, e contains a vertex of F , and hence divides F into two triangles. Once again, any other boundary edge would have to be corner cutting, so the support of Y_i is a triangle. This proves the claim, and completes this subcase.

It remains to consider the situation where Y_i has a corner cutting edge e , with vertices w_1 and w_2 . Let v be the vertex of F cut off by e , as in Figure 5. We consider two further subcases, according to whether or not w is the root of C , as illustrated in Figure 6.




FIGURE 6. Possible locations for w , with the region Y_i shaded

SUBCASE 2.2.2: Y_i has a corner cutting edge, and w is not the root of C . If w is not the root, then it is carried by some 2-face of Δ other than F . It cannot be carried by the face opposite v , since then both of the edges $\text{Conv}\{w_1, w\}$ and $\text{Conv}\{w_2, w\}$ would be interior, meaning that w_1 and w_2 would both be low excess vertices on the same component of the internal edge graph. Therefore without loss of generality, we can assume that w is carried by the 2-face $F' \neq F$ containing w_2 .

Note that the support of Y_i is a polygon, and if we now think of its boundary not in F , but in the plane containing F , there are exactly two vertices of the boundary adjacent to w_1 ; one of these is w_2 , and the other is some vertex, which we call w_3 .

The edge $\text{Conv}\{w, w_3\}$ cannot be interior, since w_3 is contained in the boundary of F , and hence has low excess, but w_1 is the root of C . Thus $w_3 \in F \cap F'$, as shown in Figure 6a. It follows that the polygon Y_i both contains and is contained in the triangle bounded by $\text{Conv}\{w_1, w_2\}$, $\text{Conv}\{w_2, w_3\}$, and $\text{Conv}\{w_1, w_3\}$. Therefore, Y_i is the triangle $\text{Conv}\{w_1, w_2, w_3\}$, and we are done.

SUBCASE 2.2.3: Y_i has a corner cutting edge, and w is the root of C .

In this case, w cannot have an interior edge to either w_1 or w_2 , since w_1 and w_2 both have low excess, and w is the root of C . Therefore, if F_1 and F_2 are the faces other than F containing w_1 and w_2 respectively, then w must lie on the edge $F_1 \cap F_2$, possibly at the vertex, though this will not matter. This situation is illustrated in Figure 6b. Since $\text{Conv}\{w, w_1, w_2\}$ is in fact a 2-face of Γ , the subcomplex with support $\text{Conv}\{w, w_1, w_2, v\}$ satisfies the hypotheses of Lemma 4.1. Thus we may assume that it is trivially subdivided. We then set Y_{i+1} to be the union of Y_i with the triangle $\text{Conv}\{w_1, w_2, v\}$. Then Y_{i+1} satisfies properties (1) and (2), and has fewer corner cutting edges than Y_i . Repeating this procedure, as Y_0 has at most three corner cutting edges, we eventually arrive at Y_n satisfying properties (1)–(3), and the theorem follows. \square

5. HIGHER DIMENSIONS

While Theorem 1.3, which is an immediate consequence of Proposition 3.4, says that the structure of the internal edge graph is essentially the same in higher dimensions, it seems that this is not sufficient for a useful classification of triangulations with vanishing local h -polynomial. Indeed, there are infinitely many different triangulations of the 4-simplex with empty internal edge graph. For instance, the join of the trifold with an arbitrary subdivision of the 1-simplex has this property. These triangulations give rise to the following negative result.

PROPOSITION 5.1. *There is no finite collection of triangulations of the 4-simplex with vanishing local h -polynomials from which all others can be obtained by a series of conical facet refinements.*

Proof. Let Γ_n be the subdivision of the 1-simplex with n interior vertices, and let Γ' be the trifold. If e is an edge of Γ_n such that both endpoints are interior, then the join of e with the center facet of Γ' is not a pyramid. Hence $\Gamma_n * \Gamma'$ is a triangulation with vanishing local h -polynomial that has $n - 1$ non-pyramid facets. The proposition follows, since, as discussed in Remark 3.7, all of the new facets introduced by a conical facet refinement are pyramids. \square

We leave the problem of classifying triangulations with vanishing local h -polynomial in dimensions 4 and higher open, for future research.

Acknowledgements. This project was carried out as part of the SUMRY undergraduate research program. We thank the other participants and mentors in this program

for their support and encouragement. We are also grateful to Matt Larson for helpful conversation related to this project.

REFERENCES

- [1] Christos A. Athanasiadis, *A survey of subdivisions and local h -vectors*, in The mathematical legacy of Richard P. Stanley, Amer. Math. Soc., Providence, RI, 2016, pp. 39–51.
- [2] Christos A. Athanasiadis and Christina Savvidou, *The local h -vector of the cluster subdivision of a simplex*, Sémin. Lothar. Combin. **66** (2011/12), Art. B66c (21 pages).
- [3] Mark A. de Cataldo, Luca Migliorini, and Mircea Mustață, *Combinatorics and topology of proper toric maps*, J. Reine Angew. Math. **744** (2018), 133–163.
- [4] Jan Denef and François Loeser, *Motivic Igusa zeta functions*, J. Algebraic Geom. **7** (1998), no. 3, 505–537.
- [5] Jun-ichi Igusa, *Complex powers and asymptotic expansions. II. Asymptotic expansions*, J. Reine Angew. Math. **278/279** (1975), 307–321.
- [6] Eric Katz and Alan Stapledon, *Local h -polynomials, invariants of subdivisions, and mixed Ehrhart theory*, Adv. Math. **286** (2016), 181–239.
- [7] Richard P. Stanley, *Subdivisions and local h -vectors*, J. Amer. Math. Soc. **5** (1992), no. 4, 805–851.
- [8] Alan Stapledon, *Formulas for monodromy*, Res. Math. Sci. **4** (2017), 8 (42 pages).

ANDRÉ DE MOURA, Viela do Mato 4 BL A RC Esq, Quinta da Beloura, 2710-695, Sintra, Portugal
E-mail : andros.moura@gmail.com

ELIJAH GUNTHER, Department of Mathematics, David Rittenhouse Lab, 209 South 33rd Street,
Philadelphia, PA 19104-6395, USA
E-mail : elijahg@sas.upenn.edu

SAM PAYNE, UT Austin, Department of Mathematics, 2515 Speedway, PMA 8.100, Austin, TX
78722, USA
E-mail : sampayne@utexas.edu

JASON SCHUCHARDT, UCLA Department of Mathematics, Math. Sciences Building 6363, 520
Portola Plaza, Los Angeles, CA 90095, USA
E-mail : jasonscho@g.ucla.edu

ALAN STAPLEDON
E-mail : astaplndn@gmail.com