



ALGEBRAIC COMBINATORICS

Xiaofeng Gu & Willem H. Haemers

Graph toughness from Laplacian eigenvalues

Volume 5, issue 1 (2022), p. 53-61.

<https://doi.org/10.5802/alco.197>

© The journal and the authors, 2022.




This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing
www.tccpublishing.org www.centre-mersenne.org





Graph toughness from Laplacian eigenvalues

Xiaofeng Gu & Willem H. Haemers

ABSTRACT The toughness $t(G)$ of a graph $G = (V, E)$ is defined as $t(G) = \min \left\{ \frac{|S|}{c(G-S)} \right\}$, in which the minimum is taken over all $S \subset V$ such that $G - S$ is disconnected, where $c(G - S)$ denotes the number of components of $G - S$. We present two tight lower bounds for $t(G)$ in terms of the Laplacian eigenvalues and provide strong support for a conjecture for a better bound which, if true, implies both bounds, and improves and generalizes known bounds by Alon, Brouwer, and the first author. As applications, several new results on perfect matchings, factors and walks from Laplacian eigenvalues are obtained, which leads to a conjecture about Hamiltonicity and Laplacian eigenvalues.

1. INTRODUCTION

Throughout this paper, $G = (V, E)$ is a simple graph of order n with nonempty vertex set V and nonempty edge set E . The minimum degree of G is denoted by δ . For a subset $S \subset V$, the subgraph of G induced by $V \setminus S$ is denoted by $G - S$, and $c(G - S)$ is the number of components of $G - S$.

The *toughness* $t(G)$ of a graph G is defined as $t(G) = \min \left\{ \frac{|S|}{c(G-S)} \right\}$, where the minimum is taken over all proper subsets $S \subset V$ such that $c(G - S) > 1$. By convention, a complete graph has infinite toughness. For any real number $r \geq 0$, G is *r-tough* if $t(G) \geq r$. The graph toughness was introduced by Chvátal [11] in 1973 and has been shown closely related to many graph properties, including connectivity, Hamiltonicity, pancyclicity, factors, spanning trees, and others (see [4]).

Toughness of regular graphs from eigenvalues of adjacency matrices has been well studied by, among others, [1, 6, 15, 16, 24, 25]. We use $\lambda_i := \lambda_i(G)$ to denote the i -th largest eigenvalue of the adjacency matrix of G , and let $\lambda = \max\{|\lambda_2|, |\lambda_n|\}$, that is, λ is the second largest absolute eigenvalue. It was Alon [1] who first showed that for any connected d -regular graph G , $t(G) > \frac{1}{3} \left(\frac{d^2}{d\lambda + \lambda^2} - 1 \right)$, through which, Alon was able to show that for every t and g there are t -tough graphs of girth strictly greater than g . This strengthened a result of Bauer, Van den Heuvel and Schmeichel [5] who showed the same for $g = 3$, and thus disproved in a strong sense a conjecture of Chvátal [11] that there exists a constant t_0 such that every t_0 -tough graph is pancyclic. Almost at the same time, Brouwer [6] independently showed that $t(G) > \frac{d}{\lambda} - 2$ for any connected d -regular graph G . He also conjectured that $t(G) \geq \frac{d}{\lambda} - 1$ in [6, 7]. Recently Brouwer's

Manuscript received 8th April 2021, revised 24th August 2021, accepted 25th August 2021.

KEYWORDS. Toughness, Laplacian eigenvalue, perfect matching, factor, Hamilton cycle.

ACKNOWLEDGEMENTS. The first author is partially supported by a grant from the Simons Foundation (522728, XG).

conjecture has been confirmed by the first author [24]. In this paper, we consider arbitrary graphs and look for lower bounds on $t(G)$ in terms of the eigenvalues of the Laplacian matrix.

The *Laplacian matrix* L (also called *combinatorial Laplacian* or *discrete Laplacian*) of a graph G , is defined by $L = D - A$, where D is the diagonal degree matrix and A is the adjacency matrix of G . Let $\mu_i := \mu_i(G)$ denote the i -th smallest eigenvalue of the Laplacian matrix of G . Then L is positive semi-definite and $\mu_1(G) = 0$. The second smallest Laplacian eigenvalue $\mu_2(G)$, is known as the *algebraic connectivity* of G . We have $\mu_2(G) = 0$ if and only if G is disconnected. Moreover, if κ is the vertex connectivity, then

$$(1) \quad \mu_2 \leq \kappa \leq \delta.$$

The complement $\overline{G}$ of G has eigenvalues $\mu_1(\overline{G}) = 0$ and $\mu_i(\overline{G}) = n - \mu_{n+2-i}(G)$ for $i = 2, \dots, n$. Therefore $\mu_n(G) \leq n$ and $\mu_n(G) = n$ if and only if $\overline{G}$ is disconnected. If G is regular of degree d , then $L = dI - A$ and therefore $\mu_i = d - \lambda_i$ for $i = 1, \dots, n$. For these and other properties of the Laplacian matrix and its eigenvalues, we refer to [9].

The paper is organized as follows. Our main results will be in the next section. The tools and the proofs will be presented in Sections 3 and 4. In Section 5, we will show applications on perfect matchings, factors and walks. Since toughness is closely related to cycle structures, we include a conjecture on Hamilton cycles from Laplacian eigenvalues.

2. RESULTS

Recently, the second author made the following conjecture [28].

CONJECTURE 2.1 (Haemers).

$$(2) \quad t(G) \geq \frac{\mu_2}{\mu_n - \delta}.$$

For d -regular graphs, this conjecture implies that $t(G) \geq \frac{d - \lambda_2}{-\lambda_n}$, which is stronger than Brouwer's conjecture. The conjecture is supported by the following theorem and proposition. The proofs will be given in Section 4.

THEOREM 2.2.

$$(3) \quad t(G) \geq \frac{\mu_n \mu_2}{n(\mu_n - \delta)},$$

and

$$(4) \quad t(G) \geq \frac{\mu_2}{\mu_n - \mu_2}.$$

PROPOSITION 2.3. *Let $S \subset V$ be such that $t(G) = |S|/c(G - S)$. Then Conjecture 2.1 is true in each of the following cases.*

- (i) *The complement of G is disconnected,*
- (ii) *All connected components of $G - S$ are singletons (i.e. $n - |S| = c(G - S)$),*
- (iii) *The union of some components of $G - S$ has order $\frac{1}{2}(n - |S|)$,*
- (iv) *$c(G - S) = 2$.*

Since $\mu_n \leq n$ and $\delta \geq \mu_2$, the conjectured bound (2) implies (3) and (4). Note that (2) and (3) coincide if $\mu_n = n$, that is, if the complement of G is disconnected. Therefore (i) of Proposition 2.3 follows from (3).

The three bounds coincide and are tight in case G is the complete multipartite graph $K_{n_1, \dots, n_m}$ ($1 < m < n$). Indeed, assume $n_1 \geq \dots \geq n_m$ then $t(G) = (n - n_1)/n_1$, $\mu_n = n$ and $\mu_2 = \delta = n - n_1$.

The bounds (3) and (4) are incomparable. For example, when G is the Petersen graph, then $t(G) = 4/3$ and $\mu_2 = 2$, $\mu_{10} = 5$ and $\delta = 3$, and so (2), (3), and (4) give $t(G) \geq 1$, $1/2$ and $2/3$, respectively. The complement $\overline{G}$ satisfies $t(\overline{G}) = 3$ and $\mu_2(\overline{G}) = 5$, $\mu_{10}(\overline{G}) = 8$ and $\delta = 6$, and the bounds (2), (3), and (4) give $t(\overline{G}) \geq 5/2$, 2 and $5/3$ respectively.

3. TOOLS

The following separation inequality from [27] provides a bridge between graph parameters and Laplacian eigenvalues. It can also be found in [9, Proposition 4.8.1].

THEOREM 3.1 ([27]). *Suppose that X and Y are two disjoint subsets of V such that there is no edge between X and Y . Then*

$$(5) \quad \frac{|X||Y|}{(n - |X|)(n - |Y|)} \leq \left(\frac{\mu_n - \mu_2}{\mu_n + \mu_2} \right)^2.$$

By the above separation inequality, a simple yet useful proposition has been proved in [26].

PROPOSITION 3.2 ([26]). *Let $S \subset V$ such that $G - S$ is disconnected. Let X and Y be disjoint vertex subsets of $V \setminus S$ such that $X \cup Y = V \setminus S$ with $|X| \leq |Y|$. Then*

$$(6) \quad |X| \leq \frac{\mu_n - \mu_2}{2\mu_n} \cdot n,$$

and

$$(7) \quad |S| \geq \frac{2\mu_2}{\mu_n - \mu_2} \cdot |X|,$$

with each equality holding only when $|X| = |Y|$.

For the purpose of completeness, we include a proof that was originally given in [26].

Proof. By (5), we have

$$(8) \quad |X||Y| \leq \left(\frac{\mu_n - \mu_2}{\mu_n + \mu_2} \right)^2 (n - |X|)(n - |Y|),$$

Let $\beta = \frac{\mu_n - \mu_2}{\mu_n + \mu_2}$. Then $0 < \beta \leq 1$, as G contains at least one edge and $G - S$ is disconnected. It follows that

$$|X|^2 \leq |X| \cdot |Y| \leq \beta^2 (n - |X|)(n - |Y|) \leq \beta^2 (n - |X|)^2,$$

that is

$$|X| \leq \beta(n - |X|),$$

and hence

$$(9) \quad |X| \leq \frac{\beta n}{1 + \beta} = \frac{\mu_n - \mu_2}{2\mu_n} \cdot n,$$

with the equality holding only when $|X| = |Y|$.

Also, since $|Y| = n - |S| - |X|$, by (8), we have

$$|X|(n - |S| - |X|) = |X| \cdot |Y| \leq \beta^2 (n - |X|)(n - |Y|) = \beta^2 (n - |X|)(|S| + |X|),$$

implying that

$$(10) \quad |X|n \leq (\beta^2 (n - |X|) + |X|)(|S| + |X|) = (\beta^2 n + (1 - \beta^2)|X|)(|S| + |X|).$$

By (9), we have

$$(1 - \beta^2)|X| \leq (1 - \beta^2) \cdot \frac{\beta n}{1 + \beta} = (\beta - \beta^2)n,$$

which, together with (10), implies that

$$|X|n \leq (\beta^2 n + (\beta - \beta^2)n)(|S| + |X|) = \beta n(|S| + |X|),$$

and we have

$$|X| \leq \beta(|S| + |X|).$$

Hence,

$$|S| \geq \frac{1-\beta}{\beta}|X| = \frac{2\mu_2}{\mu_n - \mu_2} \cdot |X|.$$

Since (9) was utilized, the equality holds in (7) only when $|X| = |Y|$. $\square$

Generalizing Hoffman's ratio bound, the following bound for the cardinality of an independent set of an arbitrary graph has been obtained independently in [21] and in [38] (see also [29]).

THEOREM 3.3 (Godsil and Newman [21], Lu, Liu and Tian [38]). *If U is an independent set of G , then*

$$|U| \leq \frac{\mu_n - \delta}{\mu_n} \cdot n.$$

4. PROOFS

Throughout this section we take $S \subset V$ such that $t(G) = |S|/c(G - S)$, and put $c = c(G - S)$. Clearly the vertex connectivity κ of G satisfies $\kappa \leq |S|$, and so by (1), we have

$$(11) \quad |S| \geq \mu_2.$$

First we prove (3) of Theorem 2.2.

Proof. By taking a vertex in each component of $G - S$ we obtain an independent set of cardinality c . Therefore Theorem 3.3 gives $c \leq n(\mu_n - \delta)/\mu_n$, and so by (11),

$$t(G) = \frac{|S|}{c} \geq \frac{\mu_n \mu_2}{n(\mu_n - \delta)}. \quad \square$$

For convenience we continue with the proof of (ii) of Proposition 2.3.

Proof. If $n - |S| = c$, then $V \setminus S$ is an independent set of G . By use of Theorem 3.3 and (1) we have

$$t(G) = \frac{|S|}{c} \geq \frac{n-c}{c} = \frac{n}{c} - 1 \geq \frac{\mu_n}{\mu_n - \delta} - 1 = \frac{\delta}{\mu_n - \delta} \geq \frac{\mu_2}{\mu_n - \delta}. \quad \square$$

Next we prove (4) of Theorem 2.2.

Proof. Let $H_1, H_2, \dots, H_c$ be the vertex sets of the components of $G - S$. Without loss of generality, suppose that $|H_1| \leq |H_2| \leq \dots \leq |H_c|$. Above we proved that (2) and therefore (4) holds if $H_1, \dots, H_c$ are singletons. Thus, we may assume that $n - |S| \geq c + 1$. We claim that $H_1, H_2, \dots, H_c$ can be partitioned into two sets X and Y such that $|Y| \geq |X| \geq c/2$. If c is even, we can simply define $X = \bigcup_{1 \leq i \leq \lfloor c/2 \rfloor} H_i$ and $Y = (V \setminus S) \setminus X$. Now we assume c is odd. If $|H_{(c-1)/2}| \geq 2$, then define $X = \bigcup_{1 \leq i \leq (c-1)/2} H_i$ and $Y = (V \setminus S) \setminus X$ as needed. The remaining case is $|H_1| = \dots = |H_{(c-1)/2}| = 1$. We can define $X = \bigcup_{1 \leq i \leq (c+1)/2} H_i$ and $Y = (V \setminus S) \setminus X$, and we need to show that $|Y| \geq |X| \geq c/2$. If $|H_{(c+1)/2}| = 1$, then $|X| = \frac{c+1}{2}$ and $|Y| = n - |S| - |X| \geq \frac{c+1}{2}$, since $n - |S| \geq c + 1$. If $|H_{(c+1)/2}| \geq 2$, then $|X| = \frac{c-1}{2} + |H_{(c+1)/2}| \geq \frac{c-1}{2} + 2 > \frac{c}{2}$ and $|Y| = \sum_{i > (c+1)/2} |H_i| \geq 2 \cdot \frac{c-1}{2} = c - 1 \geq c/2$. Switch X and Y whenever needed to get $|Y| \geq |X|$.

It follows that $c \leq 2|X|$. Thus, by (7),

$$t(G) = \frac{|S|}{c} \geq \frac{2\mu_2}{\mu_n - \mu_2} \cdot \frac{|X|}{c} \geq \frac{\mu_2}{\mu_n - \mu_2}. \quad \square$$

The last two proofs of this section deal with (iii) and (iv) of Proposition 2.3.

Proof. (iii): In this case $V \setminus S$ can be partitioned into two sets X and Y both having cardinality $\frac{1}{2}(n - |S|)$, such that there are no edges between X and Y . We apply (7) and find $|S| \geq \frac{\mu_2}{\mu_n - \mu_2}(n - |S|)$, which implies $|S| \geq n\mu_2/\mu_n$. As before, Theorem 3.3 gives $c \leq n(\mu_n - \delta)/\mu_n$ and hence

$$t(G) = \frac{|S|}{c} \geq n \frac{\mu_2}{\mu_n} \cdot \frac{\mu_n}{n(\mu_n - \delta)} = \frac{\mu_2}{\mu_n - \delta}. \quad \square$$

Proof. (iv): It is known (see [9, Section 3.9]) that $\mu_n \geq d_{\max} + 1$, when $d_{\max}$ is the maximum degree of G . If G is not regular, then $d_{\max} - \delta \geq 1$, and hence $\mu_n - \delta \geq 2$. If G is regular of degree $d = \delta$, then the adjacency matrix has smallest eigenvalue $\lambda_n = d - \mu_n$. If G is regular with $\lambda_n > -2$ then G is the complete graph K_n or an odd cycle C_n ; see [17, Theorem 2.5]. We have $t(K_n) = \infty$ and $t(C_n) = 2$ if $n \geq 4$. If n is odd, $\mu_2(C_n) = 2 - 2\cos(\pi/n)$ and $\mu_n(C_n) = 2 + 2\cos(2\pi/n)$, so (2) gives $2 \geq (1 - \cos(\pi/n))/\cos(2\pi/n)$ which is clearly true for all odd $n \geq 5$. Thus we can assume that $\lambda_n \leq -2$ and hence $\mu_n - d = \mu_n - \delta \geq 2$. By (11), it implies that

$$t(G) = \frac{|S|}{c} = \frac{|S|}{2} \geq \frac{\mu_2}{2} \geq \frac{\mu_2}{\mu_n - \delta}. \quad \square$$

5. APPLICATIONS

It was shown in [34] that if $\frac{\mu_2}{\mu_n} \geq \frac{2}{3}$, then G is 2-tough. Now we can generalize it by rewriting (4) of Theorem 2.2 as below.

THEOREM 5.1. *If $\frac{\mu_2}{\mu_n} \geq \frac{r}{r+1}$, then G is r -tough.*

Since many graph parameters and properties are related to toughness, we have various applications, including but not limited to the results in this section. We refer readers to the survey paper [4] for more toughness related results.

Spectral conditions for matchings and k -factors of regular graphs have been attracting many researchers [8, 12–14, 23, 33, 36, 37, 39], among others. However, not as much has been discovered for general graphs that are not necessarily regular. Brouwer and the second author [8] showed that if n is even and $2\mu_2 \geq \mu_n$, then G has a perfect matching. This result has been recently generalized to matching numbers in [26]. We will have other generalizations by using graph toughness.

A graph G is called *elementary* if it contains a perfect matching and if the edges which occur in at least one perfect matching in G induce a connected subgraph. A substantial study of elementary graphs has been given in [35]. It is proved in [3] that every 1-tough graph with an even number of vertices is elementary. Thus Theorem 5.1 implies the following result.

THEOREM 5.2. *If n is even and $2\mu_2 \geq \mu_n$, then G is elementary.*

Let G be an n -vertex graph with a perfect matching, and m be a positive integer with $m < n/2 - 1$. Then G is called *m -extendable* if every matching of size m extends to a perfect matching. Plummer [40] proved that if $t(G) > m$, then G is m -extendable.

THEOREM 5.3. *Suppose n is even, and let m be a positive integer such that $m < n/2 - 1$. If*

$$\frac{\mu_2}{\mu_n} > \frac{m}{m+1},$$

then G is m -extendable.

In [19], it is proved that every k -tough graph has a k -factor if $k|V(G)|$ is even and $|V(G)| \geq k + 1$, confirming a conjecture of Chvátal [11]. The result was extended to non-regular factors by Katerinis [31]. Let $a \leq b$ be positive integers. An $[a, b]$ -factor of a graph G is a spanning subgraph H such that $a \leq d_H(v) \leq b$ for each vertex. Katerinis [31] showed that for a graph G on n vertices such that $a < b$ or bn is even, if $t(G) \geq a + \frac{a}{b} - 1$, then G has an $[a, b]$ -factor. By using Theorem 5.1, we have the following Laplacian eigenvalue condition for the existence of factors.

THEOREM 5.4. *Let $a \leq b$ be positive integers such that $a < b$ or bn is even. If*

$$\frac{\mu_2}{\mu_n} \geq 1 - \frac{b}{a(b+1)},$$

then G has a $[a, b]$ -factor.

The following corollary on k -factors is a generalization of the result on perfect matching by Brouwer and the second author [8].

COROLLARY 5.5. *Let k be a positive integer such that $n \geq k + 1$ and kn is even. If*

$$\frac{\mu_2}{\mu_n} \geq \frac{k}{k+1},$$

then G has a k -factor.

A graph G is (k, s) -factor-critical if $G - X$ has a k -factor for all $X \subseteq V(G)$ with $|X| = s$. A $(1, 1)$ -factor-critical graph is usually referred to as a factor-critical graph. It is proved in [3] that every 1-tough graph with an odd number of vertices is factor-critical. By Theorem 5.1, we have the following result on factor-critical graphs from Laplacian eigenvalues, which was originally obtained in [26].

THEOREM 5.6 ([26]). *If n is odd and $2\mu_2 \geq \mu_n$, then G is $(1, 1)$ -factor-critical.*

For $2 \leq s < n$, it was shown by Favaron [20] that for a graph G on n vertices with $n + s$ even, if $t(G) > s/2$, then G is $(1, s)$ -factor-critical. By Theorem 5.1, we have the following result.

THEOREM 5.7. *Suppose $2 \leq s < n$ and $n + s$ is even. If*

$$\frac{\mu_2}{\mu_n} > \frac{s}{s+2},$$

then G is $(1, s)$ -factor-critical.

For $k = 2, 3$ or even a general k , similar results on (k, s) -factor-critical graphs from toughness can be found in the survey [4]. Thus Theorem 5.1 implies Laplacian eigenvalue conditions for (k, s) -factor-critical graphs with various values of k and s , which will be omitted here.

Spectral conditions for the existence of a spanning tree with degree bounded above by a fixed number k in a regular graph have been obtained in [15, 16]. When $k = 2$, such a spanning tree is exactly a Hamilton path, and a sufficient condition has been given by, among others, Butler and Chung [10], who borrowed the idea from [32] (both [10, 32] studied a stronger structure, i.e. Hamilton cycle). The case of $k \geq 3$ for general graphs was asked in [15] and has been solved in [26]. A theorem of Win [41] implies that if $t(G) \geq \frac{1}{k-2}$ for $k \geq 3$, then G has a spanning tree with maximum degree at most k . Thus, Theorem 5.1 implies the following result of [26].

THEOREM 5.8 ([26]). Let $k \geq 3$ be an integer. If

$$\frac{\mu_2}{\mu_n} \geq \frac{1}{k-1},$$

then G has a spanning tree with maximum degree at most k .

Generalizing the idea of a Hamilton cycle, a k -walk in a graph G is a closed spanning walk of G that visits every vertex of G at most k times. In particular, a Hamilton cycle is a 1-walk. Jackson and Wormald [30] observed that the existence of a spanning tree with maximum degree at most k actually implies the existence of a k -walk. Thus Theorem 5.8 implies the existence of a k -walk for $k \geq 3$. For $k = 2$, Ellingham and Zha [18] showed that every 4-tough graph has a 2-walk. By Theorem 5.1, we have the following Laplacian eigenvalue condition for the existence of a 2-walk.

THEOREM 5.9. If $\frac{\mu_2}{\mu_n} \geq \frac{4}{5}$, then G has a 2-walk.

REMARK 5.10. In this section we presented applications of (4) on perfect matchings, factors and walks. The bound (3) has similar applications and if Conjecture 2.1 is true, then all results in this section can be improved in a similar manner.

Similar to Theorems 5.8 and 5.9, the first author made the following conjecture for Hamilton cycles, but never published elsewhere before.

CONJECTURE 5.11 (Gu). There exists a positive constant $C < 1$ such that if $\mu_2/\mu_n \geq C$ and $n \geq 3$, then G contains a Hamilton cycle.

REMARK 5.12. Notice that $K_{s,s+1}$ contains no Hamilton cycle, but $\mu_2 = s$ and $\mu_n = 2s + 1$, which implies that μ_2/μ_n can be arbitrarily close to $1/2$ if s is sufficiently large. Thus the smallest possible C is at least $1/2$.

Chvátal [11] conjectured that there exists a constant t_0 such that every t_0 -tough graph contains a Hamilton cycle. By (4) of Theorem 2.2, clearly Chvátal's conjecture implies Conjecture 5.11. Krivelevich and Sudakov [32] conjectured that for a d -regular graph G , there exists a constant K such that $d/\lambda > K$ and n is large enough, then G contains a Hamilton cycle. It is not hard to see that Conjecture 5.11 implies the conjecture of Krivelevich and Sudakov. All three conjectures remain open.

The results in this section as well as [8, 26, 27, 42] indicate that many graph properties are related to the Laplacian eigenratio μ_2/μ_n . This eigenratio also has application aspects, and is highly related to the synchronization in small-world systems [2]. It plays a similar role as spectral gap, but may give a bit more information about the structure of the graph. It is clear that $\mu_2/\mu_n = 1$ if and only if the graph is complete. For a non-complete graph we have that $\mu_n \geq 1 + d_{\max}$ (see Section 4), and since $\mu_2 \leq \kappa$ the Laplacian eigenratio of a non-complete graph is bounded above by $\kappa/(1 + d_{\max})$ (see [22] for more about this inequality). But for many d -regular graphs with vertex connectivity d (i.e. $d_{\max} = \kappa$) the ratio is close to 1. For example the Paley graph of order n satisfies $\mu_2/\mu_n = (\sqrt{n} - 1)/(\sqrt{n} + 1)$ (see [9]). We feel that the Laplacian eigenratio is interesting for future research.

Acknowledgements. The authors would like to thank the referees for carefully reading our manuscript and for their comments.

REFERENCES

- [1] Noga Alon, *Tough Ramsey graphs without short cycles*, J. Algebraic Combin. **4** (1995), no. 3, 189–195.
- [2] Mauricio Barahona and Louis M. Pecora, *Synchronization in small-world systems*, Physical Review Letters **89** (2002), no. 5, Paper no. 054101.

- [3] Douglas Bauer, Haitze J. Broersma, Nathan Kahl, Aurora Morgana, Edward Schmeichel, and T. Surowiec, *Tutte sets in graphs. II. The complexity of finding maximum Tutte sets*, Discrete Appl. Math. **155** (2007), no. 10, 1336–1343.
- [4] Douglas Bauer, Hajo Broersma, and Edward Schmeichel, *Toughness in graphs—a survey*, Graphs Combin. **22** (2006), no. 1, 1–35.
- [5] Douglas Bauer, Jan Vandenheuevel, and Edward Schmeichel, *Toughness and triangle-free graphs*, J. Combin. Theory Ser. B **65** (1995), no. 2, 208–221.
- [6] Andries E. Brouwer, *Toughness and spectrum of a graph*, Linear Algebra Appl. **226/228** (1995), 267–271.
- [7] ———, *Spectrum and connectivity of graphs*, CWI Quarterly **9** (1996), no. 1-2, 37–40, SMC 50 jubilee (Amsterdam, 1996).
- [8] Andries E. Brouwer and Willem H. Haemers, *Eigenvalues and perfect matchings*, Linear Algebra Appl. **395** (2005), 155–162.
- [9] ———, *Spectra of graphs*, Universitext, Springer, New York, 2012.
- [10] Steve Butler and Fan Chung, *Small spectral gap in the combinatorial Laplacian implies Hamiltonian*, Ann. Comb. **13** (2010), no. 4, 403–412.
- [11] Vasek Chvátal, *Tough graphs and Hamiltonian circuits*, Discrete Math. **5** (1973), 215–228.
- [12] Sebastian M. Cioabă, *Perfect matchings, eigenvalues and expansion*, C. R. Math. Acad. Sci. Soc. R. Can. **27** (2005), no. 4, 101–104.
- [13] Sebastian M. Cioabă and David A. Gregory, *Large matchings from eigenvalues*, Linear Algebra Appl. **422** (2007), no. 1, 308–317.
- [14] Sebastian M. Cioabă, David A. Gregory, and Willem H. Haemers, *Matchings in regular graphs from eigenvalues*, J. Combin. Theory Ser. B **99** (2009), no. 2, 287–297.
- [15] Sebastian M. Cioabă and Xiaofeng Gu, *Connectivity, toughness, spanning trees of bounded degree, and the spectrum of regular graphs*, Czechoslovak Math. J. **66(141)** (2016), no. 3, 913–924.
- [16] Sebastian M. Cioabă and Wiseley Wong, *The spectrum and toughness of regular graphs*, Discrete Appl. Math. **176** (2014), 43–52.
- [17] Michael Doob and Dragoš Cvetković, *On spectral characterizations and embeddings of graphs*, Linear Algebra Appl. **27** (1979), 17–26.
- [18] Mark N. Ellingham and Xiaoya Zha, *Toughness, trees, and walks*, J. Graph Theory **33** (2000), no. 3, 125–137.
- [19] Hikoe Enomoto, Bill Jackson, Panagiotis Katerinis, and Akira Saito, *Toughness and the existence of k -factors*, J. Graph Theory **9** (1985), no. 1, 87–95.
- [20] Odile Favaron, *On k -factor-critical graphs*, Discuss. Math. Graph Theory **16** (1996), no. 1, 41–51.
- [21] Chris D. Godsil and Mike W. Newman, *Eigenvalue bounds for independent sets*, J. Combin. Theory Ser. B **98** (2008), no. 4, 721–734.
- [22] Felix Goldberg, *Bounding the gap between extremal Laplacian eigenvalues of graphs*, Linear Algebra Appl. **416** (2006), no. 1, 68–74.
- [23] Xiaofeng Gu, *Regular factors and eigenvalues of regular graphs*, European J. Combin. **42** (2014), 15–25.
- [24] ———, *A proof of Brouwer’s toughness conjecture*, SIAM J. Discrete Math. **35** (2021), no. 2, 948–952.
- [25] ———, *Toughness in pseudo-random graphs*, European J. Combin. **92** (2021), Paper no. 103255 (11 pages).
- [26] Xiaofeng Gu and Muhuo Liu, *A tight lower bound on the matching number of graphs via Laplacian eigenvalues*, European J. Combin. **101** (2022), Paper no. 103468.
- [27] Willem H. Haemers, *Interlacing eigenvalues and graphs*, Linear Algebra Appl. **226/228** (1995), 593–616.
- [28] ———, *Toughness conjecture*, <https://www.researchgate.net/publication/348437253>, 2020.
- [29] ———, *Hoffman’s ratio bound*, Linear Algebra Appl. **617** (2021), 215–219.
- [30] Bill Jackson and Nicholas Charles Wormald, *k -walks in graphs*, Australasian Journal of Combinatorics **2** (1990), 135–146.
- [31] Panagiotis Katerinis, *Toughness of graphs and the existence of factors*, Discrete Math. **80** (1990), no. 1, 81–92.
- [32] Michael Krivelevich and Benny Sudakov, *Sparse pseudo-random graphs are Hamiltonian*, J. Graph Theory **42** (2003), no. 1, 17–33.
- [33] ———, *Pseudo-random graphs*, in More sets, graphs and numbers, Bolyai Soc. Math. Stud., vol. 15, Springer, Berlin, 2006, pp. 199–262.

- [34] Bolian Liu and Siyuan Chen, *Algebraic conditions for t -tough graphs*, Czechoslovak Math. J. **60(135)** (2010), no. 4, 1079–1089.
- [35] László Lovász and Michael D. Plummer, *Matching theory. Annals of Discrete Mathematics*, 29, North-Holland Mathematics Studies, vol. 121, North-Holland Publishing Co., Amsterdam, 1986.
- [36] Hongliang Lu, *Regular factors of regular graphs from eigenvalues*, Electron. J. Combin. **17** (2010), no. 1, Paper no. R159 (12 pages).
- [37] ———, *Regular graphs, eigenvalues and regular factors*, J. Graph Theory **69** (2012), no. 4, 349–355.
- [38] Mei Lu, Huiqing Liu, and Feng Tian, *Laplacian spectral bounds for clique and independence numbers of graphs*, J. Combin. Theory Ser. B **97** (2007), no. 5, 726–732.
- [39] Suil O and Sebastian M. Cioabă, *Edge-connectivity, eigenvalues, and matchings in regular graphs*, SIAM J. Discrete Math. **24** (2010), no. 4, 1470–1481.
- [40] Michael D. Plummer, *Toughness and matching extension in graphs*, Discrete Math. **72** (1988), no. 1-3, 311–320.
- [41] Sein Win, *On a connection between the existence of k -trees and the toughness of a graph*, Graphs Combin. **5** (1989), no. 2, 201–205.
- [42] Zhifu You and Bolian Liu, *On the Laplacian spectral ratio of connected graphs*, Appl. Math. Lett. **25** (2012), no. 10, 1245–1250.

XIAOFENG GU, Department of Computing and Mathematics, University of West Georgia, Carrollton, GA 30118, USA
E-mail : xgu@westga.edu

WILLEM H. HAEMERS, Department of Econometrics and Operations Research, Tilburg University, Tilburg, The Netherlands
E-mail : haemers@uvt.nl