



ALGEBRAIC COMBINATORICS

Yifeng Huang

Counting on the variety of modules over the quantum plane

Volume 5, issue 3 (2022), p. 583-592.

<https://doi.org/10.5802/alco.230>

© The journal and the authors, 2022.




This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing
www.tccpublishing.org www.centre-mersenne.org





Counting on the variety of modules over the quantum plane

Yifeng Huang

ABSTRACT Let ζ be a fixed nonzero element in a finite field $\mathbb{F}_q$ with q elements. In this article, we count the number of pairs (A, B) of $n \times n$ matrices over $\mathbb{F}_q$ satisfying $AB = \zeta BA$ by giving a generating function. This generalizes a generating function of Feit and Fine that counts pairs of commuting matrices. Our result can be also viewed as the point count of the variety of modules over the quantum plane $XY = \zeta YX$, whose geometry was described by Chen and Lu.

1. INTRODUCTION

1.1. MAIN RESULTS. Fix a nonzero element ζ in $\mathbb{F}_q$, the finite field with q elements. Let $\text{ord}(\zeta)$ denote the smallest positive integer m such that $\zeta^m = 1$ in $\mathbb{F}_q$. We define the set of $\mathbb{F}_q$ -points of the ζ -commuting variety to be

$$(1) \quad K_{\zeta,n}(\mathbb{F}_q) := \{(A, B) \in \text{Mat}_n(\mathbb{F}_q) \times \text{Mat}_n(\mathbb{F}_q) : AB = \zeta BA\}.$$

The ζ -commuting variety $K_{\zeta,n}$ can be viewed as the variety of n -dimensional modules over the algebra of the quantum plane, namely, the noncommutative associative algebra in variables X and Y such that $XY = \zeta YX$. The geometry of the ζ -commuting variety was studied by Chen and Lu [3], where explicit descriptions of its irreducible components and of a GIT quotient were given. The combinatorics of the ζ -commuting has also been studied when $\zeta = 1$: Feit and Fine [6] gave an explicit formula for the point count of the commuting variety (namely, $K_{1,n}$) over a finite field, and Bryan and Morrison [2] proved that the “same” formula computes the motivic class of the commuting variety (over $\mathbb{C}$) in the Grothendieck ring of varieties.

The focus of this paper is to count the cardinality of $K_{\zeta,n}(\mathbb{F}_q)$ for ζ in general. As a special case, the cardinality of $K_{1,n}(\mathbb{F}_q)$, the set of pairs of commuting matrices, was determined by Feit and Fine [6] in the form of a generating function. We give a generating function for $|K_{\zeta,n}(\mathbb{F}_q)|$ that generalizes the $\zeta = 1$ case.

THEOREM 1.1. *Let $m = \text{ord}(\zeta)$; in other words, ζ is a primitive m -th root of unity of $\mathbb{F}_q$. We have the following identity of power series in x :*

$$(2) \quad \sum_{n=0}^{\infty} \frac{|K_{\zeta,n}(\mathbb{F}_q)|}{(q^n - 1)(q^n - q) \cdots (q^n - q^{n-1})} x^n = \prod_{i=1}^{\infty} F_m(x^i; q),$$

where

$$(3) \quad F_m(x; q) := \frac{1 - x^m}{(1 - x)(1 - x^m q)} \cdot \frac{1}{(1 - x)(1 - xq^{-1})(1 - xq^{-2}) \cdots}.$$

Manuscript received 29th November 2021, revised and accepted 7th February 2022.

KEYWORDS. quantum plane, matrix equations, counting over finite fields, generating functions.

ACKNOWLEDGEMENTS. The author is supported by National Science Foundation grant DMS-1701576.

We note that $|K_{\zeta,n}(\mathbb{F}_q)|$ only depends on the order m of ζ . When $m = 1$, we recover the generating function given by Feit and Fine.

Theorem 1.1 is a direct consequence of the following result, which in itself can be viewed as a refinement of Theorem 1.1. We define

$$(4) \quad U_{\zeta,n}(\mathbb{F}_q) := \{(A, B) \in \text{Mat}_n(\mathbb{F}_q) \times \text{Mat}_n(\mathbb{F}_q) : AB = \zeta BA, A \text{ nonsingular}\},$$

and

$$(5) \quad N_{\zeta,n}(\mathbb{F}_q) := \{(A, B) \in \text{Mat}_n(\mathbb{F}_q) \times \text{Mat}_n(\mathbb{F}_q) : AB = \zeta BA, A \text{ nilpotent}\}.$$

When $\zeta = -1$, the variety $N_{-1,n}$ is the semi-nilpotent anti-commuting variety, whose irreducible components and their dimensions are explicitly described by Chen and Wang [4].

For reasons of brevity, we put $|\text{GL}_n(\mathbb{F}_q)|$ in place of $(q^n - 1)(q^n - q) \dots (q^n - q^{n-1})$ in the formulas below, noting that $|\text{GL}_n(\mathbb{F}_q)| = (q^n - 1)(q^n - q) \dots (q^n - q^{n-1})$.

THEOREM 1.2. *Let $m = \text{ord}(\zeta)$. We have the following identities of power series in x :*

(a)

$$\sum_{n=0}^{\infty} \frac{|K_{\zeta,n}(\mathbb{F}_q)|}{|\text{GL}_n(\mathbb{F}_q)|} x^n = \left(\sum_{n=0}^{\infty} \frac{|U_{\zeta,n}(\mathbb{F}_q)|}{|\text{GL}_n(\mathbb{F}_q)|} x^n \right) \left(\sum_{n=0}^{\infty} \frac{|N_{\zeta,n}(\mathbb{F}_q)|}{|\text{GL}_n(\mathbb{F}_q)|} x^n \right)$$

(b)

$$\sum_{n=0}^{\infty} \frac{|U_{\zeta,n}(\mathbb{F}_q)|}{|\text{GL}_n(\mathbb{F}_q)|} x^n = \prod_{i=1}^{\infty} G_m(x^i; q),$$

where

$$G_m(x; q) := \frac{1 - x^m}{(1 - x)(1 - x^m q)}.$$

(c)

$$\sum_{n=0}^{\infty} \frac{|N_{\zeta,n}(\mathbb{F}_q)|}{|\text{GL}_n(\mathbb{F}_q)|} x^n = \prod_{i=1}^{\infty} H(x^i; q),$$

where

$$H(x; q) := \frac{1}{(1 - x)(1 - xq^{-1})(1 - xq^{-2}) \dots}.$$

Using Theorem 1.2(a), Theorem 1.1 follows from the observation $F_m(x; q) = G_m(x; q)H(x; q)$.

Note that Theorem 1.2(c) implies that $|N_{\zeta,n}(\mathbb{F}_q)|$ does not depend on m or ζ , as long as $\zeta \neq 0$. In particular, $|N_{\zeta,n}(\mathbb{F}_q)|$ always equals $|N_{1,n}(\mathbb{F}_q)|$, which was known to Feit and Fine. Therefore, the nontrivial dependence of $|K_{\zeta,n}(\mathbb{F}_q)|$ on ζ stems purely from that of $|U_{\zeta,n}(\mathbb{F}_q)|$.

1.2. HISTORY AND RELATED WORK. An important starting case in the study of varieties of modules is the commuting variety $K_{1,n} = \{(A, B) : A, B \in \text{Mat}_n, AB = BA\}$. The commuting variety over $\mathbb{C}$ was shown to be irreducible by Gerstenhaber [10] and Motzkin and Taussky [15]. Its point count was given by Feit and Fine [6]. This result was reproved by Bryan and Morrison [2] from the perspective of motivic Donaldson–Thomas theory.

The commuting variety can be viewed in the context of Lie algebras. Let $(\mathfrak{g}, [\cdot, \cdot])$ be a Lie algebra over an algebraically closed field. Define the *commuting variety* of $\mathfrak{g}$ as

$$(6) \quad C(\mathfrak{g}) := \{(x, y) \in \mathfrak{g} \times \mathfrak{g} : [x, y] = 0\},$$

then $K_{1,n}$ is the commuting variety of the Lie algebra of n by n matrices. As a generalization of the irreducibility result of $K_{1,n}$, Richardson [16] showed that the commuting

variety of any reductive Lie algebra over $\mathbb{C}$ is irreducible. Levy [14] extended this result to positive characteristic under mild restrictions on the Lie algebra. On the combinatorics side, Fulman and Guralnick [8] generalized the point-count result of Feit and Fine to commuting varieties of unitary groups and of odd characteristic symplectic groups. We also point out some papers that relate counting problems in Lie algebras to maximal tori of Lie groups; see [9] and [13].

The focus of this paper, the ζ -commuting variety $K_{\zeta,n}$, is another generalization of the commuting variety $K_{1,n}$. When $\zeta = -1$, we get the anti-commuting variety, whose geometry over $\mathbb{C}$ was studied by Chen and Wang [4]. They gave explicit descriptions of the irreducible components of $K_{-1,n}$ and of several variants. The above work was extended to general ζ by Chen and Lu [3]. It is worth noting that $K_{\zeta,n}$ is not irreducible unless $\zeta = 1$. The main contribution of our paper is the point count of $K_{\zeta,n}$.

The point count of $K_{\zeta,n}$ can also be viewed as statistical information on the classification of modules over the quantum plane. More specifically, since an n -dimensional⁽¹⁾ module over the quantum plane $\mathbb{F}_q\{X, Y\}/(XY - \zeta YX)$ can be parametrized by a pair of matrices (A, B) in $K_{\zeta,n}$, the standard orbit-stabilizer argument gives

$$(7) \quad \frac{|K_{\zeta,n}(\mathbb{F}_q)|}{|\mathrm{GL}_n(\mathbb{F}_q)|} = \sum_{\dim M=n} \frac{1}{|\mathrm{Aut} M|},$$

where M ranges over all isomorphism classes of n -dimensional modules over the quantum plane. In other words, the x^n -coefficient of the generating function in (1.1) is the weighted count of isomorphism classes of n -dimensional modules over the quantum plane, with weight inversely proportional to the size of the automorphism group (this weighing is commonly known as the *Cohen–Lenstra measure*, following the important work of Cohen and Lenstra [5] on random abelian groups). While Theorem 1.1 neither requires nor gives a classification of finite-dimensional modules, it does compute their total weight. It is unknown whether Theorem 1.1 can be verified using a classification, via the interpretation (7). For work towards the classification of finite-dimensional modules over the quantum plane, we refer the reader to Bavula [1, §3], where a classification of *simple* modules are given.

For a fixed integer $g \geq 1$, Hausel and Rodriguez-Villegas studied a related counting problem [11, Eq (3.2.3)]

$$(8) \quad N_n(q) := |\{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n(\mathbb{F}_q) : [A_1, B_1] \cdots [A_g, B_g] \zeta_n = 1\}|,$$

where $[A, B] := ABA^{-1}B^{-1}$ and ζ_n is a primitive n -th root of unity of $\mathbb{F}_q$. If $g = 1$, then the defining equation for $N_n(q)$ is $A_1B_1 = \zeta_n B_1A_1$ (replacing ζ_n^{-1} by ζ_n in the process), so we have

$$(9) \quad N_n(q) = |K_{\zeta_n,n}^{\mathrm{GL} \times \mathrm{GL}}(\mathbb{F}_q)|$$

in the notation of Remark 2.2. We emphasize that $N_n(q)$ are the *diagonal* entries of the table $|K_{\zeta_m,n}^{\mathrm{GL} \times \mathrm{GL}}(\mathbb{F}_q)|$ in m, n , which we determine in (46) in terms of a generating function.

Hausel and Rodriguez-Villegas observed a curious functional equation [11, Eq (3.5.12)] about a generating function of $N_n(q)$ that holds for all g , which reads

$$(10) \quad [x^n]E_{\zeta_n}^{\mathrm{GL} \times \mathrm{GL}}(x; q) = -q[x^n]E_{\zeta_n}^{\mathrm{GL} \times \mathrm{GL}}(x; q^{-1})$$

when $g = 1$, where $E_{\zeta_m}^{\mathrm{GL} \times \mathrm{GL}}(x; q)$ is a generating function of $K_{\zeta_m,n}^{\mathrm{GL} \times \mathrm{GL}}$ defined in Remark 2.2, and the operator $[x^n]$ refers to extracting the x^n -coefficient. From our

⁽¹⁾The dimensionality refers to the dimension as an $\mathbb{F}_q$ -vector space.

formula (46) for $E_{\zeta_m}^{\text{GL} \times \text{GL}}(x; q)$, we have

$$(11) \quad [x^n]E_{\zeta_n}^{\text{GL} \times \text{GL}}(x; q) = q - 1,$$

so the $g = 1$ case of the functional equation reads

$$(12) \quad q - 1 = -q(q^{-1} - 1).$$

2. PROOF OF THEOREM 1.2(a)

We recall that Theorem 1.2(a) claims that $|K_{\zeta,n}(\mathbb{F}_q)|$ for all n can be recovered from $|U_{\zeta,n}(\mathbb{F}_q)|$ and $|N_{\zeta,n}(\mathbb{F}_q)|$ for all n . We start by proving a decomposition lemma, following the approach of Feit and Fine [6].

Let V be an n -dimensional vector space over any field, then by Fitting's lemma (see for instance [12, p. 113]), for any linear map $A \in \text{End}(V)$, there is a unique decomposition $V = K_A \oplus I_A$ such that $A(K_A) \subseteq K_A$, $A(I_A) \subseteq I_A$, $A|_{K_A}$ is nilpotent, and $A|_{I_A}$ is nonsingular.

LEMMA 2.1. *Fix a linear map $A \in \text{End}(V)$ and a nonzero scalar ζ . Then a linear map $B \in \text{End}(V)$ satisfies $AB = \zeta BA$ if and only if*

- (a) $B(K_A) \subseteq K_A$, $B(I_A) \subseteq I_A$.
- (b) $A|_{K_A}B|_{K_A} = \zeta B|_{K_A}A|_{K_A}$, $A|_{I_A}B|_{I_A} = \zeta B|_{I_A}A|_{I_A}$.

Proof. Having the decomposition $V = K_A \oplus I_A$, any linear map $X \in \text{End}(V)$ can be written as a matrix

$$(13) \quad X = \begin{bmatrix} X_1 & X_2 \\ X_3 & X_4 \end{bmatrix}, \quad \begin{array}{l} X_1 \in \text{End}(K_A), X_2 \in \text{Hom}(I_A, K_A), \\ X_3 \in \text{Hom}(K_A, I_A), X_4 \in \text{End}(I_A). \end{array}$$

Then we have

$$(14) \quad A = \begin{bmatrix} N & 0 \\ 0 & U \end{bmatrix}$$

where $N \in \text{End}(K_A)$ is nilpotent and $U \in \text{End}(I_A)$ is nonsingular. For an arbitrary

$B = \begin{bmatrix} B_1 & B_2 \\ B_3 & B_4 \end{bmatrix}$, the equation $AB = \zeta BA$ is equivalent to

$$(15) \quad \begin{cases} NB_1 = \zeta B_1N, \\ NB_2 = \zeta B_2U, \\ UB_3 = \zeta B_3N, \\ UB_4 = \zeta B_4U. \end{cases}$$

We note that B_2 must be zero. Suppose not, since N is nilpotent, there exists an integer $r \geq 0$ such that $N^r B_2 \neq 0$ but $N^{r+1} B_2 = 0$. The second equation gives $N^{r+1} B_2 = \zeta N^r B_2 U$. The left-hand side is zero, while the right-hand side is nonzero because ζ is a nonzero scalar and U is nonsingular. This yields a contradiction.

A similar argument shows that $B_3 = 0$, completing the proof of the lemma. $\square$

Let $V = \mathbb{F}_q^n$. To choose $A, B \in \text{End}(V)$ with $AB = \zeta BA$, because of Lemma 2.1, it suffices to choose a decomposition $V = K \oplus I$, and then choose $A_K, B_K \in \text{End}(K)$, $A_I, B_I \in \text{End}(I)$ such that A_K is nilpotent, $A_K B_K = \zeta B_K A_K$, A_I is nonsingular, and $A_I B_I = \zeta B_I A_I$. We have

$$(16) \quad |K_{\zeta,n}(\mathbb{F}_q)| = \sum_{s+t=n} h(s, t) |N_{\zeta,s}(\mathbb{F}_q)| |U_{\zeta,t}(\mathbb{F}_q)|,$$

where $h(s, t)$ is the number of ordered pairs (K, I) of subspaces of V such that $\dim K = s$, $\dim I = t$.

It is noted by Feit and Fine [6, Equation (5)] that

$$(17) \quad h(s, t) = \frac{|\mathrm{GL}_n(\mathbb{F}_q)|}{|\mathrm{GL}_s(\mathbb{F}_q)||\mathrm{GL}_t(\mathbb{F}_q)|}.$$

It follows that

$$(18) \quad \sum_{n=0}^{\infty} \frac{|K_{\zeta,n}(\mathbb{F}_q)|}{|\mathrm{GL}_n(\mathbb{F}_q)|} x^n = \sum_{n=0}^{\infty} \sum_{s+t=n} \frac{|\mathrm{GL}_n(\mathbb{F}_q)|}{|\mathrm{GL}_s(\mathbb{F}_q)||\mathrm{GL}_t(\mathbb{F}_q)|} |N_{\zeta,s}(\mathbb{F}_q)| |U_{\zeta,t}(\mathbb{F}_q)| \frac{1}{|\mathrm{GL}_n(\mathbb{F}_q)|} x^n$$

$$(19) \quad = \sum_{s,t \geq 0} \frac{|N_{\zeta,s}(\mathbb{F}_q)|}{|\mathrm{GL}_s(\mathbb{F}_q)|} \frac{|U_{\zeta,t}(\mathbb{F}_q)|}{|\mathrm{GL}_t(\mathbb{F}_q)|} x^{s+t}$$

$$(20) \quad = \left(\sum_{s=0}^{\infty} \frac{|N_{\zeta,s}(\mathbb{F}_q)|}{|\mathrm{GL}_s(\mathbb{F}_q)|} x^s \right) \left(\sum_{t=0}^{\infty} \frac{|U_{\zeta,t}(\mathbb{F}_q)|}{|\mathrm{GL}_t(\mathbb{F}_q)|} x^t \right),$$

completing the proof of Theorem 1.2(a).

REMARK 2.2. The same argument can prove two other similar factorization identities below, by noting that B is nonsingular (or nilpotent) if and only if both B_K and B_I are nonsingular (or nilpotent). To state the identities, for any combination of symbols F, G taken from $\{\mathrm{Mat}, \mathrm{GL}, \mathrm{Nilp}\}$, we define

$$(21) \quad K_{\zeta,n}^{F \times G} := \{(A, B) \in F_n(\mathbb{F}_q) \times G_n(\mathbb{F}_q) : AB = \zeta BA\},$$

where $\mathrm{Nilp}_n(\mathbb{F}_q)$ denotes the set of n by n nilpotent matrices over $\mathbb{F}_q$. Let

$$(22) \quad E_{\zeta}^{F \times G}(x; q) := \sum_{n=0}^{\infty} \frac{|K_{\zeta,n}^{F \times G}(\mathbb{F}_q)|}{|\mathrm{GL}_n(\mathbb{F}_q)|} x^n.$$

Then

$$(23) \quad E_{\zeta}^{\mathrm{Mat} \times \mathrm{GL}}(x; q) = E_{\zeta}^{\mathrm{GL} \times \mathrm{GL}}(x; q) E_{\zeta}^{\mathrm{Nilp} \times \mathrm{GL}}(x; q);$$

$$(24) \quad E_{\zeta}^{\mathrm{Mat} \times \mathrm{Nilp}}(x; q) = E_{\zeta}^{\mathrm{GL} \times \mathrm{Nilp}}(x; q) E_{\zeta}^{\mathrm{Nilp} \times \mathrm{Nilp}}(x; q).$$

Note that Theorem 1.2(a) can be restated as

$$(25) \quad E_{\zeta}^{\mathrm{Mat} \times \mathrm{Mat}}(x; q) = E_{\zeta}^{\mathrm{Mat} \times \mathrm{GL}}(x; q) E_{\zeta}^{\mathrm{Mat} \times \mathrm{Nilp}}(x; q).$$

3. PROOF OF THEOREM 1.2(b)

Recall that the goal of Theorem 1.2(b) is to determine $|U_{\zeta,n}(\mathbb{F}_q)|$, namely, to enumerate pairs of matrices $(A, B) \in \mathrm{Mat}_n(\mathbb{F}_q) \times \mathrm{Mat}_n(\mathbb{F}_q)$ such that $AB = \zeta BA$ and A is nonsingular. To do so, following the approach of Feit and Fine, let β be a similarity class of $n \times n$ matrices. By a standard orbit-stabilizer argument, for B in β , the number of nonsingular matrices A such that $ABA^{-1} = \zeta B$ is either $|\mathrm{GL}_n(\mathbb{F}_q)|/|\beta|$ or zero. Moreover, this number is not zero if and only if B is similar to ζB . We now give a sufficient and necessary condition for it in terms of β .

We recall that each class β corresponds to a unique rational canonical form. It is characterized by an n -dimensional module M_{β} of the polynomial ring $\mathbb{F}_q[t]$. Such a module can be uniquely expressed in the form of

$$(26) \quad M_{\beta} = \frac{\mathbb{F}_q[t]}{(g_1(t))} \oplus \frac{\mathbb{F}_q[t]}{(g_2(t))} \oplus \cdots \oplus \frac{\mathbb{F}_q[t]}{(g_r(t))}$$

for monic polynomials $g_1, \dots, g_r$ such that g_i divides g_{i+1} for all $1 \leq i \leq r-1$. For a positive integer m , we say a monic polynomial g to be in $\mathcal{P}_m$ if $g(t) = t^b G(t^m)$ for some nonnegative integer b and monic polynomial G . For example, a polynomial is in $\mathcal{P}_2$ if it is either even or odd.

LEMMA 3.1. *Let B be an $n \times n$ matrix over any field, and let ζ be an m -th root of unity. Then B is similar to ζB if and only if the polynomials $g_1, \dots, g_r$ associated to the rational canonical form of B are in $\mathcal{P}_m$.*

Proof. We denote the ground field by $\mathbb{F}$. An endomorphism B over a vector space V determines a module over the polynomial ring $\mathbb{F}[t]$ by letting $t \cdot v = Bv$ for $v \in V$. We denote this $\mathbb{F}[t]$ -module by $(B \curvearrowright V)$. The isomorphism class of this $\mathbb{F}[t]$ -module determines the rational canonical form of B .

Let $g_1, \dots, g_h$ be the polynomials associated to the rational canonical form of B . Then

$$(27) \quad (B \curvearrowright V) \cong \frac{\mathbb{F}[t]}{(g_1(t))} \oplus \frac{\mathbb{F}[t]}{(g_2(t))} \oplus \cdots \oplus \frac{\mathbb{F}[t]}{(g_r(t))}.$$

We now compute $(\zeta B \curvearrowright V)$. We have

$$(28) \quad (\zeta B \curvearrowright V) \cong (\zeta t \curvearrowright M_B)$$

$$(29) \quad \cong \bigoplus_{i=1}^r \left(\zeta t \curvearrowright \frac{\mathbb{F}[t]}{(g_i(t))} \right)$$

$$(30) \quad \cong \bigoplus_{i=1}^r \frac{\mathbb{F}[t]}{(g_i(\zeta^{-1}t))},$$

where the last isomorphism follows from (a): the action of ζt on $\frac{\mathbb{F}[t]}{(g_i(t))}$ is cyclic, and

(b): the polynomial $x \mapsto g_i(\zeta^{-1}x)$ is a minimal polynomial for ζt acting on $\frac{\mathbb{F}[t]}{(g_i(t))}$.

Hence, B is similar to ζB if and only if

$$(31) \quad \bigoplus_{i=1}^r \frac{\mathbb{F}[t]}{(g_i(t))} \cong \bigoplus_{i=1}^r \frac{\mathbb{F}[t]}{(g_i(\zeta^{-1}t))}$$

as $\mathbb{F}[t]$ -modules. Since $g_i(t)$ divides $g_{i+1}(t)$ for all i , we have that $g_i(\zeta^{-1}t)$ divides $g_{i+1}(\zeta^{-1}t)$ as well. By the uniqueness statement about the polynomials associated to the rational canonical form, for each i , the monic polynomials $g_i(t)$ and $\zeta^{\deg g_i} g_i(\zeta^{-1}t)$ must be equal. Write $g_i(t) = t^d + c_1 t^{d-1} + \cdots + c_{d-1} t + c_d$, then $\zeta^d g_i(\zeta^{-1}t) = t^d + \zeta c_1 t^{d-1} + \cdots + \zeta^{d-1} c_{d-1} t + \zeta^d c_d$. Since ζ is an m -th root of unity, we observe that $g_i(t) = \zeta^d g_i(\zeta^{-1}t)$ if and only if $c_j = 0$ for all j not divisible by m . This is equivalent to saying that $g_i(t)$ is in $\mathcal{P}_m$. $\square$

Let $\mathcal{S}_{\zeta,n}(\mathbb{F}_q)$ denote the set of similarity classes β of $n \times n$ matrices over $\mathbb{F}_q$ such that some (equivalently, every) matrix B in β is similar to ζB . We have

$$(32) \quad |U_{\zeta,n}(\mathbb{F}_q)| = \sum_{B \in \text{Mat}_n(\mathbb{F}_q)} |\{A \in \text{GL}_n(\mathbb{F}_q) : ABA^{-1} = \zeta B\}|$$

$$(33) \quad = \sum_{\beta} \sum_{B \in \beta} |\{A \in \text{GL}_n(\mathbb{F}_q) : ABA^{-1} = \zeta B\}|$$

$$(34) \quad = \sum_{\beta \in \mathcal{S}_{\zeta,n}(\mathbb{F}_q)} |\beta| \frac{|\text{GL}_n(\mathbb{F}_q)|}{|\beta|} + \sum_{\beta \notin \mathcal{S}_{\zeta,n}(\mathbb{F}_q)} 0$$

$$(35) \quad = |\text{GL}_n(\mathbb{F}_q)| |\mathcal{S}_{\zeta,n}(\mathbb{F}_q)|.$$

We now count $|\mathcal{S}_{\zeta,n}(\mathbb{F}_q)|$. By Lemma 3.1, a similarity class in $\mathcal{S}_{\zeta,n}(\mathbb{F}_q)$ is characterized by monic polynomials $g_1, g_2, \dots, g_r$ in $\mathcal{P}_m$ such that every polynomial divides the next. Let $h_i = g_{r+1-i}/g_{r-i}$ for $1 \leq i \leq t$, where $g_0 = 1$. It is easily checked from the definition of $\mathcal{P}_m$ that $g_1, \dots, g_r$ are all in $\mathcal{P}_m$ if and only if $h_1, \dots, h_r$ are all in $\mathcal{P}_m$. Let $b_i = \deg h_i$. The only restriction on the monic h_i is that h_i is in $\mathcal{P}_m$ and that

$\sum_{i=1}^r ib_i = n$. We observe the important fact that the number of monic polynomials in $\mathcal{P}_m$ of degree b_i is $q^{\lfloor b_i/m \rfloor}$. Hence to give $g_1, \dots, g_r$, we first choose $(b_i)_{i \geq 1}$ such that $\sum ib_i = n$, and then independently choose h_i in $\mathcal{P}_m$ of degree b_i . It follows that

$$(36) \quad |\mathcal{S}_{\zeta, n}(\mathbb{F}_q)| = \sum_{\substack{b_i \geq 0 \\ \sum ib_i = n}} q^{\lfloor b_i/m \rfloor}.$$

Therefore,

$$(37) \quad \sum_{n=0}^{\infty} \frac{|U_{\zeta, n}(\mathbb{F}_q)|}{|\mathrm{GL}_n(\mathbb{F}_q)|} x^n = \sum_{n=0}^{\infty} |\mathcal{S}_{\zeta, n}(\mathbb{F}_q)| x^n$$

$$(38) \quad = \sum_{n \geq 0} \sum_{\substack{b_i \geq 0 \\ \sum ib_i = n}} q^{\lfloor b_i/m \rfloor} x^n$$

$$(39) \quad = \sum_{b_1, b_2, \dots \geq 0} q^{\lfloor b_i/m \rfloor} x^{\sum ib_i}$$

$$(40) \quad = \prod_{i=1}^{\infty} \sum_{b=0}^{\infty} q^{\lfloor b/m \rfloor} (x^i)^b.$$

By writing $b = km + l$ with $0 \leq l < m$, we get

$$(41) \quad \sum_{b=0}^{\infty} q^{\lfloor b/m \rfloor} x^b = \sum_{l=0}^{m-1} \sum_{k=0}^{\infty} q^k x^{km+l}$$

$$(42) \quad = \sum_{l=0}^{m-1} \frac{x^l}{1 - qx^m}$$

$$(43) \quad = \frac{1 + x + \dots + x^{m-1}}{1 - qx^m}$$

$$(44) \quad = \frac{1 - x^m}{(1 - x)(1 - qx^m)}.$$

Hence, if we define $G_m(x; q) = \frac{1 - x^m}{(1 - x)(1 - qx^m)}$, then we have

$$(45) \quad \sum_{n=0}^{\infty} \frac{|U_{\zeta, n}(\mathbb{F}_q)|}{|\mathrm{GL}_n(\mathbb{F}_q)|} x^n = \prod_{i=1}^{\infty} G_m(x^i; q),$$

finishing the proof of Theorem 1.2(b).

REMARK 3.2. The same argument can compute a similar generating function below, by noting that B is nonsingular if and only if each polynomial $g_i(t)$ that appears in the rational canonical form has a nonzero constant term. In the notation of Remark 2.2, we have

$$(46) \quad E_{\zeta}^{\mathrm{GL} \times \mathrm{GL}}(x; q) = \prod_{i=1}^{\infty} \frac{1 - x^{im}}{1 - x^{im}q},$$

where $m = \mathrm{ord}(\zeta)$.

Similarly, if we instead notice that B is nilpotent if and only if each $g_i(t)$ is a power of t , then we get

$$(47) \quad E_{\zeta}^{\mathrm{GL} \times \mathrm{Nilp}}(x; q) = \prod_{i=1}^{\infty} \frac{1}{1 - x^i}.$$

We notice that the above two formulas, together with Theorem 1.2(b), verify (23) explicitly. We also observe that $E_{\zeta}^{\text{GL} \times \text{GL}}(x; q)$ is a power series in x^m . In particular, this implies that if $AB = \zeta BA$ and A, B are both nonsingular, then the size n of the matrices A, B must be a multiple of the order of ζ .

4. PROOF OF THEOREM 1.2(c)

We follow the idea of Fine and Herstein [7] to determine $|N_{\zeta, n}(\mathbb{F}_q)|$, namely, the number of matrix pairs $(A, B) \in \text{Mat}_n(\mathbb{F}_q) \times \text{Mat}_n(\mathbb{F}_q)$ such that $AB = \zeta BA$ and A is nilpotent. In fact, we will show that the situation is completely the same as the case $\zeta = 1$ studied in [7].

Associate to each similarity class of n by n nilpotent matrices a partition π of n :

$$(48) \quad \pi : n = a_1 \cdot 1 + a_2 \cdot 2 + \dots,$$

so that a representative of the similarity class associated to π is given by

$$(49) \quad A_{\pi} = \begin{bmatrix} 0_{a_1} & & & & & \\ & 0_{a_2} & 1_{a_2} & & & \\ & & 0_{a_2} & & & \\ & & & 0_{a_3} & 1_{a_3} & \\ & & & & 0_{a_3} & 1_{a_3} \\ & & & & & 0_{a_3} \\ & & & & & & \ddots \end{bmatrix},$$

where 0_a and 1_a denote the a by a zero matrix and the a by a identity matrix, respectively.

Let $\alpha(\pi)$ denote the similarity class associated to π . Since the number of matrices B such that $AB = \zeta BA$ only depends on the similarity class of A , we have

$$(50) \quad |N_{\zeta, n}(\mathbb{F}_q)| = \sum_{\pi \vdash n} |\alpha(\pi)| |\{B \in \text{Mat}_n(\mathbb{F}_q) : A_{\pi} B = \zeta B A_{\pi}\}|.$$

For any fixed scalar $\zeta \neq 0$, it is elementary to check that $A_{\pi} B = \zeta B A_{\pi}$ if and only if B is of the following form:

$$(51) \quad \begin{bmatrix} B_{1,1}^1 & B_{1,2}^1 & B_{1,3}^1 & B_{1,4}^1 & \cdots \\ B_{2,1}^1 & B_{2,2}^1 & B_{2,3}^1 & B_{2,4}^1 & \cdots \\ & \zeta B_{2,2}^1 & \zeta B_{2,3}^1 & \zeta B_{2,4}^1 & \cdots \\ B_{3,1}^1 & B_{3,2}^1 & B_{3,3}^1 & B_{3,4}^1 & \cdots \\ & \zeta B_{3,2}^1 & \zeta B_{3,3}^1 & \zeta B_{3,4}^1 & \cdots \\ & & \zeta^2 B_{3,3}^1 & \zeta^2 B_{3,4}^1 & \cdots \\ B_{4,1}^1 & B_{4,2}^1 & B_{4,3}^1 & B_{4,4}^1 & \cdots \\ & \zeta B_{4,2}^1 & \zeta B_{4,3}^1 & \zeta B_{4,4}^1 & \cdots \\ & & \zeta^2 B_{4,3}^1 & \zeta^2 B_{4,4}^1 & \cdots \\ & & & \zeta^3 B_{4,4}^1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

where each $B_{i,j}^k$ is an arbitrary a_i by a_j matrix, chosen independently. We note that the count $|\{B \in \text{Mat}_n(\mathbb{F}_q) : A_{\pi} B = \zeta B A_{\pi}\}|$ does not depend on ζ . Hence,

$$(52) \quad |N_{\zeta, n}(\mathbb{F}_q)| = |N_{1, n}(\mathbb{F}_q)|.$$

It is known [6, Equation (6)] that

$$(53) \quad |N_{1, n}(\mathbb{F}_q)| = |\text{GL}_n(\mathbb{F}_q)| \sum_{\pi \vdash n} \frac{1}{f(a_1) f(a_2) \cdots},$$

where $f(a) := (1 - q^{-1})(1 - q^{-2}) \dots (1 - q^{-a})$.

Hence,

$$(54) \quad \sum_{n=0}^{\infty} \frac{|N_{\zeta,n}(\mathbb{F}_q)|}{|\mathrm{GL}_n(\mathbb{F}_q)|} x^n = \sum_{n=0}^{\infty} \sum_{\pi \vdash n} \frac{1}{f(a_1)f(a_2)\dots} x^n$$

$$(55) \quad = \sum_{a_1, a_2, \dots \geq 0} \frac{1}{f(a_1)f(a_2)\dots} x^{\sum i a_i}$$

$$(56) \quad = \prod_{i=1}^{\infty} \sum_{a=0}^{\infty} \frac{1}{f(a)} (x^i)^a$$

$$(57) \quad = \prod_{i=1}^{\infty} H(x^i; q),$$

where

$$(58) \quad H(x; q) := \sum_{a=0}^{\infty} \frac{1}{f(a)} x^a = \frac{1}{(1-x)(1-xq^{-1})(1-xq^{-2})\dots}$$

by a classical identity due to Euler. This concludes the proof of Theorem 1.2(c), and hence proves Theorem 1.2 and Theorem 1.1.

REMARK 4.1. Combining Theorem 1.2(c), formula (47) and the decomposition formula (24), we get (in the notation of Remark 2.2)

$$(59) \quad E_{\zeta}^{\mathrm{Nilp} \times \mathrm{Nilp}}(x; q) = \prod_{i=1}^{\infty} \frac{1}{(1-x^i q^{-1})(1-x^i q^{-2})\dots}.$$

At this point, we have computed $E_{\zeta}^{\mathrm{F} \times \mathrm{G}}(x; q)$ for all combinations of $\mathrm{F}, \mathrm{G} \in \{\mathrm{Mat}, \mathrm{GL}, \mathrm{Nilp}\}$. We notice that $E_{\zeta}^{\mathrm{F} \times \mathrm{G}}(x; q)$ does not depend on ζ whenever F or G is Nilp . This should not be surprising in light of the argument of Theorem 1.2(c).

5. DISCUSSIONS

We note from the work of Bryan and Morrison [2, §3.1] that $|U_{1,n}(\mathbb{F}_q)|$ and $|N_{1,n}(\mathbb{F}_q)|$ “determine” each other. The key ingredient is that either of the quantities above is the point count of the variety of modules over the “commutative” plane $\mathrm{Spec} \mathbb{F}_q[x, y]$ supported on a certain subset of closed points. A module is determined by its localizations at closed points in its support, so both $|U_{1,n}(\mathbb{F}_q)|$ and $|N_{1,n}(\mathbb{F}_q)|$ are determined by the point count of the variety of modules supported at a point. Since the commutative plane “looks the same everywhere” locally in light of the Cohen structure theorem (the complete localization of $\mathbb{F}_q[x, y]$ at any closed point is isomorphic to $\mathbb{F}[[x, y]]$ for some field extension $\mathbb{F}$ of $\mathbb{F}_q$), we can reverse the process, so that either of $|U_{1,n}(\mathbb{F}_q)|$ and $|N_{1,n}(\mathbb{F}_q)|$ determines the point count of the variety of modules supported at a point, and hence determines each other.

However, for $\zeta \neq 1$, Theorem 1.2 shows that $|N_{\zeta,n}(\mathbb{F}_q)|$ does not depend on ζ while $|U_{\zeta,n}(\mathbb{F}_q)|$ does. Is it still possible to recover $|U_{\zeta,n}(\mathbb{F}_q)|$ from $|N_{\zeta,n}(\mathbb{F}_q)|$ together with the geometry of the quantum plane $xy = \zeta yx$ (which will depend on ζ)?

Acknowledgements. We thank Jason Bell for proposing questions that inspire this work. We thank Jeffrey Lagarias and Weiqiang Wang for fruitful conversations. We thank the referees for helpful comments and informing the author about related work from [11].

REFERENCES

- [1] Vladimir Bavula, *Classification of the simple modules of the quantum Weyl algebra and the quantum plane*, in Quantum groups and quantum spaces (Warsaw, 1995), Banach Center Publ., vol. 40, Polish Acad. Sci. Inst. Math., Warsaw, 1997, pp. 193–201.
- [2] Jim Bryan and Andrew Morrison, *Motivic classes of commuting varieties via power structures*, J. Algebraic Geom. **24** (2015), no. 1, 183–199.
- [3] Xinhong Chen and Meining Lu, *Varieties of modules over the quantum plane*, 2019, <https://arxiv.org/abs/1910.02198>.
- [4] Xinhong Chen and Weiqiang Wang, *Anti-commuting matrices*, Trans. Amer. Math. Soc. **373** (2020), no. 3, 1597–1617.
- [5] H. Cohen and H. W. Lenstra, Jr., *Heuristics on class groups of number fields*, in Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983), Lecture Notes in Math., vol. 1068, Springer, Berlin, 1984, pp. 33–62.
- [6] Walter Feit and N. J. Fine, *Pairs of commuting matrices over a finite field*, Duke Math. J. **27** (1960), 91–94.
- [7] N. J. Fine and I. N. Herstein, *The probability that a matrix be nilpotent*, Illinois J. Math. **2** (1958), 499–504.
- [8] Jason Fulman and Robert Guralnick, *Enumeration of commuting pairs in Lie algebras over finite fields*, Ann. Comb. **22** (2018), no. 2, 295–316.
- [9] Jason Fulman, Rita Jiménez Rolland, and Jennifer C. H. Wilson, *Generating functions and statistics on spaces of maximal tori in classical Lie groups*, New York J. Math. **23** (2017), 165–191.
- [10] Murray Gerstenhaber, *On dominance and varieties of commuting matrices*, Ann. of Math. (2) **73** (1961), 324–348.
- [11] Tamás Hausel and Fernando Rodriguez-Villegas, *Mixed Hodge polynomials of character varieties. With an appendix by Nicholas M. Katz*, Invent. Math. **174** (2008), no. 3, 555–624.
- [12] Nathan Jacobson, *Basic algebra. II*, second ed., W. H. Freeman and Company, New York, 1989.
- [13] G. I. Lehrer, *Rational tori, semisimple orbits and the topology of hyperplane complements*, Comment. Math. Helv. **67** (1992), no. 2, 226–251.
- [14] Paul Levy, *Commuting varieties of Lie algebras over fields of prime characteristic*, J. Algebra **250** (2002), no. 2, 473–484.
- [15] T. S. Motzkin and Olga Taussky, *Pairs of matrices with property L. II*, Trans. Amer. Math. Soc. **80** (1955), 387–401.
- [16] R. W. Richardson, *Commuting varieties of semisimple Lie algebras and algebraic groups*, Compositio Math. **38** (1979), no. 3, 311–327.

YIFENG HUANG, University of Michigan, Dept. of Mathematics, 530 Church St, Ann Arbor, MI 48109 (USA)
 E-mail : huangyf@umich.edu