

ALGEBRAIC COMBINATORICS


Dongkwan Kim & Pavlo Pylyavskyy

Sign insertion and Kazhdan–Lusztig cells of affine symmetric groups

Volume 6, issue 1 (2023), p. 213-241.

<https://doi.org/10.5802/alco.233>


© The author(s), 2023.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Sign insertion and Kazhdan–Lusztig cells of affine symmetric groups

Dongkwan Kim & Pavlo Pylyavskyy

ABSTRACT Combinatorics of Kazhdan–Lusztig cells in affine type A was originally developed by Lusztig, Shi, and Xi. Building on their work, Chmutov, Pylyavskyy, and Yudovina introduced the affine matrix-ball construction (abbreviated AMBC) which gives an analog of Robinson–Schensted correspondence for affine symmetric groups. An alternative approach to Kazhdan–Lusztig theory in affine type A was developed by Blasiak in his work on catabolism. He introduced a sign insertion algorithm and conjectured that if one fixes the two-sided cell, the recording tableau of the sign insertion process determines uniquely and is determined uniquely by the left cell. In this paper we unite these two approaches by proving Blasiak’s conjecture. In the process, we show that certain new operations we introduce called partial rotations connect the elements in the intersection of a left cell and a right cell. Lastly, we investigate the connection between Blasiak’s sign insertion and the standardization map acting on the set of semi-standard Young tableaux defined by Lascoux and Schützenberger.

1. INTRODUCTION

In their groundbreaking paper [8], Kazhdan and Lusztig developed a new approach to the representation theory of Hecke algebras. This gave birth to a whole area called Kazhdan–Lusztig theory. Of particular importance in this theory are the objects called cells. Their definition can be summarized as follows. Each Hecke algebra is associated with a Coxeter group W . Kazhdan and Lusztig define a pre-order \leq_L on elements of W . Some pairs v, w of elements of W satisfy both $v \leq_L w$ and $w \leq_L v$, in which case they are said to be left-equivalent, denoted $v \sim_L w$. Similarly one can define right equivalence \sim_R . The respective equivalence classes are called the left cells and the right cells.

In (finite) type A , i.e. when W is the symmetric group, the Kazhdan–Lusztig cell structure is understood in terms of so-called Robinson–Schensted correspondence. This is a bijective correspondence between elements of the symmetric group and pairs (P, Q) of standard Young tableaux of the same shape. It is well known [1, 2, 7, 8] that two permutations lie in the same left (resp. right) cell if and only if they have the same recording tableau Q (resp. insertion tableau P).

In affine type A , i.e. when W is an affine symmetric group, Chmutov, Pylyavskyy, and Yudovina [6] constructed a bijection $W \rightarrow \Omega$, where Ω is the set of triples (P, Q, ρ) such that P and Q are tabloids of the same shape and ρ is an integer vector satisfying certain inequalities. This bijection is called the affine matrix-ball

Manuscript received 30th October 2020, revised 17th November 2021, accepted 19th February 2022.

KEYWORDS. affine symmetric group, affine matrix-ball construction, Kazhdan–Lusztig cells, Lascoux–Schützenberger standardization, sign insertion.

construction, abbreviated AMBC. They show that, parallel to the finite type, fixing the tabloid Q gives all affine permutations in a left cell, while fixing the tabloid P gives all affine permutations in a right cell. Further properties of AMBC were developed by Chmutov–Lewis–Pylyavskyy [5], Chmutov–Frieden–Kim–Lewis–Yudovina [4], and Kim–Pylyavskyy [9].

In his work on catabolism and Garsia–Procesi modules, Blasiak [3] developed an alternative algorithm to determine the left cell to which an affine permutation belongs. He called his algorithm sign insertion. However, some of the key properties of the algorithm remained open. In particular it was in fact Blasiak’s conjecture that once one fixes a two-sided cell, recording tableaux of sign insertion algorithm are in one to one correspondence with left cells. In this paper we prove this conjecture using AMBC, thus relating the two existing approaches to Kazhdan–Lusztig combinatorics in affine type A .

Structure of Kazhdan–Lusztig cells can be encoded by W -graphs, introduced in the original work of Kazhdan and Lusztig [8]. The edges of the W -graphs fall into two categories: directed and undirected. The undirected edges are the star operations, and are generally easier to understand than the directed edges. In type A they coincide with Knuth moves on permutations, and they are known to connect each Kazhdan–Lusztig cell into a single connected component. In affine type A , however, these star operations are not sufficient to connect the whole cell. The connected components in that case were studied in [5].

The main tool we use in the proof of Blasiak’s conjecture is the new operations called partial rotations and inverse partial rotations. We show that, under certain assumption, they preserve both the left and the right cell of an affine permutation, and thus the only piece of data they change is the weight ρ in terms of AMBC. We also prove that (inverse) partial rotations turn the intersection of a left cell and a right cell into a single connected component. We believe that this makes partial rotations to have independent interest for future study of Kazhdan–Lusztig combinatorics in affine type A .

Lastly, we discuss Blasiak’s sign insertion algorithm with the standardization map of Lascoux and Schützenberger [12, 11]. The latter is a function which maps a semistandard Young tableau to another tableau of the same shape but of different content. This function is known to preserve the cocharge statistic of semistandard Young tableaux. More precisely, the set of semistandard Young tableaux of fixed content is equipped with a poset structure whose grading is given by cocharge statistic. Then the function of Lascoux and Schützenberger yields a graded poset embedding between such posets of different content.

In this paper, we describe how Blasiak’s sign insertion can be understood in terms of the standardization map of Lascoux and Schützenberger. The validity of Blasiak’s conjecture states that with each recording tableau of the sign insertion algorithm one may associate corresponding row-standard Young tableaux which parametrize corresponding left cells. As these cells are in different two-sided cells, these tableaux should have different shapes. Then our theorem states that the images of (the reading words of) such tableaux under RSK correspondence are closely related in terms of the aforementioned standardization map. (See Theorem 7.1 for the exact statement.)

The paper proceeds as follows. In Section 2 we introduce notation we use when working with affine symmetric groups. In Section 3 we remind the reader of the key properties of Kazhdan–Lusztig cells and AMBC. In Section 4 we describe Blasiak’s sign insertion and state our main theorem. In Section 5 we introduce partial rotations, and show how they interact with AMBC and sign insertion. In Section 6 we prove Blasiak’s conjecture. We also prove that partial rotations connect elements in the

intersection of a left cell and a right cell. Finally in Section 7 we show how Blasiak’s sign insertion is related to the standardization map of Lascoux and Schützenberger.

2. NOTATIONS AND DEFINITIONS

For a set A , we write $\#A$ to be its cardinal. For $a, b \in \mathbb{Z}$, we set $[a, b] = \{x \in \mathbb{Z} \mid a \leq x \leq b\}$. A word is a finite sequence of integers, e.g. $w = (w_1, w_2, \dots, w_k)$ for $k \in \mathbb{N} = \{0, 1, 2, \dots\}$ and $w_1, \dots, w_k \in \mathbb{Z}$. For such a word we also write $w = w_1 \cdots w_k$. When $k = 0$, we also write $w = \emptyset$. For two words x and y , we write $x \frown y$ to be their concatenation.

We say that α is a composition of n , denoted $\alpha \models n$, if $\alpha = (a_1, \dots, a_k)$ for some $k, a_1, \dots, a_k \in \mathbb{N} = \{0, 1, 2, \dots\}$ where $\sum_{i=1}^k a_i = n$. Here k is said to be the length of α , also denoted by $l(\alpha)$. Unless otherwise specified, the length of α is the largest integer such that $a_{l(\alpha)} \neq 0$. However, sometimes we allow it to have a zero tail at the end. A composition $\alpha = (a_1, \dots, a_k) \models n$ is called a partition of n , denoted $\alpha \vdash n$, if $a_1 \geq \dots \geq a_k > 0$. For a composition $\alpha = (a_1, \dots, a_{l(\alpha)}) \models n$, we set $\alpha^{\text{rev}} = \{a_{l(\alpha)}, \dots, a_1\}$, which is also a composition of n .

A tabloid is a sequence $T = (T_1, \dots, T_k)$ where each T_i is a sequence of positive integers. We also use its Young diagram to depict T where the i -th row is equal to T_i . We define its shape $\text{sh}(T)$ to be the composition given by the lengths of parts of T . When $\text{sh}(T)$ is a partition we say that T is a tableau. We define the length of the tabloid T to be that of its shape, usually denoted by $l(T)$. For a tabloid T , its reading word, denoted $\text{rw}(T)$, is defined to be the concatenation of its rows from bottom to top. For example, if $T = ((3, 6, 7, 9), (4, 8, 10), (1, 5), (2))$ then $\text{rw}(T) = (2, 1, 5, 4, 8, 10, 3, 6, 7, 9)$. The content of a tabloid T is a composition, say $\alpha = (a_1, a_2, \dots)$, where each a_i is the number of letters in T equal to i .

For a composition $\alpha = (a_1, \dots, a_k) \models n$, we write $\text{RSYT}(\alpha)$ to be the set of row-standard Young tabloid of shape α , i.e. the set of tabloids $T = (T_1, \dots, T_k)$ where each element of $[1, n]$ appears exactly once in T and each T_i is an increasing sequence of length a_i . Also we let $\text{RSYT}(n) = \bigsqcup_{\lambda \vdash n} \text{RSYT}(\lambda)$, the set of row-standard Young tableaux of size n (only of partition shape). For a composition $\alpha = (\alpha_1, \alpha_2, \dots) \models n$ and a partition $\lambda \vdash n$, we denote by $\text{SSYT}(\lambda, \alpha)$ the set of semistandard Young tableaux of shape λ and content α . We also set $\text{SSYT}(n, \alpha) = \bigsqcup_{\lambda \vdash n} \text{SSYT}(\lambda, \alpha)$. When $\alpha = (1^n)$ we write $\text{SYT}(\lambda)$ and $\text{SYT}(n)$ instead of $\text{SSYT}(\lambda, (1^n))$ and $\text{SSYT}(n, (1^n))$, called the set of standard Young tableaux.

For $n \in \mathbb{Z}_{>0}$, we set $\widetilde{\mathcal{S}}_n$ to the extended affine symmetric group defined by

$$\widetilde{\mathcal{S}}_n = \{w: \mathbb{Z} \rightarrow \mathbb{Z} \mid w \text{ is bijective, } w(i+n) = w(i) + n \text{ for } i \in \mathbb{Z}\}.$$

For $w \in \widetilde{\mathcal{S}}_n$, its window notation is defined to be $[w(1), \dots, w(n)]$. We often identify w with its window notation. We define the affine symmetric group $\overline{\mathcal{S}}_n$ to be

$$\overline{\mathcal{S}}_n = \{w \in \widetilde{\mathcal{S}}_n \mid w(1) + \dots + w(n) = n(n+1)/2\},$$

which is a subgroup of $\widetilde{\mathcal{S}}_n$. We set $s_i = [1, \dots, i-1, i+1, i, i+2, \dots, n]$ for $i \in [1, n-1]$ and $s_0 = s_n = [0, 2, \dots, n-1, n+1]$. Then $(\overline{\mathcal{S}}_n, \{s_1, \dots, s_{n-1}, s_n = s_0\})$ is an affine Weyl group of type A . Moreover, if we set $\omega = [2, 3, \dots, n+1] \in \widetilde{\mathcal{S}}_n$ then $\langle \omega \rangle$ is an infinite cyclic group and $\overline{\mathcal{S}}_n = \langle \omega \rangle \times \overline{\mathcal{S}}_n$. Here ω is called the shift element.

For $n \in \mathbb{Z}_{>0}$, we say that w is a (affine) partial permutation if there exists $X \subset [1, n]$ such that w is an injection from $X + n\mathbb{Z}$ to \mathbb{Z} satisfying $w(i+n) = w(i) + n$ for any $i \in X + n\mathbb{Z}$. Note that $X = [1, n]$ if and only if $w \in \widetilde{\mathcal{S}}_n$. We identify w with its window notation $[w(1), \dots, w(n)]$ where we adopt the convention that $w(i) = \emptyset$ whenever $i \notin X$. We often identify a partial permutation $w : X + n\mathbb{Z} \rightarrow \mathbb{Z}$ for $X \subset [1, n]$ with

its graph $\{(x, w(x)) \mid x \in X + n\mathbb{Z}\}$. Thus for example, for two partial permutations $u : X + n\mathbb{Z} \rightarrow \mathbb{Z}$ and $v : X' + n\mathbb{Z} \rightarrow \mathbb{Z}$, we write $u \subset v$ if $X \subset X'$ and $u(x) = v(x)$ for $x \in X$. In addition, we consider the graphs to be placed on the xy -plane such that the x -axis directs to the south and the y -axis directs to the east, i.e. it is obtained from the xy -plane with the conventional direction by rotating 90° clockwise.


For $n \in \mathbb{Z}_{>0}$, let $\mathcal{R} : \mathbb{Z} \rightarrow \mathbb{Z}$ be an involution defined by $\mathcal{R}(x) = n + 1 - x$. This also induces an involution on the set of partial permutations defined by $w \mapsto \mathcal{R} \circ w \circ \mathcal{R}$, which we again denote by \mathcal{R} . Pictorially, it corresponds to the rotation of the diagram of w by 180 degrees which preserves the square $[1, n] \times [1, n]$.

For a sequence $A = (A_1, \dots, A_r)$ and a set $X = \{x_1, \dots, x_s\} \subset [1, r]$ so that $x_1 < x_2 < \dots < x_s$, we set $A_X = (A_{x_1}, \dots, A_{x_s})$. If $X = [1, s]$ (resp. $X = [s, r]$), we also write $A_{\leq s}$ (resp. $A_{\geq s}$) instead of A_X . We set $A|_X = (\emptyset, A_{x_1}, \emptyset, \dots, \emptyset, A_{x_s}, \emptyset)$ (i.e replacing A_i with \emptyset if $i \notin X$). When $w \in \widetilde{\mathcal{S}}_n$ and $X \subset [1, n]$ we set $w|_X = (w(1), \dots, w(n))|_X$, i.e. we identify w with its window notation and discard $w(i)$ for $i \notin X$. Note that this becomes a window notation of some partial permutation; we abuse notation and also denote it by $w|_X$.

3. CELLS, STAR OPERATIONS, AND AMBC

Kazhdan and Lusztig [8] introduced (two-sided, left, and right) *cells* for each Coxeter group and defined (right and left) star operations. (The notion of cells is extended to extended affine Weyl groups by Lusztig [13].) Here we focus on the case of (extended) affine symmetric groups in terms of the affine matrix-ball construction.

For a row-standard Young tableau T and $i \in [1, l(T) - 1]$ suppose that $T_i = (a_1, \dots, a_s)$ and $T_{i+1} = (b_1, \dots, b_t)$. Then we define $\text{lch}_i(T)$, called the local charge in row i of T , to be the smallest $d \in \mathbb{N}$ such that $a_{l-d} < b_l$ for $l \in [d+1, t]$.⁽¹⁾ Pictorially, this corresponds to how much one needs to shift T_i to the right so that (T_i, T_{i+1}) becomes standard. For example, if $T_i = (3, 5, 7, 8)$ and $T_{i+1} = (1, 2, 4, 6)$ then we have $\text{lch}_i(T) = 2$ as depicted below.



For $P, Q \in \text{RSYT}(\lambda)$ where $\lambda = (\lambda_1, \dots, \lambda_l)$ is a partition, we define $\#_{P,Q} = (s_1, \dots, s_l) \in \mathbb{Z}^l$, called the symmetrized offset constant of (P, Q) , to be as follows.

$$s_i = \begin{cases} 0 & \text{if } i = 1 \text{ or } \lambda_{i-1} > \lambda_i, \\ s_{i-1} + \text{lch}_{i-1}(P) - \text{lch}_{i-1}(Q) & \text{otherwise.} \end{cases}$$

In other words, we have $s_i - s_{i-1} = \text{lch}_{i-1}(P) - \text{lch}_{i-1}(Q)$ whenever $\lambda_{i-1} = \lambda_i$. We say that $\# = (\rho_1, \dots, \rho_l)$ is *dominant* with respect to (P, Q) if $\rho_{i-1} - s_{i-1} \leq \rho_i - s_i$ whenever $\lambda_{i-1} = \lambda_i$.

The affine matrix-ball construction (abbreviated AMBC), defined in [6], is a systematic method to understand the combinatorial properties of $\widetilde{\mathcal{S}}_n$ and yields two functions

$$\begin{aligned} \Phi : \widetilde{\mathcal{S}}_n &\rightarrow \bigsqcup_{\lambda \vdash n} \text{RSYT}(\lambda) \times \text{RSYT}(\lambda) \times \mathbb{Z}^{l(\lambda)}, \\ \Psi : \bigsqcup_{\lambda \vdash n} \text{RSYT}(\lambda) \times \text{RSYT}(\lambda) \times \mathbb{Z}^{l(\lambda)} &\rightarrow \widetilde{\mathcal{S}}_n. \end{aligned}$$

⁽¹⁾One can easily show that this definition coincides with [5, Definition 5.3].

Here, we have $\text{im } \Phi = \{(P, Q, \overset{\#}{\rho}) \mid \overset{\#}{\rho} \text{ is dominant with respect to } (P, Q)\}$ and $\text{im } \Psi = \widetilde{\mathcal{S}}_n$. Moreover, $\Psi|_{\text{im } \Phi}$ and Φ are inverse to each other. (Their constructions are explained in Section 5.1 in more detail.)

The two-sided cells of $\widetilde{\mathcal{S}}_n$ are parametrized by partitions of n and we denote such cells by \underline{c}_λ for $\lambda \vdash n$. The left/right cells contained in \underline{c}_λ are parametrized by row-standard Young tableaux of shape λ , and we write Γ_T (resp. $(\Gamma_T)^{-1}$) for $T \in \text{RSYT}(\lambda)$ for such a left (resp. right) cell. Here, we have $w \in \Gamma_T$ if and only if $\Phi(w) = (P, T, \overset{\#}{\rho})$ for some P and $\overset{\#}{\rho}$, and similarly $w \in (\Gamma_T)^{-1}$ if and only if $\Phi(w) = (T, Q, \overset{\#}{\rho})$ for some Q and $\overset{\#}{\rho}$. Also we have $\underline{c}_\lambda = \bigsqcup_{T \in \text{RSYT}(\lambda)} \Gamma_T = \bigsqcup_{T \in \text{RSYT}(\lambda)} (\Gamma_T)^{-1}$.

For $w \in \widetilde{\mathcal{S}}_n$ and $i \in [1, n]$ such that either $w(i) < w(i+2) < w(i+1)$, $w(i+1) < w(i+2) < w(i)$, $w(i) < w(i-1) < w(i+1)$, or $w(i+1) < w(i-1) < w(i)$, we define the right star operation $w \mapsto w^*$ for $* \sim i$ to be the one obtained from w by swapping $w(i+kn)$ and $w(i+1+kn)$ for each $k \in \mathbb{Z}$. Similarly, we define the left star operation $w \mapsto {}^*w$ for $* \sim i$ to be the composition $w \mapsto w^{-1} \mapsto (w^{-1})^* \mapsto ((w^{-1})^*)^{-1}$ when the corresponding right star operation $w^{-1} \mapsto (w^{-1})^*$ for $* \sim i$ is well-defined. Then [9, Proposition 5.10] states that when $\Phi(w) = (P, Q, \overset{\#}{\rho}) = (P, Q, \overset{\#}{\mathbb{S}}_{P,Q} + \overset{\#}{\bullet}\rho)$ we have

$$\begin{aligned} \Phi({}^*w) &= \begin{cases} (P^*, Q, \overset{\#}{\mathbb{S}}_{P^*,Q} + \overset{\#}{\bullet}\rho) & \text{if } * \not\sim n, \\ (P^*, Q, \overset{\#}{\mathbb{S}}_{P^*,Q} + \overset{\#}{\bullet}\rho - \overset{\#}{\delta}(P, 1) + \overset{\#}{\delta}(P, n)) & \text{if } * \sim n. \end{cases} \\ \Phi(w^*) &= \begin{cases} (P, Q^*, \overset{\#}{\mathbb{S}}_{P,Q^*} + \overset{\#}{\bullet}\rho) & \text{if } * \not\sim n, \\ (P, Q^*, \overset{\#}{\mathbb{S}}_{P,Q^*} + \overset{\#}{\bullet}\rho + \overset{\#}{\delta}(Q, 1) - \overset{\#}{\delta}(Q, n)) & \text{if } * \sim n, \end{cases} \end{aligned}$$

where P^* and Q^* are Knuth moves for $* \sim i$ defined in [9, Definition 5.7]. Also, for $T \in \text{RSYT}(\lambda)$, $a \in [1, l(T)]$, and $s \in T_a$, we define $\overset{\#}{\delta}(T, s) = (\delta_1, \dots, \delta_{l(T)})$ to be

$$\delta_i = \begin{cases} 1 & \text{if } \lambda_{a-1} > \lambda_a = \lambda_i \text{ (here we set } \lambda_0 = \infty), \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

(In particular, $\delta_i = \delta_j$ whenever $\lambda_i = \lambda_j$.)

Multiplication by $\omega = [2, 3, \dots, n+1] \in \widetilde{\mathcal{S}}_n$ preserves each two-sided cell but permutes left/right cells. More precisely, when $\Phi(w) = (P, Q, \overset{\#}{\rho}) = (P, Q, \overset{\#}{\mathbb{S}}_{P,Q} + \overset{\#}{\bullet}\rho)$ we have (see [9, Proposition 5.5])

$$\begin{aligned} \Phi(\omega \cdot w) &= (\omega(P), Q, \overset{\#}{\mathbb{S}}_{\omega(P),Q} + \overset{\#}{\bullet}\rho + \overset{\#}{\delta}(P, n)), \text{ and} \\ \Phi(w \cdot \omega^{-1}) &= (P, \omega(Q), \overset{\#}{\mathbb{S}}_{P,\omega(Q)} + \overset{\#}{\bullet}\rho - \overset{\#}{\delta}(Q, n)). \end{aligned}$$

Here, $\omega(T)$ is a tableau obtained from T by replacing each entry i with $i+1$ for $i \neq n$ and n with 1 (and reordering entries in each row if necessary so that the result becomes row-standard).

EXAMPLE 3.1. Suppose that $w = [1, 6, 8, 14, 17, 5, 0, 19, 3, 22] \in \widetilde{\mathcal{S}}_{10}$. Then we have

$$\begin{aligned} {}^*w &= [1, 6, 8, 14, 17, 5, 0, 19, 2, 23] \text{ for } * \sim 2, \\ w^* &= [12, 6, 8, 14, 17, 5, 0, 19, 3, 11] \text{ for } * \sim 10, \\ \omega w &= [2, 7, 9, 15, 18, 6, 1, 20, 4, 23], \\ w\omega^{-1} &= [12, 1, 6, 8, 14, 17, 5, 0, 19, 3]. \end{aligned}$$

In each case we have (all the symmetrized offset constants are zero as the parts of $(4, 3, 2, 1)$ are pairwise different)

$$\begin{aligned} \Phi(w) &= (((3, 5, 8, 10), (1, 4, 9), (2, 6), (7)), ((4, 5, 8, 10), (1, 2, 3), (6, 9), (7)), (4, 0, 0, 0)), \\ \Phi(*w) &= (((2, 5, 8, 10), (1, 4, 9), (3, 6), (7)), ((4, 5, 8, 10), (1, 2, 3), (6, 9), (7)), (4, 0, 0, 0)), \\ \Phi(w^*) &= (((3, 5, 8, 10), (1, 4, 9), (2, 6), (7)), ((1, 4, 5, 8), (2, 3, 10), (6, 9), (7)), (3, 1, 0, 0)), \\ \Phi(\omega w) &= (((1, 4, 6, 9), (2, 5, 10), (3, 7), (8)), ((4, 5, 8, 10), (1, 2, 3), (6, 9), (7)), (5, 0, 0, 0)), \\ \Phi(w\omega^{-1}) &= (((3, 5, 8, 10), (1, 4, 9), (2, 6), (7)), ((1, 5, 6, 9), (2, 3, 4), (7, 10), (8)), (3, 0, 0, 0)), \end{aligned}$$

which are as expected from the observation above.

4. SIGN INSERTION AND THE MAIN THEOREM

Here we recall the sign insertion algorithm defined in [3] and state the first main result of this paper (Theorem 4.5). Suppose that we have words \mathfrak{P}, Ω, w such that $w \neq \emptyset$ and $\mathfrak{P} \frown w$ is a window notation of some element in $\widetilde{\mathcal{S}}_n$. (Recall that $x \frown y$ is the concatenation of x and y .) Also we assume that \mathfrak{P} and Ω are both increasing and of the same length. Let us write $w = (a) \frown w''$ where $a \in \mathbb{Z}$ is the first entry of w and w'' is the remaining part. For $i \in \mathbb{Z}_{>0}$ bigger than any entry in Ω , we define $F(i - 1, \mathfrak{P}, \Omega, w) = (i, \mathfrak{P}', \Omega', w')$ as follows.

- If a is bigger than any entry of \mathfrak{P} , then we set $\mathfrak{P}' = \mathfrak{P} \frown (a)$, $\Omega' = \Omega \frown (i)$, and $w' = w''$.
- Otherwise, let b be the smallest element in \mathfrak{P} bigger than a . Then we set \mathfrak{P}' to be \mathfrak{P} after replacing b with a . Also we set $\Omega' = \Omega$ and $w' = w'' \frown (n + b)$.

For $w \in \widetilde{\mathcal{S}}_n$, we start with the quadruple $(0, \emptyset, \emptyset, w)$ and apply F recursively until the last component becomes the empty word, say $(i, \mathfrak{P}, \Omega, \emptyset)$. We write $\text{sgn}_{\mathfrak{P}}(w) = \mathfrak{P}$ and $\text{sgn}_{\Omega}(w) = \Omega$. The process $w \mapsto (\text{sgn}_{\mathfrak{P}}(w), \text{sgn}_{\Omega}(w))$ is called sign insertion.

EXAMPLE 4.1. Let $w = [17, 13, 4, 20, 9, 24] \in \widetilde{\mathcal{S}}_6$. Each step of the sign insertion applied to w is described as follows.

i	\mathfrak{P}	Ω	w
0	\emptyset	\emptyset	17, 13, 4, 20, 9, 24
1	17	1	13, 4, 20, 9, 24
2	13	1	4, 20, 9, 24, 23
3	4	1	20, 9, 24, 23, 19
4	4, 20	1, 4	9, 24, 23, 19
5	4, 9	1, 4	24, 23, 19, 26
6	4, 9, 24	1, 4, 6	23, 19, 26
7	4, 9, 23	1, 4, 6	19, 26, 30
8	4, 9, 19	1, 4, 6	26, 30, 29
9	4, 9, 19, 26	1, 4, 6, 9	29
10	4, 9, 19, 26, 30	1, 4, 6, 9, 10	29
11	4, 9, 19, 26, 29	1, 4, 6, 9, 10	36
12	4, 9, 19, 26, 29, 36	1, 4, 6, 9, 10, 12	\emptyset

Thus we have $\text{sgn}_{\mathfrak{P}}(w) = (4, 9, 19, 26, 29, 36)$ and $\text{sgn}_{\Omega}(w) = (1, 4, 6, 9, 10, 12)$.

REMARK 4.2. Note that it is not exactly the same as, but equivalent to, the original definition of Blasiak [3, Algorithm 9.7]. Indeed, in our convention the window notation of $w \in \widetilde{\mathcal{S}}_n$ is $[w(1), w(2), \dots, w(n)]$, whereas in [3] it is defined to be $[n + 1 - w^{-1}(1), n + 1 - w^{-1}(2), \dots, n + 1 - w^{-1}(n)]$. Furthermore, our sign insertion algorithm uses the first element of w in each step, whereas in [3] the last element is used. Therefore,

Blasiak’s insertion algorithm applied to w is equivalent to our algorithm applied to $\mathcal{R}(w^{-1})$.

Let us discuss some basic properties of sign insertion.

LEMMA 4.3. *Suppose that $F(i - 1, \mathfrak{P}, \Omega, w) = (i, \mathfrak{P}', \Omega', w')$. Then either $\mathfrak{P} \frown w = \mathfrak{P}' \frown w'$ or $\mathfrak{P}' \frown w'$ is obtained from $\mathfrak{P} \frown w$ by applying a series of right star operations followed by multiplication by ω on the right. As a result, w and $\text{sgn}_{\mathfrak{P}}(w)$ are in the same left cell for $w \in \widetilde{\mathcal{S}}_n$.*

Proof. Suppose that $\mathfrak{P} \frown w \neq \mathfrak{P}' \frown w'$. If we let $w = (a) \frown w''$, then \mathfrak{P}' is obtained from \mathfrak{P} by the “bumping process”, i.e. regarding \mathfrak{P} as an one-row tableau and inserting a to it using the usual Robinson–Schensted insertion algorithm. Thus if we let b be the element bumped out from \mathfrak{P} , then it is clear that $(b) \frown \mathfrak{P}'$ is obtained from $\mathfrak{P} \frown (a)$ by a series of right star operations. Now the result follows as $((b) \frown \mathfrak{P}' \frown w'') \cdot \omega = \mathfrak{P}' \frown w'' \frown (b + n) = \mathfrak{P}' \frown w'$. \square

LEMMA 4.4. *Suppose that $w \in \widetilde{\mathcal{S}}_n$.*

- (1) $\text{sgn}_{\Omega}(w) = \text{sgn}_{\Omega}(\omega w)$.
- (2) *If $*w$ is well-defined for some $* \sim i$, then $\text{sgn}_{\Omega}(*w) = \text{sgn}_{\Omega}(w)$.*

Proof. This is a reformulation of [3, Proposition 9.9 (h), (i)] with respect to our convention. In [3] he only considers when $* \not\sim n$ (see the comment right above [3, Example 9.3]), however the argument still applies when $* \sim n$. \square

Now we state the first main result of this paper, originally conjectured by J. Blasiak.

THEOREM 4.5. [3, Conjecture 9.17(b)] *For $\lambda \vdash n$ and Ω , the set $\{w \in \underline{c}_{\lambda} \mid \text{sgn}_{\Omega}(w) = \Omega\}$ is a single left cell (if nonempty).*

REMARK 4.6. Since his $\text{sgn}_{\Omega}(w)$ is equal to our $\text{sgn}_{\Omega}(\mathcal{R}(w^{-1}))$, his conjecture is translated to that $\{w \in \underline{c}_{\lambda} \mid \text{sgn}_{\Omega}(\mathcal{R}((w^{-1})^{-1})) = \text{sgn}_{\Omega}(\mathcal{R}(w)) = \Omega\}$ is a single left cell. However, since $\mathcal{R} : \widetilde{\mathcal{S}}_n \rightarrow \mathcal{S}_n$ sends a left cell to a left cell (see [4, Theorem 3.1] for more detail), these two statements are equivalent.

5. AMBC, PROPER PARTIAL ROTATIONS, AND SIGN INSERTION

Here we first review the construction of AMBC in detail. Then we introduce *proper partial rotations* which play a central role in our paper and discuss how they are related to both AMBC and Blasiak’s sign insertion. Eventually we verify the Diamond Lemma (Lemma 5.15) which is a key step for the proof of Blasiak’s conjecture.

5.1. NOTIONS FOR AMBC. Here we recap some definitions from [6] that will be used in this paper. We fix a positive integer n .

For a partial permutation w , we define its Shi poset to be the poset $P = P_w$ on $\{x \in [1, n] \mid w(x) \neq \emptyset\}$ such that $i <_P j$ if either $[i > j \text{ and } w(i) < w(j)]$ or $[w(j) > w(i) + n]$. For such a poset, we define its Greene–Kleitman partition to be $\lambda = (\lambda_1, \lambda_2, \dots)$ where $\sum_{i=1}^k \lambda_i$ is the maximum of the number of elements in the union of k -antichains in P_w . Then by [14], if $w \in \widetilde{\mathcal{S}}_n$ is an actual permutation then the Greene–Kleitman partition λ of P_w parametrizes the two-sided cell that contains w , i.e. we have $w \in \underline{c}_{\lambda}$. In such a case we define the width of P_w to be the first row of λ , which is also equal to the maximum length of an antichain in P_w .

EXAMPLE 5.1. Let $w = [8, 1, 19, 14, 16, 2, 25, 13, 10, 27] \in \widetilde{\mathcal{S}}_{10}$. Its Shi poset P_w is described in Figure 1. Direct calculation shows that its Greene–Kleitman partition is $(3, 3, 3, 1)$, which means that $w \in \underline{c}_{(3,3,3,1)}$.




FIGURE 1. Hasse diagram of $P_{[8,1,19,14,16,2,25,13,10,27]}$

We say that a partial permutation $S : X + n\mathbb{Z} \rightarrow \mathbb{Z}$ for some $X \subset [1, n]$ is a stream if S is an increasing function. In such a case its density is defined to be $\#X$. When there exists another partial permutation w such that $S \subset w$, we say that S is a channel of w if the density of S is equal to the width of P_w . Indeed, a stream $S \subset w$ is a channel if and only if its density is maximum among the streams contained in w . For a stream $S : X + n\mathbb{Z} \rightarrow \mathbb{Z}$ its altitude $a(S)$ is defined to be $\sum_{x \in X} (\lceil w(x)/n \rceil - 1)$ where $\lceil t \rceil$ is the smallest integer not smaller than t .

For $(x, y), (x', y') \in \mathbb{Z}^2$ we write $(x, y) \leq_{SW} (x', y')$ if $x \geq x'$ and $y \leq y'$ and say that (x, y) is southwest of (x', y') or equivalently (x', y') is northeast of (x, y) . Similarly, we write $(x, y) \leq_{NW} (x', y')$ if $x \leq x'$ and $y \leq y'$ and say that (x, y) is northwest of (x', y') or equivalently (x', y') is southeast of (x, y) . (Note that this is compatible with our convention of the xy -plane which is the clockwise rotation of the usual convention by 90 degrees.) For two channels $C, C' \subset w$ we say that C is southwest of C' , denoted $C \leq_{SW} C'$, if for any $b \in C$ there exists $b' \in C'$ such that $b \leq_{SW} b'$. If neither $C \leq_{SW} C'$ nor $C \geq_{SW} C'$ then we write $C \not\leq_{SW} C'$. By [6, Proposition 3.13], for any partial permutation w there always exist channels $C, C' \subset w$ such that $C \leq_{SW} C'' \leq_{SW} C'$ for any channel $C'' \subset w$. We call C (resp. C') the southwest (resp. northeast) channel of w .

For a partial permutation w , a sequence $((x_1, w(x_1)), \dots, (x_k, w(x_k)))$ is called a path in w if $x_1 < \dots < x_k$ and $w(x_1) < \dots < w(x_k)$, i.e. it is a chain with respect to the southeast direction. Or equivalently, if (b_1, \dots, b_k) is a path in w then we have $b_1 \leq_{NW} \dots \leq_{NW} b_k$.⁽²⁾ A numbering $d : w \rightarrow \mathbb{Z}$ is called monotone if for any path (b_1, b_2, \dots, b_k) we have $d(b_1) < d(b_2) < \dots < d(b_k)$.

For a stream S we say that $\tilde{d} : S \rightarrow \mathbb{Z}$ is a proper numbering if $\tilde{d}(x, S(x)) < \tilde{d}(y, S(y))$ whenever $x < y$ and $\tilde{d}(x + n, S(x + n)) - \tilde{d}(x, S(x))$ equals the density of S for any x . Note that a proper numbering is unique up to shift. When C is a channel of w and $\tilde{d} : C \rightarrow \mathbb{Z}$ is a proper numbering, we define the channel numbering $d_w^C = d^C : w \rightarrow \mathbb{Z}$ to be

$$d^C(b) = \max\{\tilde{d}(b') + k \mid \text{there exists a path } (b' = b_0, b_1, \dots, b_k = b) \text{ in } w \text{ where } b' \in C\}.$$

Again d^C is uniquely determined up to shift. When C is the southwest channel of w we also write $d = d^C$ (or $d_w = d_w^C$), called the southwest channel numbering of w .

For two channels $C, C' \subset w$, let us fix shifts of d^C and $d^{C'}$ so that they coincide on some/any element in C . (These two conditions are equivalent since the restriction of a proper numbering to a channel is also proper, see [6, Lemma 3.15].) Then the distance between C and C' is defined to be $h(C, C') = |d^{C'}(b) - d^C(b)|$ for some/any $b \in C'$. This is indeed well-defined and satisfies that $h(C, C') + h(C', C'') \geq h(C, C'')$

⁽²⁾This is called a reverse path in [6] and a path is defined to be of the opposite direction. Here we stick to our definition in order to avoid confusion to which way a path is directed.

for any channels $C, C', C'' \subset w$. Furthermore, if $C \not\sim_{SW} C'$ then $h(C, C') = 0$ by [6, Lemma 11.12]. (The converse is not true in general.) We say that R is a river of w if it is of the form $R = \{C' \subset w \mid C' \text{ is a channel of } w, h(C, C') = 0\}$ for some channel $C \subset w$. (Later by Lemma 5.4, such C can be chosen to be any channel in R .)

EXAMPLE 5.2. Let $w = [8, 1, 19, 14, 16, 2, 25, 13, 10, 27] \in \widetilde{\mathcal{S}}_{10}$ as in Example 5.1. There are three channels of w as follows:

$$\begin{aligned} C_1 &= [\emptyset, 1, \emptyset, \emptyset, \emptyset, 2, \emptyset, \emptyset, 10, \emptyset], \\ C_2 &= [8, \emptyset, \emptyset, 14, 16, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset], \\ C_3 &= [\emptyset, \emptyset, 19, \emptyset, \emptyset, \emptyset, 25, \emptyset, \emptyset, 27]. \end{aligned}$$

Here C_1 is the southwest channel of w and C_3 is the northeast channel of w . Let us fix the proper numbering \tilde{d} of C_1 such that $\tilde{d}(2, 1) = 0$. Then the southwest channel numbering $d = d^{C_1} : w \rightarrow \mathbb{Z}$ is defined to be

$$\begin{aligned} (1, 8), (2, 1) &\mapsto 0, & (4, 14), (6, 2) &\mapsto 1, & (3, 19), (5, 16), (8, 13), (9, 10) &\mapsto 2, \\ (7, 25) &\mapsto 3, & (10, 27) &\mapsto 4. \end{aligned}$$

Similarly, $d^{C_2} : w \rightarrow \mathbb{Z}$ is defined to be

$$\begin{aligned} (2, 1) &\mapsto 0, & (1, 8), (6, 2) &\mapsto 1, & (4, 14), (8, 13), (9, 10) &\mapsto 2, \\ (5, 16), (3, 19) &\mapsto 3, & (7, 25) &\mapsto 4, & (10, 27) &\mapsto 5, \end{aligned}$$

and $d^{C_3} : w \rightarrow \mathbb{Z}$ is defined to be

$$\begin{aligned} (2, 1) &\mapsto 0, & (1, 8), (6, 2) &\mapsto 1, & (4, 14), (8, 13), (9, 10) &\mapsto 2, & (5, 16) &\mapsto 3, \\ (3, 19) &\mapsto 4, & (7, 25) &\mapsto 5, & (10, 27) &\mapsto 6. \end{aligned}$$

Therefore we see that $h(C_1, C_2) = 1$, $h(C_2, C_3) = 1$, and $h(C_1, C_3) = 2$.

For a partial permutation w and its channel $C \subset w$, we define $\text{fw}_C(w)$ and $\mathfrak{st}_C(w)$ as follows. Let $d = d_w^C : w \rightarrow \mathbb{Z}$ be the channel numbering defined above. For each $m \in \mathbb{Z}$, we take $x_1, \dots, x_k \in \mathbb{Z}$ such that $x_1 > \dots > x_k$ and $d(x_i, w(x_i)) = m$ for $i \in [1, k]$. Then we have $w(x_1) < \dots < w(x_k)$. (It follows from the monotone property of d .) Set $Z_m = Z'_m \sqcup Z''_m \sqcup Z'''_m$ where

$$\begin{aligned} Z'_m &= \{(x_i, w(x_i)) \mid i \in [1, k]\}, \\ Z''_m &= \{(x_i, w(x_{i+1})) \mid i \in [1, k-1]\}, \\ Z'''_m &= \{(x_k, w(x_1))\}. \end{aligned}$$

The set Z_m is called the zigzag labeled m . Then $w \cap Z_m = Z'_m$. We set $\text{fw}_C(w) = \bigsqcup_{m \in \mathbb{Z}} Z''_m$ and $\mathfrak{st}(w) = \bigsqcup_{m \in \mathbb{Z}} Z'''_m$. Then $\text{fw}_C(w)$ is a partial permutation and $\mathfrak{st}_C(w)$ is a stream whose density is equal to the width of P_w . When C is the southwest channel of w we simply write $\text{fw}(w)$ and $\mathfrak{st}(w)$ instead of $\text{fw}_C(w)$ and $\mathfrak{st}_C(w)$.

EXAMPLE 5.3. Let $w = [8, 1, 19, 14, 16, 2, 25, 13, 10, 27] \in \widetilde{\mathcal{S}}_{10}$ as in Example 5.1 and 5.2. Then with respect to $d = d^{C_1}$ we have

$$\begin{aligned} Z'_0 &= \{(2, 1), (1, 8), (-3, 15)\}, & Z''_0 &= \{(2, 8), (1, 15)\}, & Z'''_0 &= \{(-3, 1)\}, \\ Z'_1 &= \{(6, 2), (4, 14), (0, 17)\}, & Z''_1 &= \{(6, 14), (4, 17)\}, & Z'''_1 &= \{(0, 2)\}, \\ Z'_2 &= \{(9, 10), (8, 13), (5, 16), (3, 19)\}, & Z''_2 &= \{(9, 13), (8, 16), (5, 19)\}, & Z'''_2 &= \{(3, 10)\}. \end{aligned}$$

Also we have $\text{fw}(w) = (Z''_0 \sqcup Z''_1 \sqcup Z''_2) + (n, n)\mathbb{Z}$ and $\mathfrak{st}(w) = (Z'''_0 \sqcup Z'''_1 \sqcup Z'''_2) + (n, n)\mathbb{Z}$.

The map $\Phi : \widetilde{\mathcal{S}}_n \rightarrow \bigsqcup_{\lambda \vdash n} \text{RSYT}(\lambda) \times \text{RSYT}(\lambda) \times \mathbb{Z}^{l(\lambda)}$ is defined as follows. Starting with $w_0 = w \in \widetilde{\mathcal{S}}_n$, we successively calculate $w_{i+1} = \text{fw}(w_i)$ and $S_{i+1} = \mathfrak{st}(w_i)$ until we have an empty partial permutation. For each S_i we set $P_i, Q_i \subset [1, n]$ such that $S_i : Q_i + n\mathbb{Z} \rightarrow P_i + n\mathbb{Z}$ is a bijection. Then we have $\Phi(w) = (P, Q, \frac{\#}{\#})$ where

$P = (P_1, P_2, \dots)$, $Q = (Q_1, Q_2, \dots)$, and $\#p = (a(S_1), a(S_2), \dots)$. Moreover, $\#p$ is always dominant with respect to (P, Q) .

A stream S is said to be compatible with a partial permutation w if $S \cap w = \emptyset$, $S \cup w$ is still a partial permutation, and the density of S is not (strictly) smaller than the width of P_w . In such a case we define the backward numbering $d_w^{\text{bk}, S} = d^{\text{bk}, S} : w \rightarrow \mathbb{Z}$ as follows. First we fix a proper numbering $\tilde{d} : S \rightarrow \mathbb{Z}$, and for $(x, w(x)) \in w$ we let $d(x, w(x)) = \max\{\tilde{d}(y, S(y)) \in S \mid y < x, S(y) < w(x)\}$. Now we repeat the following process:

- If $d(x, w(x)) < d(y, w(y))$ for any x, y such that $x < y$ and $w(x) < w(y)$, then we terminate the process.
- Otherwise, choose $(x, w(x))$ such that:
 - there exists y such that $d(x, w(x)) \geq d(y, w(y))$, $x < y$, and $w(x) < w(y)$, and
 - for any z such that $z < x$ and $w(z) < w(x)$ we have $d(z, w(z)) < d(x, w(x))$.
- For each $i \in \mathbb{Z}$ we lower the value of $d(x + in, w(x) + in)$ by 1.

After this process is finished we set $d_w^{\text{bk}, S} = d$. This numbering is always well-defined.

For a partial permutation w and a compatible stream S , we define $\text{bk}_S(w)$ as follows. Let \tilde{d} be the proper numbering on S and $d = d_w^{\text{bk}, S}$ be the induced backward numbering on w . Then for each $m \in \mathbb{Z}$, we take $y, x_1, \dots, x_k \in \mathbb{Z}$ such that $x_1 > \dots > x_k$ and $\tilde{d}(y, S(y)) = d(x_i, w(x_i)) = m$ for $i \in [1, k]$. Then we have $w(x_1) < \dots < w(x_k)$. (It follows from the monotone property of d .) Set $Z_m = Z'_m \sqcup Z''_m \sqcup Z'''_m$ where

$$Z'_m = \{(x_{i+1}, w(x_i)) \mid i \in [1, k - 1]\} \sqcup \{(x_1, S(y))\} \sqcup \{(y, w(x_k))\},$$

$$Z''_m = \{(x_i, w(x_i)) \mid i \in [1, k]\}, \quad Z'''_m = \{(y, S(y))\}.$$

The set Z_m is called the zigzag labeled m . Then $w \cap Z_m = Z''_m$ and $S \cap Z_m = Z'''_m$. We set $\text{bk}_S(w) = \bigsqcup_{m \in \mathbb{Z}} Z'_m$. Then $\text{bk}_S(w)$ is a partial permutation the width of whose Shi poset is equal to the density of S .

The map $\Psi : \bigsqcup_{\lambda \vdash n} \text{RSYT}(\lambda) \times \text{RSYT}(\lambda) \times \widetilde{\mathcal{S}}_n \rightarrow \widetilde{\mathcal{S}}_n$ is defined as follows. For $P = (P_1, \dots, P_l)$, $Q = (Q_1, \dots, Q_l)$, and $\#p = (\rho_1, \dots, \rho_l)$, we define S_i to be the unique stream which maps $Q_i + n\mathbb{Z}$ to $P_i + n\mathbb{Z}$ and which satisfies $a(S_i) = \rho_i$. Now starting with the empty partial permutation w_l we successively define $w_{i-1} = \text{bk}_{S_i}(w_i)$ for $i \in [1, l]$. This process is well-defined and we set $\Psi(P, Q, \#p) = w_0$. (Here the dominance of $\#p$ is not required.)

5.2. CHANNELS AND RIVERS. Let us describe some properties of channels and rivers. We start with observing the following lemma which strengthens [6, Proposition 3.17].

LEMMA 5.4. *Suppose that C_1, C_2, C_3 are three (not necessarily disjoint) channels of a partial permutation w . Then we have $h(C_1, C_3) \leq h(C_1, C_2) + h(C_2, C_3)$. Furthermore, if $h(C_1, C_2) = 0$, $h(C_2, C_3) = 0$, or $C_1 \leq_{\text{SW}} C_2 \leq_{\text{SW}} C_3$, then we have $h(C_1, C_2) + h(C_2, C_3) = h(C_1, C_3)$.*

Proof. It is indeed shown in the proof of [6, Proposition 3.17]. □

For a partial permutation of w whose Shi poset is of width r , we successively choose C_i to be the southwest channel of $w - \bigsqcup_{j=1}^{i-1} C_j$ until we fail to find a stream of density r and let $\{C_1, \dots, C_m\}$ be its result. (This is always possible due to [6, Proposition 3.13].) Then (C_1, \dots, C_m) is a disjoint collection of channels of w where $C_1 \leq_{\text{SW}} \dots \leq_{\text{SW}} C_m$. By [6, Definition 3.7] and the argument thereafter, we see that m equals the multiplicity of r (the width of P_w) in the Greene–Kleitman partition of P_w . This argument (and its reflection along the anti-diagonal of the xy -plane) shows

the existence, but not uniqueness, of the following sequence. (Throughout this paper only the existence of such a sequence is needed.)

DEFINITION 5.5. For a partial permutation w , we denote by $\mathfrak{C}_w = (C_1, \dots, C_m)$ a sequence of the channels of w such that

- (1) m is the multiplicity of λ_1 in the Greene–Kleitman partition λ of P_w ,
- (2) $C_1 \leq_{\text{SW}} C_2 \leq_{\text{SW}} \dots \leq_{\text{SW}} C_m$,
- (3) C_i 's are pairwise disjoint, and
- (4) this sequence contains all the northeast channels of rivers of w .

Similarly, we denote by $\mathfrak{C}'_w = (C'_1, \dots, C'_m)$ a similar sequence which satisfies (1), (2), (3), and

- (4') it contains all the southwest channels of rivers of w .

REMARK 5.6. By Lemma 5.4, for a given w and two sequences $\mathfrak{C}_w = (C_1, \dots, C_m)$ and $\mathfrak{C}'_w = (C'_1, \dots, C'_m)$ as above, there exist $0 = m_0, m_1, m_2, \dots, m_l = m$ where $0 < m_1 < m_2 < \dots < m_{l-1} < m$ such that $h(C_i, C_j) = h(C'_i, C'_j) = 0$ if and only if $i, j \in [1 + m_{k-1}, m_k]$ for some $k \in [0, l - 1]$. In other words, there are l rivers $R_1, \dots, R_l \subset w$ such that $C_i, C'_i \in R_k$ for $k \in [1, l]$ and $i \in [1 + m_{k-1}, m_k]$. Then by maximality of \mathfrak{C}_w (resp. \mathfrak{C}'_w), the intersection of $C_{1+m_{k-1}}$ and the southwest channel of R_k (resp. the intersection of C'_{m_k} and the northeast channel of R_k) is nonempty. In particular, the channel numbering corresponding to the southwest (resp. northeast) channel of R_k is the same as that of $C_{1+m_{k-1}}$ (resp. C'_{m_k}). In addition, when $m_{k-1} + 1 < m_k$ it is possible to assume that both \mathfrak{C}_w and \mathfrak{C}'_w contain both the southwest and northeast channel of R_k . (When $m_{k-1} + 1 = m_k$, the northeast and the southwest channel of R_k have nonempty intersection but they may differ in general.)

EXAMPLE 5.7. Suppose that $w = [6, 1, 18, 3, 19, 24, 12, 15, 17, 10]$. Then there are four channels

$$\begin{aligned} C_1 &= [\emptyset, 1, \emptyset, 3, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset, 10], \\ C'_2 &= [6, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset, 12, 15, \emptyset, \emptyset], \\ C_2 &= [\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset, 12, 15, 17, \emptyset], \\ C_3 &= [\emptyset, \emptyset, 18, \emptyset, 19, 24, \emptyset, \emptyset, \emptyset, \emptyset]. \end{aligned}$$

Here $\{C_1, C_2, C'_2\}$ and $\{C_3\}$ are rivers of w . Also the sequence (C_1, C_2, C_3) satisfies the properties of both \mathfrak{C}_w and \mathfrak{C}'_w .

Suppose that $m = \#\mathfrak{C}_w \geq 2$. Then by [6, Corollary 14.11], there exists a sequence $\mathfrak{C}_{\text{fw}(w)} = (D_1, \dots, D_{m-1})$ as in Definition 5.5 such that each D_i is between C_i and C_{i+1} . (When $\#\mathfrak{C}_w = 1$, the width of $P_{\text{fw}(w)}$ is strictly smaller than that of P_w .) If we let $\Phi(w) = (P, Q, \overset{\#}{\rho}) = (P, Q, \overset{\#}{\mathbb{S}}_{P,Q} + \overset{\#}{\rho})$ where $\overset{\#}{\rho} = (\rho_1, \rho_2, \dots)$, $\overset{\#}{\mathbb{S}}_{P,Q} = (s_1, s_2, \dots)$, and $\overset{\bullet}{\rho} = (\bullet\rho_1, \bullet\rho_2, \dots)$, then [6, Theorem 8.1] shows that $h(C_1, C_2) = \rho_2 - \rho_1 - s_2 = \bullet\rho_2 - \bullet\rho_1$ and $h(C_i, C_{i+1}) = h(D_{i-1}, D_i)$ for $i \in [2, m - 1]$.

The distances between channels can be obtained from AMBC due to the following lemma.

LEMMA 5.8. Suppose that $\Phi(w) = (P, Q, \overset{\#}{\mathbb{S}}_{P,Q} + \overset{\#}{\rho})$ where $\overset{\#}{\rho} = (\bullet\rho_1, \bullet\rho_2, \dots)$ and $\mathfrak{C}_w = (C_1, \dots, C_m)$. Then for $i \in [1, m - 1]$ we have $h(C_i, C_{i+1}) = \bullet\rho_{i+1} - \bullet\rho_i$.

Proof. When $i = 1$ we already observed that $h(C_1, C_2) = \bullet\rho_2 - \bullet\rho_1$ which follows from [6, Theorem 8.1]. However, the second part of the aforementioned theorem states that $h(C_i, C_{i+1}) = h(D_{i-1}, D_i)$ where C_i, D_j are as above, and thus the result follows from induction on i and the fact that $\Phi(\text{fw}(w)) = (P_{\geq 2}, Q_{\geq 2}, (\overset{\#}{\mathbb{S}}_{P,Q} + \overset{\#}{\rho})_{\geq 2})$. \square

In particular, the indices $0 = m_0, m_1, m_2, \dots, m_l = m$ in the above remark are chosen in a way that $\bullet\rho_i = \bullet\rho_j$ if and only if $i, j \in [1 + m_{k-1}, m_k]$ for some $k \in [1, l]$, i.e. each river of w corresponds to a subset of $[1, m]$ having the same values of $\# \bullet\rho$.

For a partial permutation w and a compatible stream S , the indexing river of $\text{bk}_S(w)$ corresponding to (w, S) is defined to be the collection of channels $C \in \text{bk}_S(w)$ such that (if we fix a shift of $d_{\text{bk}_S(w)}^C$ properly then) $d_{\text{bk}_S(w)}^C(b) = m$ for any $b \in \text{bk}_S(w)$ if and only if $b \in Z'_m$ where Z'_m is as in 5.1 (the set of elements labeled m). Then the collection is indeed a river. Moreover, when $(w, S) = (\text{fw}(v), \text{st}(v))$ for some partial permutation v then the southwest channel of v is contained in the indexing river of (w, S) since the forward numbering (i.e. the southwest channel numbering) and the backward numberings are compatible in this case.

For a partial permutation w and a compatible stream S with the proper numbering $\tilde{d} : S \rightarrow \mathbb{Z}$ and $d = d_w^{\text{bk}, S} : w \rightarrow \mathbb{Z}$, we say that $(x, w(x)) \in w$ is N -terminal (resp. W -terminal) with respect to S if there exists $(y, S(y)) \in S$ such that $\tilde{d}(y, S(y)) = d(x, w(x)) + 1$ and $x < y$ (resp. $w(x) < w(y)$). Also we say that $b \in w$ is N -terminating (resp. W -terminating) with respect to S if there exists a path $b = b_0, \dots, b_k \in w$ such that b_k is N -terminal (resp. W -terminal) and $d(b_{i-1}) + 1 = d(b_i)$ for $i \in [1, k]$.

By [6, Lemma 14.22] and [6, Lemma 16.15], if S is a stream compatible with w and R is a river of w then either every $b \in \bigcup_{C \in R} C$ is N -terminating or every $b \in \bigcup_{C \in R} C$ is W -terminating, thus we may say that the river R is either N -terminating or W -terminating. Furthermore, by [6, Corollary 16.17] at most one river of w is both N - and W -terminating. These notions are related to the indexing river of $\text{bk}_S(w)$ as described in the lemma below. (Also see Figure 2.)

LEMMA 5.9. *Suppose that v is a partial permutation, S is a stream compatible with v , and $w = \text{bk}_S(v)$. Then $(x, w(x)) \in w$ is in the indexing river corresponding to (v, S) if and only if in the zigzag Z containing $(x, w(x))$ (in the backward step of AMBC) the elements $(y, v(y)) \in Z \cap v$ such that $x < y$ (resp. $w(x) < v(y)$) are W -terminating (resp. N -terminating.)*

Proof. The if part follows directly from [6, Remark 16.7]. For the converse, suppose that $(x, w(x))$ is in the indexing river, say R . Let $(c, w(c)), (c', w(c')) \in w$ be such that c is the minimum among the elements in $\bigcup_{C \in R} C \cap Z$ and c' is the maximum. Then we have $(c', w(c')) \leq_{SW} (x, w(x)) \leq_{SW} (c, w(c))$. By [6, Lemma 16.5, 16.8] and their “reflection statements along the anti-diagonal” (or equivalently considering w^{-1} instead), if we let $(y, v(y)) \in v$ (resp. $(y', v(y')) \in v$) be the most northeast W -terminating element (resp. the most southwest N -terminating element) of $Z \cap v$ then $c < y$ and $w(c) = v(y)$ (resp. $c' = y'$ and $w(c') < v(y')$). (If no such $(y, v(y))$ exists then $(c, w(c)) = (c', w(c')) = (x, w(x))$ is the most southwest element of Z , and if no such $(y', v(y'))$ exists then $(c, w(c)) = (c', w(c')) = (x, w(x))$ is the most northeast element of Z .) Thus the result follows. \square

As a result, elements of v which are northeast of the indexing river of $\text{bk}_S(v)$ corresponding to (v, S) are N -terminating but not W -terminating, and a similar statement holds for balls southwest of the indexing river.

EXAMPLE 5.10. Figure 2 describes a zigzag which appears in the backward step of AMBC. Here, the striped ball is an element of S , the white ones are the elements of v , and the shaded ones are the elements of $w = \text{bk}_S(v)$. The curvy lines denote channels in the indexing river. The elements $C = (c, w(c)), C' = (c', w(c'))$ in Lemma 5.9 are depicted as in the diagram, and the white balls labeled N (resp. W , resp. NW) are N -terminating but not W -terminating (resp. W -terminating but not N -terminating, resp. both N - and W -terminating).




FIGURE 2. Indexing river and N/W -terminating elements

5.3. PROPER PARTIAL ROTATION AND AMBC. Here we define (proper) partial rotations and study their interactions with AMBC.

DEFINITION 5.11. For a partial permutation w and a stream $S \subset w$, let us set x_i for $i \in \mathbb{Z}$ so that $S = \{(x_i, w(x_i)) \mid i \in \mathbb{Z}\}$ and $i < j$ if and only if $x_i < x_j$. We define the partial rotation on w with respect to S , denoted $\text{pr}_S(w)$, as follows.

$$\text{pr}_S(w)(a) = \begin{cases} w(x_{i+1}) & \text{if } a = x_i, \\ w(a) & \text{otherwise.} \end{cases}$$

Likewise, we also define the inverse partial rotation $\text{pr}_S^{-1}(w)$ as follows.

$$\text{pr}_S^{-1}(w)(a) = \begin{cases} w(x_{i-1}) & \text{if } a = x_i, \\ w(a) & \text{otherwise.} \end{cases}$$

We often abbreviate the partial rotation (resp. inverse partial rotation) to PR (resp. IPR). We say that the partial rotation pr_S (resp. the inverse partial rotation pr_S^{-1}) applied to w is proper if S is the northeast (resp. southwest) channel of some river of w .

In [6], the term *shift* is used in order to denote the PR/IPRs. Moreover, the notations $w\langle 1 \rangle_S$, $w\langle -1 \rangle_S$, $S\langle 1 \rangle$, and $S\langle -1 \rangle$ in [6] are translated to $\text{pr}_S(w)$, $\text{pr}_S^{-1}(w)$, $\text{pr}_S(S)$, and $\text{pr}_S^{-1}(S)$ in our language. When R is a river of w and C (resp. C') is the northeast (resp. southwest) channel of R , then $w\langle 1 \rangle_R$ and $w\langle -1 \rangle_R$ in [6] are equal to $\text{pr}_C(w)$ (resp. $\text{pr}_{C'}^{-1}(w)$) which is by definition a proper partial rotation (resp. a proper inverse partial rotation). We stick to our notations in this paper, but allow ourselves to use $S\langle 1 \rangle$ and $S\langle -1 \rangle$ instead of $\text{pr}_S(S)$ and $\text{pr}_S^{-1}(S)$.

Let us describe the relation between proper PR/IPRs and AMBC.

THEOREM 5.12. *For a partial permutation w with a sequence $\mathfrak{C}_w = (C_1, \dots, C_m)$ (resp. $\mathfrak{C}'_w = (C'_1, \dots, C'_m)$) as in Definition 5.5, assume that R is a river of w whose northeast (resp. southwest) channel is C_r for some r (resp. $C'_{r'}$ for some r'). Note that both pr_{C_r} and $\text{pr}_{C'_{r'}}^{-1}$ are proper with respect to w . Suppose that $\Phi(w) = (P, Q, \#p)$. Then we have*

$$\Phi(\text{pr}_{C_r}(w)) = (P, Q, \#p + \#e_r) \quad \text{and} \quad \Phi(\text{pr}_{C'_{r'}}^{-1}(w)) = (P, Q, \#p - \#e_{r'}).$$

Here, $\#e_i$ is a vector whose i -th coordinate is 1 and 0 elsewhere.

Proof. We first prove $\Phi(\text{pr}_{C_r}(w)) = (P, Q, \#p + \#e_r)$. Assume that $r = 1$. Let us write $(v, S) = (\text{fw}(w), \text{st}(w))$ so that $w = \text{bk}_S(v)$, $\Phi(v) = (P_{\geq 2}, Q_{\geq 2}, \#p_{\geq 2})$, and $a(S) = \rho_1$. Since C_1 has a nontrivial intersection with the southwest channel of w , R is the indexing river of w corresponding to (v, S) . Thus by [6, Theorem 16.9] we have $\text{pr}_{C_1}(w) = \text{bk}_{S\langle 1 \rangle}(v)$. Since $h(C_1, C_2) > 0$ by assumption, we have $\bullet\rho_1 < \bullet\rho_2$ by Lemma 5.8, which implies that $\#p + \#e_1$ is still dominant with respect to the pair (P, Q) . Thus from the definition of backward algorithm we see that $\Phi(\text{pr}_{C_1}(w)) = \Phi(\text{bk}_{S\langle 1 \rangle}(v)) = (P, Q, \#p + \#e_1)$ (note that $a(S\langle 1 \rangle) = \rho_1 + 1$) as desired.

In general, we successively define $(v_1, S_1) = (\text{fw}(w), \text{st}(w))$ and $(v_i, S_i) = (\text{fw}(v_{i-1}), \text{st}(v_{i-1}))$ for $i \in [2, r]$. Let $C_r = D^0 \subset w, D^1 \subset v_1, \dots, D^r \subset v_r$ be the $(m + 1 - r)$ -th northeast channel in $\mathfrak{C}_w, \mathfrak{C}_{v_1}, \dots, \mathfrak{C}_{v_r}$, respectively. Note that each of them is the northeast channel of some river, say $D^i \in R^i$, by Lemma 5.8 since so is $C_r = D^0 \in R^0 = R$.

The above argument for $r = 1$ case shows that $\text{pr}_{D^{r-1}}(v_{r-1}) = \text{bk}_{S_{r-1}\langle 1 \rangle}(v_r)$ and thus $\Phi(\text{pr}_{D^{r-1}}(v_{r-1})) = (P_{\geq r}, Q_{\geq r}, (\#p + \#e_r)_{\geq r})$. Now recall that the southwest channel of v_{r-2} is in the indexing river corresponding to (v_{r-1}, S_{r-1}) . In particular, any ball in v_{r-1} is N -terminating by Lemma 5.9. Thus (the reflection statement along the anti-diagonal of) [6, Proposition 16.39] shows that $\text{bk}_{S_{r-1}}(\text{pr}_{D^{r-1}}(v_{r-1})) = \text{pr}_{D^{r-2}}(v_{r-2})$. (Here, D^{r-2} is the northeast channel of R^{r-2} , and R^{r-2} contains the most southwest channel among the ones of v_{r-2} that are northeast of D^{r-1} .) Thus it follows that $\Phi(\text{pr}_{D^{r-2}}(v_{r-2})) = \Phi(\text{bk}_{S_{r-1}}(\text{pr}_{D^{r-1}}(v_{r-1}))) = (P_{\geq r-1}, Q_{\geq r-1}, (\#p + \#e_r)_{\geq r-1})$.

Now we iterate this process and eventually reach the conclusion that $\text{pr}_{C_r}(w) = \text{bk}_{S_1}(\text{pr}_{D^1}(v_1)) = (\text{bk}_{S_1} \circ \text{bk}_{S_2})(\text{pr}_{D^2}(v_2)) = \dots = (\text{bk}_{S_1} \circ \dots \circ \text{bk}_{S_{r-1}} \circ \text{bk}_{S_r\langle 1 \rangle})(v_r)$. Therefore from the definition of backward algorithm we see that $\Phi(\text{pr}_{C_r}(w)) = \Phi((\text{bk}_{S_1} \circ \dots \circ \text{bk}_{S_{r-1}} \circ \text{bk}_{S_r\langle 1 \rangle})(v_r)) = (P, Q, \#p + \#e_r)$ as desired. (Here we use the fact that $\#p + \#e_r$ is dominant with respect to (P, Q) .)

To prove the second claim $\Phi(\text{pr}_{C'_{r'}}^{-1}(w)) = (P, Q, \#p - \#e_{r'})$ one may similarly argue as above. The case when $r' = 1$ is almost the same as the above, and for the inductive argument one needs to use (the reflection statement along the anti-diagonal of) [6, Proposition 16.38] instead of [6, Proposition 16.39]. We omit the details. \square

EXAMPLE 5.13. Suppose that $w = [6, 1, 18, 3, 19, 24, 12, 15, 17, 10] \in \widetilde{\mathcal{S}}_{10}$ as in Example 5.7. Then we have

$$\begin{aligned} \Phi(w) &= (P, Q, \#p) \\ &= (((1, 3, 10), (2, 5, 6), (4, 7, 9), (8)), ((3, 5, 6), (7, 8, 9), (1, 4, 10), (2)), (2, 3, 2, 0)) \end{aligned}$$

and $\#p_{P,Q} = (0, 1, 1, 0) - (0, 0, 2, 0) = (0, 1, -1, 0)$. Thus $\#p = \#p - \#p_{P,Q} = (2, 2, 3, 0)$. (This again shows that there are two rivers $R_1 \ni C_1, C_2$ and $R_2 \ni C_3$.) If we calculate

$\text{pr}_C(w)$ and $\text{pr}_C^{-1}(w)$ for each $C \in \mathfrak{C}_w$ when the operations are proper then

$$\begin{aligned} \text{pr}_{C_1}^{-1}(w) &= [6, 0, 18, 1, 19, 24, 12, 15, 17, 3] \\ &\xrightarrow{\Phi} (((1, 3, 10), (2, 5, 6), (4, 7, 9), (8)), ((3, 5, 6), (7, 8, 9), (1, 4, 10), (2)), (1, 3, 2, 0)), \\ \text{pr}_{C_2}(w) &= [6, 1, 18, 3, 19, 24, 15, 17, 22, 10] \\ &\xrightarrow{\Phi} (((1, 3, 10), (2, 5, 6), (4, 7, 9), (8)), ((3, 5, 6), (7, 8, 9), (1, 4, 10), (2)), (2, 4, 2, 0)), \\ \text{pr}_{C_3}(w) &= [6, 1, 14, 3, 18, 19, 12, 15, 17, 10] \\ &\xrightarrow{\Phi} (((1, 3, 6, 0), (2, 5, 9), (4, 7), (8)), ((3, 5, 6, 9), (4, 7, 8), (1, 10), (2)), (2, 3, 1, 0)), \\ \text{pr}_{C_3}^{-1}(w) &= [6, 1, 19, 3, 24, 28, 12, 15, 17, 10] \\ &\xrightarrow{\Phi} (((1, 3, 6, 0), (2, 5, 9), (4, 7), (8)), ((3, 5, 6, 9), (4, 7, 8), (1, 10), (2)), (2, 3, 3, 0)), \end{aligned}$$

as expected.

5.4. PROPER PARTIAL ROTATION AND SIGN INSERTION. Here we show that the proper PR/IPRs do not affect the result of sign insertion as the following theorem states.

THEOREM 5.14. *Suppose that pr_C and $\text{pr}_{C'}^{-1}$ are proper with respect to $w \in \widetilde{\mathcal{S}}_n$ for some $C, C' \subset w$. Then we have $\text{sgn}_\Omega(w) = \text{sgn}_\Omega(\text{pr}_C(w)) = \text{sgn}_\Omega(\text{pr}_{C'}^{-1}(w))$.*

The main tool to prove this theorem is the following Diamond Lemma discussing interactions between right star operations and proper PR/IPRs.

LEMMA 5.15 (Diamond lemma). *Suppose that we are given a partial permutation w with $\mathfrak{C}_w = (C_1, \dots, C_m)$, $p \in [1, n]$, and $q \in [1, m]$. Assume that:*

- *Either $w(p-1)$ is between $w(p)$ and $w(p+1)$ so that a right star operation $w \mapsto w^*$ that switches $w(p)$ and $w(p+1)$ is well-defined, or $w(p+1)$ is between $w(p-1)$ and $w(p)$ so that a right star operation $w \mapsto w^*$ that switches $w(p-1)$ and $w(p)$ is well-defined. Let us call it “the right star operation centered at p ”.*
- *C_q is a northeast channel of a river of w .*

Then, we have:

- *The q -th southwest channel C_q^* of $\mathfrak{C}_{w^*} = (C_1^*, \dots, C_m^*)$, where \mathfrak{C}_{w^*} is as in Definition 5.5, is a northeast channel of a river of w^* .*
- *The right star operation centered at p is well-defined for $\text{pr}_{C_q}(w)$, say $\text{pr}_{C_q}(w)^*$.*

Furthermore, we have $\text{pr}_{C_q}(w)^* = \text{pr}_{C_q^*}(w^*)$. In other words, we may “canonically complete the commutative diagram” when we are given $w \mapsto w^*$ and $w \mapsto \text{pr}_{C_q}(w)$. The same result also holds when one replaces proper partial rotations with proper inverse partial rotations and/or replaces right star operations with multiplication by ω on the right.

Since star operations and multiplication by ω preserve each two-sided cell, or equivalently the Greene–Kleitman partition of a Shi poset, it makes sense to take the q -th southwest channel in \mathfrak{C}_{w^*} and $\mathfrak{C}_{w \cdot \omega}$ in the lemma above. Also note that the statement for multiplication by ω on the right is straightforward because the channels of $w \cdot \omega$ are simply the shifts of those of w .

EXAMPLE 5.16. Let $w = [6, 1, 18, 3, 19, 24, 12, 15, 17, 10]$ as in Example 5.7 and 5.13. Then the star operation $w \mapsto w^*$ centered at 7 is well defined and $w^* = [6, 1, 18, 3, 19, 12, 24, 15, 17, 10]$. Let us consider

$$\text{pr}_{C_2}(w) = [6, 1, 18, 3, 19, 24, 15, 17, 22, 10].$$

Then $\text{pr}_{C_2}(w) \mapsto \text{pr}_{C_2}(w)^*$ centered at 7 is well defined and

$$\text{pr}_{C_2}(w)^* = [6, 1, 18, 3, 19, 15, 24, 17, 22, 10].$$

Moreover, $w^* \mapsto \text{pr}_{C_2}(w)^*$ is the proper partial rotation with respect to the channel $[\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, 12, \emptyset, 15, 17, \emptyset]$.

First let us discuss how to use Lemma 5.15 to prove Theorem 5.14.

Proof of Theorem 5.14, assuming Lemma 5.15. It suffices only to prove for the proper partial rotation, i.e. $\text{sgn}_\Omega(w) = \text{sgn}_\Omega(\text{pr}_C(w))$. Let us set $w_0 = w$, $\mathfrak{P}_0 = \Omega_0 = \emptyset$ and sequentially define $(i, \mathfrak{P}_i, \Omega_i, w_i) = F(i - 1, \mathfrak{P}_{i-1}, \Omega_{i-1}, w_{i-1})$ for $i \in [1, N]$ where $w_N = \emptyset$. Thus in particular we have $\text{sgn}_{\mathfrak{P}}(w) = \mathfrak{P}_N$ and $\text{sgn}_\Omega(w) = \Omega_N$. Let us set $v_i = \mathfrak{P}_i \cap w_i$ for $i \in [0, N]$. Note that either $v_i = v_{i-1}$ or v_i is obtained from v_{i-1} by applying right star operations followed by multiplication by ω on the right (cf. the proof of Lemma 4.3).

Let $\tilde{w} = \text{pr}_C(w)$ where pr_C applied to w is proper. Then by Lemma 5.15, we may (uniquely) find $\tilde{w} = \tilde{v}_0, \tilde{v}_1, \dots, \tilde{v}_N$ such that $v_i \mapsto \tilde{v}_i$ is a proper partial rotation, $v_i = v_{i-1}$ if and only if $\tilde{v}_i = \tilde{v}_{i-1}$, and if $\tilde{v}_i \neq \tilde{v}_{i-1}$ then \tilde{v}_i is obtained from \tilde{v}_{i-1} by a series of right star operations followed by multiplication by ω on the right. We claim that there exist $\tilde{\mathfrak{P}}_i, \tilde{w}_i$ for $i \in [0, N]$ such that $\tilde{v}_i = \tilde{\mathfrak{P}}_i \cap \tilde{w}_i$ for $i \in [0, N]$ and $(i, \tilde{\mathfrak{P}}_i, \tilde{\Omega}_i, \tilde{w}_i) = F(i - 1, \tilde{\mathfrak{P}}_{i-1}, \tilde{\Omega}_{i-1}, \tilde{w}_{i-1})$ for $i \in [1, N]$. (In particular the lengths of $\tilde{\mathfrak{P}}_i, \mathfrak{P}_i$, and Ω_i are all equal.)

We prove by induction on i . It is obvious when $i = 0$ by setting $\mathfrak{P}_0 = \emptyset$ and $\tilde{w}_0 = \tilde{w}$, so suppose that it is true up to $i - 1$. Let p be the length of \mathfrak{P}_{i-1} . Then we have $v_{i-1}(1) < \dots < v_{i-1}(p)$ and $\tilde{v}_{i-1}(1) < \dots < \tilde{v}_{i-1}(p)$. If $v_{i-1}(p) < v_{i-1}(p + 1)$, then it means that $v_{i-1} = v_i$, i.e. $v_{i-1} \mapsto v_i$ is a trivial step, which implies $\tilde{v}_{i-1} = \tilde{v}_i$ by assumption. We claim that $\tilde{v}_{i-1}(p) < \tilde{v}_{i-1}(p + 1)$ so that we may set $\tilde{\mathfrak{P}}_i$ and \tilde{w}_i similar to \mathfrak{P}_i and w_i . Indeed, if $\tilde{v}_{i-1}(p) > \tilde{v}_{i-1}(p + 1)$ then it is possible to apply the right star operation centered at p on \tilde{v}_{i-1} . However, by Lemma 5.15, it means that the same star operation can be applied to v_{i-1} , which is a contradiction.

It remains to consider the case when $v_{i-1}(p) > v_{i-1}(p + 1)$. Then v_i is obtained from v_{i-1} by applying the right star operations centered at $p, p - 1, \dots, 2$, respectively, followed by multiplication by ω on the right. By Lemma 5.15, the same process applies to \tilde{v}_{i-1} to get \tilde{v}_i . However, as $\tilde{v}_{i-1}(1) < \dots < \tilde{v}_{i-1}(p)$ and $\tilde{v}_{i-1}(p) > \tilde{v}_{i-1}(p + 1)$ by assumption, one can easily show that this is the usual “bumping process” (which inserts $\tilde{v}_{i-1}(p + 1)$ to $\tilde{\mathfrak{P}}_{i-1}$) followed by multiplication by ω on the right. This is clearly a valid step of the sign insertion algorithm. Thus, if we set $\tilde{\mathfrak{P}}_i$ to be the first p -th characters of \tilde{v}_i and \tilde{w}_i to be the remainder then the induction step is also valid in this case. It suffices for the proof. \square

EXAMPLE 5.17. We compare the sign insertion of $w = [6, 1, 18, 3, 19, 24, 12, 15, 17, 10]$ and $\text{pr}_{C_2}(w) = [6, 1, 18, 3, 19, 24, 15, 17, 22, 10]$. The underlined numbers are where the proper partial rotation is applied. It is clear that the sign insertion processes for w and $\text{pr}_{C_2}(w)$ are parallel.

i	\mathfrak{P}	Ω	w
0	\emptyset	\emptyset	6 1 18 3 19 24 <u>12</u> <u>15</u> <u>17</u> 10
1	6	1	1 18 3 19 24 <u>12</u> <u>15</u> <u>17</u> 10
2	1	1	18 3 19 24 <u>12</u> <u>15</u> <u>17</u> 10 16
3	1 18	1 3	3 19 24 <u>12</u> <u>15</u> <u>17</u> 10 16
4	1 3	1 3	19 24 <u>12</u> <u>15</u> <u>17</u> 10 16 28
5	1 3 19	1 3 5	24 <u>12</u> <u>15</u> <u>17</u> 10 16 28
6	1 3 19 24	1 3 5 6	<u>12</u> <u>15</u> <u>17</u> 10 16 28
7	1 3 <u>12</u> 24	1 3 5 6	<u>15</u> <u>17</u> 10 16 28 29

8	1 3 <u>12</u> <u>15</u>	1 3 5 6	<u>17</u> 10 16 28 29 34
9	1 3 <u>12</u> <u>15</u> <u>17</u>	1 3 5 6 7	10 16 28 29 34
10	1 3 10 <u>15</u> <u>17</u>	1 3 5 6 7	16 28 29 34 <u>22</u>
11	1 3 10 <u>15</u> 16	1 3 5 6 7	28 29 34 <u>22</u> <u>27</u>
12	1 3 10 <u>15</u> 16 28	1 3 5 6 7 12	29 34 <u>22</u> <u>27</u>
13	1 3 10 <u>15</u> 16 28 29	1 3 5 6 7 12 13	34 <u>22</u> <u>27</u>
14	1 3 10 <u>15</u> 16 28 29 34	1 3 5 6 7 12 13 14	<u>22</u> <u>27</u>
15	1 3 10 <u>15</u> 16 <u>22</u> 29 34	1 3 5 6 7 12 13 14	<u>27</u> 38
16	1 3 10 <u>15</u> 16 <u>22</u> <u>27</u> 34	1 3 5 6 7 12 13 14	38 39
17	1 3 10 <u>15</u> 16 <u>22</u> <u>27</u> 34 38	1 3 5 6 7 12 13 14 17	39
18	1 3 10 <u>15</u> 16 <u>22</u> <u>27</u> 34 38 39	1 3 5 6 7 12 13 14 18	\emptyset

i	\mathfrak{P}	\mathfrak{Q}	w
0	\emptyset	\emptyset	6 1 18 3 19 24 <u>15</u> <u>17</u> <u>22</u> 10
1	6	1	1 18 3 19 24 <u>15</u> <u>17</u> <u>22</u> 10
2	1	1	18 3 19 24 <u>15</u> <u>17</u> <u>22</u> 10 16
3	1 18	1 3	3 19 24 <u>15</u> <u>17</u> <u>22</u> 10 16
4	1 3	1 3	19 24 <u>15</u> <u>17</u> <u>22</u> 10 16 28
5	1 3 19	1 3 5	24 <u>15</u> <u>17</u> <u>22</u> 10 16 28
6	1 3 19 24	1 3 5 6	<u>15</u> <u>17</u> <u>22</u> 10 16 28
7	1 3 <u>15</u> 24	1 3 5 6	<u>17</u> <u>22</u> 10 16 28 29
8	1 3 <u>15</u> <u>17</u>	1 3 5 6	<u>22</u> 10 16 28 29 34
9	1 3 <u>15</u> <u>17</u> <u>22</u>	1 3 5 6 7	10 16 28 29 34
10	1 3 10 <u>17</u> <u>22</u>	1 3 5 6 7	16 28 29 34 <u>25</u>
11	1 3 10 16 <u>22</u>	1 3 5 6 7	28 29 34 <u>25</u> <u>27</u>
12	1 3 10 16 <u>22</u> 28	1 3 5 6 7 12	29 34 <u>25</u> <u>27</u>
13	1 3 10 16 <u>22</u> 28 29	1 3 5 6 7 12 13	34 <u>25</u> <u>27</u>
14	1 3 10 16 <u>22</u> 28 29 34	1 3 5 6 7 12 13 14	<u>25</u> <u>27</u>
15	1 3 10 16 <u>22</u> <u>25</u> 29 34	1 3 5 6 7 12 13 14	<u>27</u> 38
16	1 3 10 16 <u>22</u> <u>25</u> <u>27</u> 34	1 3 5 6 7 12 13 14	38 39
17	1 3 10 16 <u>22</u> <u>25</u> <u>27</u> 34 38	1 3 5 6 7 12 13 14 17	39
18	1 3 10 16 <u>22</u> <u>25</u> <u>27</u> 34 38 39	1 3 5 6 7 12 13 14 18	\emptyset

5.5. PROOF OF DIAMOND LEMMA. It remains to prove Lemma 5.15. Recall the definition of \mathfrak{C}_w given in Definition 5.5.

LEMMA 5.18. *For a partial permutation w , right star operations and multiplication by ω on the right do not change the distances between channels in \mathfrak{C}_w .*

Proof. This follows from Lemma 5.8 together with the last part of Section 3. (Here, it is crucial that $\delta(T, s) = (\delta_1, \dots, \delta_l)$ satisfies that $\delta_i = \delta_j$ whenever $\lambda_i = \lambda_j$.) \square

The following lemma describes the reason why the proper PR/IPRs are useful.

LEMMA 5.19. *If $C \subset w \in \widetilde{\mathcal{S}}_n$ is a channel then $\text{pr}_C(w)$ (resp. $\text{pr}_C^{-1}(w)$) and w are in the same two-sided cell if and only if S is the northeast (resp. southwest) channel of a river of w . (In this case such an element is in the same left and right cell of w by Theorem 5.12.)*

Proof. Here we prove only for partial rotations; the inverse case can be proved similarly. The if part follows from Theorem 5.12 and it remains to prove the only if part. Assume that $C \subset w$ is a channel which is not the northeast in its river, but its partial rotation yields a permutation in the same two-sided cell. By assumption there exists a channel $C' \neq C$ such that $C \leq_{SW} C'$ and $h(C, C') = 0$, i.e. d^C and $d^{C'}$ are equal up to shift by [6, Definition 3.16]. We fix shifts of d^C and $d^{C'}$ such that they coincide, and let $c_i \in C$ and $b_i \in C'$ be balls such that $d^C(c_i) = d^{C'}(c_i) = d^C(b_i) = d^{C'}(b_i) = i$

for $i \in \mathbb{Z}$. Note that by assumption either $b_i = c_i$ or $c_i \leq_{SW} b_i$, and there exists $i \in \mathbb{Z}$ such that $b_i \neq c_i$.

We claim that there exists $j \in \mathbb{Z}$ such that $b_j = (y, w(y)) \in C' - C$ (thus $c_j \leq_{SW} b_j$) and $c_{j-1} = (x, w(x)) \in C$ satisfies $x < y$ and $w(x) < w(y)$. If $c_{j-1} = b_{j-1}$ then the claim is obvious, thus it suffices to consider the case when $C \cap C' = \emptyset$. However, if the assumption is false for any $j \in \mathbb{Z}$ then we may lower the labels of the balls in C' by 1 and the numbering still satisfies the monotone property, which is a contradiction. Thus such j always exists.

Now for $i \in \mathbb{Z}$ let d_i be a ball in the partial rotation of C so that c_i and d_i share the same x -coordinate, or equivalently the y -coordinate of d_i is the same as that of c_{i+1} . Let $b_j \in C' - C$ be as above. Then it follows that x -coordinate of b_j is between those of c_{j-1} and c_j and the y -coordinate of b_j is bigger than that of c_j , which in turn means that d_{j-1} is northeast of b_j . In addition, similar argument shows that there exists $k \in \mathbb{Z}$ such that b_k is northeast of d_k .

By adjusting the shift of d^C and $d^{C'}$ necessary, we may assume that $1 \leq j \leq k \leq r$ where r is the density of C . Then the conditions above imply that $(\{b_i \mid j \leq i \leq k\} \cup \{d_i \mid 1 \leq i \leq j - 1 \text{ or } k \leq i \leq r\}) + \mathbb{Z}(n, n) \subset w$ is a stream of density $r + 1$, which contradicts the assumption that $\text{pr}_C(w)$ is in the same two-sided cell of w (cf. [6, Corollary 11.5]). Thus the result follows. \square

EXAMPLE 5.20. Suppose that $w = [6, 1, 18, 3, 19, 24, 12, 15, 17, 10]$ as in Example 5.13 and 5.7. We already calculated $\text{pr}_{C_1}^{-1}(w)$, $\text{pr}_{C_2}(w)$, $\text{pr}_{C_3}(w)$, and $\text{pr}_{C_3}^{-1}(w)$, and their images under Φ . Let us calculate the remaining possibilities.

$$\begin{aligned} \text{pr}_{C_1}(w) &= [6, 3, 18, 10, 19, 24, 12, 15, 17, 11] \\ &\xrightarrow{\Phi} (((1, 3, 6, 10), (2, 5, 9), (4, 7), (8)), ((3, 5, 6, 9), (4, 7, 8), (1, 10), (2)), (3, 4, 1, 0)), \\ \text{pr}_{C_2}^{-1}(w) &= [6, 1, 18, 3, 19, 24, 7, 12, 15, 10] \\ &\xrightarrow{\Phi} (((1, 3, 7, 10), (2, 5, 6), (4, 8, 9)), ((1, 3, 5, 6), (7, 8, 9), (2, 4, 10)), (1, 3, 2)), \\ \text{pr}_{C_2'}(w) &= [12, 1, 18, 3, 19, 24, 15, 16, 17, 10] \\ &\xrightarrow{\Phi} (((1, 3, 7, 10), (2, 5, 6), (4, 8, 9)), ((1, 3, 5, 6), (7, 8, 9), (2, 4, 10)), (3, 3, 2)), \\ \text{pr}_{C_2'}^{-1}(w) &= [5, 1, 18, 3, 19, 24, 6, 12, 17, 10] \\ &\xrightarrow{\Phi} (((1, 3, 6, 10), (2, 5, 9), (4, 7), (8)), ((3, 5, 6, 9), (4, 7, 8), (1, 10), (2)), (1, 4, 1, 0)). \end{aligned}$$

Note that $w \in \underline{c}_{(3,3,3,1)}$ but these elements are not contained in this two-sided cell.

We are ready to prove Lemma 5.15. To recap, we assume that to an affine permutation w one can apply two operations: (1) a right star operation centered at p for $p \in [1, n]$; (2) a proper PR/IPR applied to the q -th southwest channel C in \mathfrak{C}_w . Let us denote by w^* the outcome of the right star operation and by \tilde{w} the outcome of such a proper PR/IPR. First we claim the following.

LEMMA 5.21. *Suppose that there exists $\tilde{w}^* \in \widetilde{\mathcal{S}}_n$ such that $\tilde{w} \mapsto \tilde{w}^*$ is again the right star operation centered at p and $w^* \mapsto \tilde{w}^*$ is a PR/IPR. Then the latter operation is proper and applied to the q -th southwest channel of w^* .*

Proof. By Theorem 5.12, $\Phi(w)$ and $\Phi(\tilde{w})$ differ by a single value of $\frac{\#}{\rho}$ on the top block. Since $w, \tilde{w}, w^*, \tilde{w}^*$ all lie within the same two-sided cell by assumption, Lemma 5.19 tells us that the PR/IPR $w^* \mapsto \tilde{w}^*$ should be proper. Since the distances between channels of w^* (resp. \tilde{w}^*) are the same as those of w (resp. \tilde{w}) by Lemma 5.18, we see that the PR/IPR $w^* \mapsto \tilde{w}^*$ should be applied to the q -th channel of w^* . \square

Thus, it is enough to show that the commutative diagram can be completed by a PR/IPR (without checking properness) applied to a channel of w^* (without checking that it is the q -th southwest channel). Meanwhile, without loss of generality we may assume that $p \in [2, n - 1]$; if $p \in \{1, n\}$, then we may multiply ω on the right so that the right star operation is centered in $[2, n - 1]$; this process also respects proper PR/IPRs. (See the comment after Lemma 5.15. Also, here $n \geq 3$ since the right star operation is not well-defined for $n \leq 2$.) Moreover, it suffices only to consider partial rotations because the inverse case can be deduced by considering w^{-1} .

We write $\tilde{w} = \text{pr}_C(w)$ where $C \subset w$ is a northeast channel of a river. We set $a, b, c \in \mathbb{Z}$ to the y -coordinates of some elements in C such that $w^{-1}(a) < w^{-1}(b) < w^{-1}(c)$. (Recall that the y -axis directs to the east in our convention.) We assume that no other elements in C are between them, i.e. if $(y, w(y)) \in C$ and $a < w(y) < c$ then $b = w(y)$. (If the density of C is 2 (resp. 1) then we have $c = a + n$ (resp. $c = b + n = a + 2n$.)

We argue case-by-case based on the size of $[p - 1, p + 1] \cap X$ where X is a subset of $[1, n]$ such that $X + n\mathbb{Z}$ is the domain of C . Here we will verify that (1) the right star operation centered at p is well-defined for \tilde{w} , and (2) there exists a channel $C' \subset w^*$ such that its corresponding partial rotation coincide with \tilde{w}^* .

Case 0. Suppose that $\#[p - 1, p + 1] \cap X = 0$. In this case the two operations simply commute, and the commutative diagram can be trivially completed.

Case 1. Suppose that $\#[p - 1, p + 1] \cap X = 1$.

- (1) The right star operation looks like (here either side can be w):

$$\dots xby \dots c \dots \rightsquigarrow \dots bxy \dots c \dots .$$

- (a) Suppose that $x < y < b$. Since $x < y < c$, after the partial rotation we get a valid right star operation

$$\dots b \dots xcy \dots \rightsquigarrow \dots b \dots cxy \dots ,$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (b) Suppose that $b < y < x$. If $c > y$ then y and its translates can be inserted to C to yield a longer stream in w , which is a contradiction. Thus $c < y < x$ and after the partial rotation we get a valid right star operation

$$\dots b \dots xcy \dots \rightsquigarrow \dots b \dots cxy \dots ,$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (2) The right star operation looks like (here either side can be w):

$$\dots yxb \dots c \dots \rightsquigarrow \dots ybx \dots c \dots .$$

- (a) Suppose that $x < y < b$. Since $x < y < c$, after the partial rotation we get a valid right star operation

$$\dots b \dots yxc \dots \rightsquigarrow \dots b \dots ycx \dots ,$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (b) Suppose that $b < y < x$. If $c > y$ then y and its translates can be inserted to $C \langle 1 \rangle$ to yield a longer stream in \tilde{w} , which is a contradiction. Thus $c < y < x$ and after the partial rotation we get a valid right star operation

$$\dots b \dots yxc \dots \rightsquigarrow \dots b \dots ycx \dots ,$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (3) The right star operation looks like (here either side can be w):

$$\dots yxb \dots c \dots \rightsquigarrow \dots xyb \dots c \dots .$$

Without loss of generality we may assume that $x < y$, i.e. $x < b < y$. If $c > y$ then y and its translates can be inserted to $C\langle 1 \rangle$ to yield a longer stream in \tilde{w} , which is a contradiction. Thus $x < c < y$ and after the partial rotation we get a valid right star operation

$$\dots b \dots yxc \dots \rightsquigarrow \dots b \dots xyc \dots ,$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (4) The right star operation looks like (here either side can be w):

$$\dots byx \dots c \dots \rightsquigarrow \dots bxy \dots c \dots .$$

Without loss of generality we may assume that $x < y$, i.e. $x < b < y$. If $c > y$ then y and its translates can be inserted to C to yield a longer stream in w , which is a contradiction. Thus $x < c < y$ and after the partial rotation we get a valid right star operation

$$\dots b \dots cyx \dots \rightsquigarrow \dots b \dots cxy \dots ,$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

Case 2. Suppose that $\#([p - 1, p + 1] \cap X) = 2$.

- (1) The right star operation looks like:

$$w = \dots azb \dots c \dots \rightsquigarrow w^* = \dots zab \dots c \dots .$$

Here we should have $a < b < z$. If $z < c$, then z and its translates can be inserted to $C\langle 1 \rangle$ to yield a longer stream in \tilde{w} , which is a contradiction. Thus $b < c < z$ and after the partial rotation we get a valid right star operation

$$\tilde{w} = \dots a \dots bzc \dots \rightsquigarrow \tilde{w}^* = \dots a \dots zbc \dots$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (2) The right star operation looks like:

$$w = \dots zab \dots c \dots \rightsquigarrow w^* = \dots azb \dots c \dots .$$

Here we should have $a < b < z$.

- (a) If $c < z$, then after the partial rotation we get a valid right star operation

$$\tilde{w} = \dots a \dots zbc \dots \rightsquigarrow \tilde{w}^* = \dots a \dots bzc \dots$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (b) If $c > z$, then after the partial rotation we get a valid right star operation

$$\tilde{w} = \dots a \dots zbc \dots \rightsquigarrow \tilde{w}^* = \dots a \dots zcb \dots$$

and a valid channel $C' = \dots, a, z, c, \dots \subset w^*$ (i.e. replacing b with z in C) so that the commutative diagram can be completed.

- (3) The right star operation looks like (here either side can be w):

$$\cdots axb \cdots c \cdots \rightsquigarrow \cdots abx \cdots c \cdots .$$

Here we should have $x < a < b < c$. After the partial rotation we get a valid right star operation

$$\cdots a \cdots bxc \cdots \rightsquigarrow \cdots a \cdots bcx \cdots$$

and a valid channel $C' \subset w^*$ whose y -coordinates are identical to C so that the commutative diagram can be completed.

- (4) The right star operation looks like:

$$w = \cdots aby \cdots c \cdots \rightsquigarrow w^* = \cdots bay \cdots c \cdots .$$

Here we should have $a < y < b < c$. After the partial rotation we get a valid right star operation

$$\tilde{w} = \cdots a \cdots bcy \cdots \rightsquigarrow \tilde{w}^* = \cdots a \cdots byc \cdots$$

and a valid channel $C' = \dots, a, y, c, \dots \subset w^*$ (i.e. replacing b with y in C) so that the commutative diagram can be completed.

- (5) The right star operation looks like:

$$w = \cdots yab \cdots c \cdots \rightsquigarrow w^* = \cdots yba \cdots c \cdots .$$

Here we should have $a < y < b < c$. However, it is impossible since y and its translates can be inserted to $C\langle 1 \rangle$ to yield a longer stream in \tilde{w} .

Case 3. Suppose that $\#([p - 1, p + 1] \cap X) = 3$. However, this is impossible since it means that $w(p - 1) < w(p) < w(p + 1)$ and the right star operation cannot be applied.

We exhausted all the possibilities and this finishes the proof of the Diamond Lemma.

6. PROOF OF BLASIAK’S CONJECTURE

6.1. SAME LEFT CELL IMPLIES SAME Ω . We are ready to prove half of Theorem 4.5, i.e. $\{w \in \underline{c}_\lambda \mid \text{sgn}_\Omega(w) = \Omega\}$ is a union of left cells. To this end the following lemma is essential.

LEMMA 6.1. For $v, w \in \widetilde{\mathcal{S}}_n$, let us write $v(i, j) = in + v(j)$ and $w(i, j) = in + w(j)$ where $i, j \in \mathbb{Z}$. For $X \subset [1, n]$, suppose that

- (a) the relative orders of $\{v(i, x)\}_{i \in [0, n-1], x \in X}$ and $\{w(i, x)\}_{i \in [0, n-1], x \in X}$ are the same, i.e. for any $i, i' \in [0, n - 1], j, j' \in X$ we have $v(i, j) \leq v(i', j')$ if and only if $w(i, j) \leq w(i', j')$.
- (b) $v(y) + n^2 < v(x)$ and $w(y) + n^2 < w(x)$ for $x \in X$ and $y \in [1, n] - X$, and
- (c) $\text{sgn}_\Omega(v_{[1, n] - X}) = \text{sgn}_\Omega(w_{[1, n] - X})$.

Then $\text{sgn}_\Omega(v) = \text{sgn}_\Omega(w)$.

Proof. First suppose that $X = [1, n]$, so that the conditions (b) and (c) are vacuous. In this case, the statement directly follows from the definition of the sign insertion algorithm. More precisely, first note that the entries which appear in one of \mathfrak{P}_i or v_i (resp. w_i) in the sign insertion algorithm are contained in $\{v(i, j)\}_{i \in [0, n-1], j \in [1, n]}$ (resp. $\{w(i, j)\}_{i \in [0, n-1], j \in [1, n]}$) since an entry cannot be bumped more than n times during the process. Thus, one can show by induction on the steps of sign insertion that indeed Ω_i are the same for v and w , using the assumption that the relative orders of $\{v(i, j)\}_{i \in [0, n-1], j \in [1, n]}$ and $\{w(i, j)\}_{i \in [0, n-1], j \in [1, n]}$ are the same.

In general, condition (b) implies that $v(i, y) < v(i', x)$ and $w(i, y) < w(i', x)$ for any $i, i' \in [0, n - 1], x \in X$, and $y \in [1, n] - X$. In other words, $v(i', x)$ and $w(i', x)$

for $i' \in [0, n - 1]$ and $x \in X$ behave as if they are very large compared to $v(i, y)$ and $w(i, y)$ for $i \in [0, n - 1]$ and $y \in [1, n] - X$. Since condition (c) gives the equality of two Ω 's when the sign insertion algorithm is applied to the “smaller part” of v and w , respectively, similarly to above one can use induction on the steps of sign insertion to conclude the claim. \square

Let us prove that $\text{sgn}_\Omega(v) = \text{sgn}_\Omega(w)$ when v and w are in the same left cell indexed by $Q \in \text{RSYT}(\lambda)$. Let m be the multiplicity of λ_1 in λ and let $\mathfrak{C}_w = (C_1, \dots, C_m)$ be a sequence in Definition 5.5. Let $X_i = \{x \in [1, n] \mid (x, w(x)) \in C_i\}$. If we let $X_i = \{a_1, \dots, a_k\}$ where $a_1 < \dots < a_k$ then by definition of the channel we have $w(a_1) < \dots < w(a_k) < w(a_1) + n$. Moreover, since proper PR/IPR do not affect $\text{sgn}_\Omega(w)$ and the left cell of w , by applying them repeatedly we may assume that $0 \ll \rho_1 \ll \dots \ll \rho_m$, which results in $w(y) \ll w(x_1) \ll \dots \ll w(x_m)$ for any $y \in [1, n] - \bigsqcup_{i=1}^m X_i, x_1 \in X_1, \dots, x_m \in X_m$.

Let us recall the description of AMBC in terms of asymptotic realization, i.e. [6, Section 7]. We observe the following statement.

LEMMA 6.2. *For $w \in \widetilde{\mathcal{S}}_n$ and $X \subset [1, n]$, suppose that $\{(x, w(x)) \mid x \in X + n\mathbb{Z}\}$ is a channel of w . Also assume that $w(y) \ll w(x)$ for any $x \in X$ and $y \in [1, n] - X$. Then $\Phi(w) = (P, Q, \#p)$ for some P and $\#p$ if and only if $Q_1 = X$ (as sets) and $\Phi(w|_{[1, n] - X}) = (P', Q_{\geq 2}, \#p')$ for some P' and $\#p'$.*

Proof. Here we prove the only if part, but its converse is proved similarly. Note that Q is equal to the recording tableau modulo n where the usual Robinson–Schensted algorithm is applied to an infinite sequence $(w(1), w(2), \dots)$. (This is implicit in [6, Section 7] and shown in the proof of [10, Theorem 10.3].) The condition implies that $w(in + x) > w(y)$ for any $i \in \mathbb{Z}, x \in X$, and $y \in [1, in + x - 1]$. Thus when $w(in + x)$ is inserted to the infinite insertion tableau it is always located at the end of the first row. Thus it follows that the first row of Q should contain all the elements in X . However, since $\#X$ is equal to the length of the first row of Q_1 because $\{(x, w(x)) \mid x \in X + n\mathbb{Z}\}$ is a channel, it follows that $Q_1 = X$.

On the other hand, the bumping process of $w(in + y)$ for $i \in \mathbb{N}$ and $y \in [1, n] - X$ is almost the same as considering the partial permutation $w|_{[1, n] - X}$, except that at the end of each step it bumps some element of the form $w(jn + x)$ for $j \in \mathbb{N}$ and $x \in X$. It means that the recording tableau of the infinite word corresponding to $w|_{[1, n] - X}$ is the same as that corresponding to w after the first row is removed. Thus by taking residues modulo n we obtain the second claim. \square

In our situation, by iterating the above lemma we see that $X_i = Q_{m+1-i}$ for $i \in [1, m]$. The most important part of this observation is that X_1, \dots, X_m only depend on Q whenever $0 \ll \rho_1 \ll \dots \ll \rho_m$. More precisely, if we set $\#p$ such that $\rho_i + N < \rho_{i+1}$ for $i \in [1, m - 1]$ and $\rho_1 > N$ for a sufficiently large N , then the set of x -coordinates of each disjoint channel C_1, \dots, C_m stabilizes as N grows. Moreover, if we set $X = \bigsqcup_{i=1}^m X_i$, then $w(y) + n^2 < w(x)$ for $x \in X$ and $y \in [1, n] - X$, and $[w(i, j) \leq w(i', j')] \text{ for } i, i' \in [0, n - 1] \text{ and } j, j' \in X]$ if and only if either $[j, j' \in X_k \text{ and } j \leq j' \text{ for some } k \in [1, m]]$ or $[j \in X_k \text{ and } j' \in X_{k'} \text{ for some } 1 \leq k < k' \leq m]$.

Now assume that both $\#p$ and ϕ satisfy $0 \ll \rho_1 \ll \dots \ll \rho_m$ and $0 \ll \phi_1 \ll \dots \ll \phi_m$, which is again possible. By the argument above and Lemma 6.1, in order to prove $\text{sgn}_\Omega(v) = \text{sgn}_\Omega(w)$ it suffices to show $\text{sgn}_\Omega(v|_{[1, n] - X}) = \text{sgn}_\Omega(w|_{[1, n] - X})$ where $X = \bigsqcup_{i=1}^m X_i = \bigsqcup_{i=1}^m Q_i$. However, Lemma 6.2 implies that the partial permutations $v|_{[1, n] - X}$ and $w|_{[1, n] - X}$ have the same Q tableau under AMBC. Therefore, the result follows from induction on n .

6.2. SAME Ω AND SAME TWO-SIDED CELL IMPLIES SAME LEFT CELL. Let us continue with the other half of Theorem 4.5, i.e. here we prove that $\text{sgn}_\Omega(w) \neq \text{sgn}_\Omega(v)$ if w and v are in the same two-sided cell but in different left cells. Suppose that a partition $\lambda \vdash n$ of length $l = l(\lambda)$ and a tableau $T \in \text{RSYT}(\lambda)$ is given. Let us write $L_i = \sum_{j=i+1}^l \lambda_j$ for $i \in [0, l]$ so that $L_{i-1} - L_i = \lambda_i$. Recall that we set $A_X = (A_{x_1}, \dots, A_{x_s})$ for a sequence $A = (A_1, \dots, A_r)$ and a set $X = \{x_1, \dots, x_s\} \subset [1, r]$ such that $x_1 < x_2 < \dots < x_s$. For $N \in \mathbb{Z}$, we consider an element $w_{T,N} \in \widetilde{\mathcal{S}}_n$ where

$$(w_{T,N})_{T_i} = (Nn(l-i) + L_i + 1, Nn(l-i) + L_i + 2, \dots, Nn(l-i) + L_{i-1}).$$

Usually N will be a sufficiently large natural number.

When $N \gg n > 0$, it is not hard to show that $(\text{sgn}_{\mathfrak{P}}(w_{T,N}))_{[L_i+1, L_{i-1}]}$ is obtained from $(w_{T,N})_{T_i}$ by applying partial rotations, i.e. there exists $m_i \in [0, n-1]$ such that $(\text{sgn}_{\mathfrak{P}}(w_{T,N}))_{[L_i+1, L_{i-1}]} = (\text{pr}_{T_i}^{m_i}(w_{T,N}))_{T_i}$. (This can be proved using induction on the steps of sign insertion.) On the other hand, let $C_i = \{(jn+x, jn+w_{T,N}(x)) \mid x \in T_i, j \in \mathbb{Z}\}$ be a stream of $w_{T,N}$. Similarly, we regard $\text{sgn}_{\mathfrak{P}}(w_{T,N})$ as an affine permutation and set $C'_i = \{(jn+x, jn+\text{sgn}_{\mathfrak{P}}(w_{T,N})(x)) \mid x \in [L_i+1, L_{i-1}], j \in \mathbb{Z}\}$ which becomes a stream of $\text{sgn}_{\mathfrak{P}}(w_{T,N})$. Then it is easy to show that when $0 < k \ll N$ we have $\text{pr}_{C'_i}^k(\text{sgn}_{\mathfrak{P}}(w_{T,N})) = \text{sgn}_{\mathfrak{P}}(\text{pr}_{C_i}^k(w_{T,N}))$, i.e. the partial rotation “commutes with” $\text{sgn}_{\mathfrak{P}}$. (This follows from a similar argument to the proof of Lemma 6.1 when $X = [1, n]$.) Therefore, if we set $k_i = n\lambda_i - m_i > 0$ for $i \in [1, l]$ and define $\tilde{w}_{T,N} = (\text{pr}_{C_1}^{k_1} \circ \dots \circ \text{pr}_{C_l}^{k_l})(w_{T,N})$ then it satisfies

$$\begin{aligned} (\text{sgn}_{\mathfrak{P}}(\tilde{w}_{T,N}))_{[L_i+1, L_{i-1}]} &= (Nn(l-i) + n^2 + L_{i-1} + 1, \dots, Nn(l-i) + n^2 + L_i) \\ &= (w_{T,N})_{T_i} + (n^2, n^2, \dots, n^2). \end{aligned}$$

(Note that $\text{sgn}_{\mathfrak{P}}(\tilde{w}_{T,N})$ only depends on λ and N but not T .)

In general $\text{pr}_{C_i}^{k_i}$ are not proper because C_i 's do not need to be channels. However, as $N \gg 0$, using inductive argument based on Lemma 6.2, one can easily show in this case that $\tilde{w}_{T,N}$ is contained in the left cell parametrized by T , i.e. $\Phi(\tilde{w}_{T,N}) = (P, T, \overset{\#}{p})$ for some P and $\overset{\#}{p}$.

EXAMPLE 6.3. Let $T = ((3, 6, 7, 9), (4, 8, 10), (1, 5), (2)) \in \text{RSYT}(4, 3, 2, 1)$. Then we have

$$w_{T,10} = [102, 1, 307, 204, 103, 308, 309, 205, 310, 206].$$

(Here, $N = 10$ is sufficiently large for our purpose.) Direct calculation shows that

$$\text{sgn}_{\mathfrak{P}}(w_{T,10}) = (1, 103, 112, 206, 214, 215, 318, 319, 320, 327).$$

Here $m_1 = 5, m_2 = 2, m_3 = 1, m_4 = 0$. Thus if we set $\tilde{w}_{T,10} = (\text{pr}_{T_1}^{35} \circ \text{pr}_{T_2}^{28} \circ \text{pr}_{T_3}^{19} \circ \text{pr}_{T_4}^{10})(w_{T,10}) = [193, 101, 390, 295, 202, 397, 398, 296, 399, 304]$ then we have

$$\text{sgn}_{\mathfrak{P}}(\tilde{w}_{T,10}) = [101, 202, 203, 304, 305, 306, 407, 408, 409, 410]$$

as desired. Furthermore, we have

$$\Phi(\tilde{w}_{T,10}) = \left(\begin{array}{c|c} \overline{1 \ 3 \ 6 \ 10} & \overline{3 \ 6 \ 7 \ 9} \\ \hline \overline{2 \ 5 \ 9} & \overline{4 \ 8 \ 10} \\ \hline \overline{4 \ 8} & \overline{1 \ 5} \\ \hline \overline{7} & \overline{2} \end{array}, \begin{pmatrix} 155 \\ 88 \\ 39 \\ 10 \end{pmatrix} \right),$$

i.e. $\tilde{w}_{T,10}$ is contained in the left cell parametrized by T .

We are ready to prove the other half of Theorem 4.5. For $w, v \in \widetilde{\mathcal{S}}_n$, suppose that $w \in \Gamma_T$ and $v \in \Gamma_S$ where $S, T \in \text{RSYT}(\lambda)$ for some $\lambda \vdash n$ but $S \neq T$. Here we prove that $\text{sgn}_\Omega(w) \neq \text{sgn}_\Omega(v)$. Since we already showed that the elements in the

same left cell have the same image under sgn_Ω , it suffices to assume that $w = \tilde{w}_{T,N}$ and $v = \tilde{w}_{S,N}$ for sufficiently large N .

By the argument above, we have $\text{sgn}_{\mathfrak{P}}(\tilde{w}_{T,N}) = \text{sgn}_{\mathfrak{P}}(\tilde{w}_{S,N})$, i.e. $\text{sgn}_{\mathfrak{P}}(\tilde{w}_{T,N})$ does not depend on T but only on λ once $N \gg 0$ is fixed. However, the map $w \mapsto (\text{sgn}_{\mathfrak{P}}(w), \text{sgn}_\Omega(w))$ is injective by the argument in [3, p.2333]. It follows that $\text{sgn}_\Omega(\tilde{w}_{T,N}) = \text{sgn}_\Omega(\tilde{w}_{S,N})$ if and only if $\tilde{w}_{T,N} = \tilde{w}_{S,N}$ if and only if $T = S$. Therefore, we have $\text{sgn}_\Omega(\tilde{w}_{T,N}) \neq \text{sgn}_\Omega(\tilde{w}_{S,N})$ as desired.

6.3. PARTIAL ROTATIONS AND LEFT CELLS. As a by-product of our argument, we may connect the elements in the intersection of a right cell and a left cell by (inverse) partial rotation. To this end first we observe the following lemma.

LEMMA 6.4. *For a partial permutation w and $X \subset [1, n]$, suppose that w_X is a channel of w and $w(y) \ll w(x)$ for $x \in X$ and $y \in [1, n] - X$. Let $X' = \{n + 1 - x \mid x \in \text{im } w_X \cap [1, n]\}$. If we let $\Phi(w) = (P, Q, -)$ and $\Phi(w_{[1, n] - X}) = (P', Q', -)$, then we have $P = \text{evac}((X') \frown \text{evac}(P'))$ and $Q = (X) \frown Q'$ where evac is the affine evacuation defined in [4].*

Proof. By Lemma 6.2 we have $X = Q_1$. Moreover, if we consider $\mathcal{R}(w^{-1})$ and X' instead then the conditions of Lemma 6.2 is still valid and X' is equal to the first row of $\text{evac}(P)$ because $\Phi(\mathcal{R}(w^{-1})) = (\text{evac}(Q), \text{evac}(P), -)$. Since $\mathcal{R}(w^{-1})_{[1, n] - X'} = \mathcal{R}((w_{[1, n] - X})^{-1})$, it follows that $\text{evac}(P) = (X') \frown \text{evac}(P')$, i.e. $P = \text{evac}((X') \frown \text{evac}(P'))$ as desired. \square

The main statement of this section is as follows.

THEOREM 6.5. *Suppose that $P, Q \in \text{RSYT}(\lambda)$ and $w, \tilde{w} \in (\Gamma_P)^{-1} \cap \Gamma_Q$. Then there exists a sequence $w = w_0, w_1, \dots, w_k = \tilde{w}$, which all lie in $(\Gamma_P)^{-1} \cap \Gamma_Q$, such that for $i \in [1, k]$ either $w_i = \text{pr}_{S_{i-1}}(w_{i-1})$ or $w_i = \text{pr}_{S_{i-1}}^{-1}(w_{i-1})$ where S_{i-1} is a stream (not necessarily a channel) of w_{i-1} .*

Proof. Let $\Phi(w) = (P, Q, \#p)$ and $\Phi(\tilde{w}) = (P, Q, \#\varphi)$ for some $\#p = (\rho_1, \rho_2, \dots)$ and $\#\varphi = (\varphi_1, \varphi_2, \dots)$. By applying the proper partial rotations to the northeast channels of w and \tilde{w} , respectively, we may assume that $\rho_1, \varphi_1 \gg 0$. Then by Lemma 6.2, we see that w_{Q_1} and \tilde{w}_{Q_1} are channels of w and \tilde{w} , respectively. Moreover, Lemma 6.4 implies that $\text{im } w_{Q_1} = \text{im } \tilde{w}_{Q_1} = \{n + 1 - x \mid x \in \text{evac}(P)_1\} + n\mathbb{Z}$. Therefore, by applying proper partial rotation we may assume that $w_{Q_1} = \tilde{w}_{Q_1}$ while $w(y) \ll w(x)$ and $\tilde{w}(y) \ll \tilde{w}(x)$ for $x \in X$ and $y \in [1, n] - X$. Now again by Lemma 6.4, it suffices to show that $w_{[1, n] - X}$ and $\tilde{w}_{[1, n] - X}$ are connected by a similar sequence described in the statement. Thus the claim follows from induction on the length of P and Q . \square

7. SIGN INSERTION AND LASCOUX–SCHÜTZENBERGER STANDARDIZATION

Let us set $\Upsilon_\Omega = \{T \in \text{RSYT}(n) \mid \text{sgn}_\Omega(w) = \Omega \text{ for some } w \in \Gamma_T\}$. Then Theorem 4.5 implies that $\Upsilon_\Omega \cap \text{RSYT}(\lambda)$ is a singleton if nonempty. Our goal in this section is to provide a relation between elements in Υ_Ω , necessarily in different shapes.

7.1. TABLEAUX, CRYSTAL OPERATORS, AND ROBINSON–SCHENSTED ALGORITHM.

For $\alpha \models n$ and a row-standard Young tabloid $T \in \text{RSYT}(\alpha)$ we consider its associated two-row array $\mathcal{A}(T)$ whose first row records $(\text{length of } \alpha) + 1 - (\text{row number})$ and whose second row is $\text{rw}(T)$. For example, if $T = ((2), \emptyset, (3, 5, 6), (1, 4))$ then $\mathcal{A}(T) = \begin{pmatrix} 1 & 1 & 2 & 2 & 2 & 4 \\ 1 & 4 & 3 & 5 & 6 & 2 \end{pmatrix}$. Then we may apply the usual Robinson–Schensted–Knuth algorithm (see [17, Chapter 7] for more detail) to $\mathcal{A}(T)$. Let us denote this map by $\text{RSK} : \text{RSYT}(\alpha) \rightarrow \bigsqcup_{\lambda \vdash n} \text{SYT}(\lambda) \times \text{SSYT}(\lambda, \alpha^{\text{rev}})$; it sends $T \mapsto (\text{RSK}_P(T), \text{RSK}_Q(T))$. This is indeed a bijection.

Let \mathcal{S}_∞ be the set of permutations of $\mathbb{Z}_{>0}$ of finite support. (In other words, $\mathcal{S}_\infty = \bigcup_{n \geq 1} \mathcal{S}_n$.) Let $s_i \in \mathcal{S}_\infty$ be the transposition swapping i and $i + 1$. For a composition $\alpha \models n$, we may regard it as an infinite sequence all of whose entries but finite are zero, say $\alpha = (\alpha_1, \alpha_2, \dots)$. Then $s_i \cdot \alpha = (\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_{i+1}, \alpha_i, \alpha_{i+2}, \dots)$ is well-defined and there exists a bijection $s_i : \text{SSYT}(n, \alpha) \rightarrow \text{SSYT}(n, s_i \cdot \alpha)$ such that $\text{rw}(T) \mapsto \text{rw}(s_i(T))$ is a crystal reflection operator defined by Lascoux and Schützenberger [12] (see also [16, 3.2]). These operations are involutions and satisfy the braid relation, and thus these yield an action of \mathcal{S}_∞ on $\bigsqcup_{\alpha \models n} \text{SSYT}(n, \alpha)$.

Moreover, there exists a bijection $R_i : \text{RSYT}(\alpha) \rightarrow \text{RSYT}(s_i \cdot \alpha)$, called a combinatorial R -matrix, which comes from an isomorphism of tensor products of one-row Kirillov–Reshetikhin crystals. (See [15, 4.8] or [4, Definition 3.11].) These operations are also involutions and satisfy the braid relation, and thus these yield an action of \mathcal{S}_∞ on $\bigsqcup_{\alpha \models n} \text{RSYT}(\alpha)$, denoted by R_σ for $\sigma \in \mathcal{S}_\infty$. Then by [15, Proposition 5.1] (see also [4, Proposition 3.21] and the footnote thereafter), we have $\text{RSK}_P(R_i(T)) = \text{RSK}_P(T)$ and $\text{RSK}_Q(R_i(T)) = s_{i(T)-i} \text{RSK}_Q(T)$. Therefore, if $\sigma \in \mathcal{S}_{l(T)}$ then $\text{RSK}_P(R_\sigma(T)) = \text{RSK}_P(T)$ and $\text{RSK}_Q(R_\sigma(T)) = (w_0 \sigma w_0) \text{RSK}_Q(T)$ where w_0 is the longest element of $\mathcal{S}_{l(T)}$.

7.2. LASCoux–SCHÜTZENBERGER STANDARDIZATION. We recall the standardization map of Lascoux and Schützenberger. Here we mainly follow the argument and notations from [16]. Suppose that we are given two compositions $\alpha, \beta \models n$ whose rearrangements are partitions $\alpha^+, \beta^+ \vdash n$, respectively. Assume that $\alpha^+ \geq \beta^+$ with respect to dominance order. Then there exists an injective map $\theta_\alpha^\beta : \text{SSYT}(n, \alpha) \rightarrow \text{SSYT}(n, \beta)$ defined as follows.

- If $\alpha^+ = \beta^+$ then choose any permutation $\sigma \in \mathcal{S}_\infty$ so that $\sigma \cdot \alpha = \beta$ and let $\theta_\alpha^\beta = \sigma$.
- If $\beta_i = \alpha_i$ for $i \geq 3$ and $\beta_1 + 1 = \alpha_1 > \beta_2 = \alpha_2 + 1$, then θ_α^β acts on a tableau by changing the rightmost letter 1 to 2 (which is always possible).
- In general, there exists a sequence $\alpha = \alpha_0, \alpha_1, \dots, \alpha_k = \beta$ where each $\alpha_i \rightarrow \alpha_{i+1}$ is either in the first or the second case. Then we define $\theta_\alpha^\beta = \theta_{\alpha_{k-1}}^{\alpha_k} \circ \dots \circ \theta_{\alpha_0}^{\alpha_1}$.

Then by [11] this map is well-defined (i.e. it does not depend on the choice of $\alpha = \alpha_0, \alpha_1, \dots, \alpha_k = \beta$ on the last part) and also satisfies $\theta_\beta^\gamma \circ \theta_\alpha^\beta = \theta_\alpha^\gamma$.

For $\alpha \models n$, note that $\text{SSYT}(n, \alpha)$ is endowed with a graded poset structure by cyclage where the grading is given by cocharge. By [12] and [11], the map $\theta_\alpha^\beta : \text{SSYT}(n, \alpha) \rightarrow \text{SSYT}(n, \beta)$ is indeed a grade-preserving poset embedding (see also [16, Theorem 44, 45]).

7.3. INVESTIGATION OF Υ_Ω . The main result of this section is the following theorem.

THEOREM 7.1. *Suppose that $\Upsilon_\Omega = \{T \in \text{RSYT}(n) \mid \text{sgn}_\Omega(w) = \Omega \text{ for some } w \in \Gamma_T\}$ is nonempty. Then,*

- $\text{RSK}_P(\Upsilon_\Omega)$ consists of a single element.
- If $S, T \in \Upsilon_\Omega$ and $\text{sh}(S) \geq \text{sh}(T)$ then $\theta_{\text{sh}(S)^{\text{rev}}}^{\text{sh}(T)^{\text{rev}}}(\text{RSK}_Q(S)) = \text{RSK}_Q(T)$. Moreover, if $T \in \Upsilon_\Omega$ and $\text{sh}(T) \geq \lambda$ for some partition $\lambda \vdash n$ then $\theta_{\text{sh}(T)^{\text{rev}}}^{\lambda^{\text{rev}}}(\text{RSK}_Q(T)) \in \text{RSK}_Q(\Upsilon_\Omega)$. In other words, $\text{RSK}_Q(\Upsilon_\Omega)$ is “stable under θ ”.

As a corollary we obtain the following.

COROLLARY 7.2. *For each $n \in \mathbb{Z}_{>0}$, $\{\text{sgn}_\Omega(w) \mid w \in \widetilde{\mathcal{S}}_n\}$ is canonically bijective with $\text{RSYT}(1^n)$.*

Proof. If we define $\text{RSYT}(1^n) \rightarrow \{\text{sgn}_\Omega(w) \mid w \in \widetilde{\mathcal{S}}_n\} : T \mapsto \text{sgn}_\Omega(v_T)$ where v_T is any element in Γ_T , then this map is well-defined and injective by Theorem 4.5. We claim that this is also surjective. Indeed, suppose that $\Upsilon_\Omega \neq \emptyset$ and $T \in \Upsilon_\Omega$. Then Theorem 7.1 implies that $\text{RSK}_P(\Upsilon_\Omega) = \{\text{RSK}_P(T)\}$ and $\text{RSK}_Q(\Upsilon_\Omega) \ni \theta_{\text{sh}(T)}^{(1^n)\text{rev}}(\text{RSK}_Q(T))$. Now if we let $S \in \text{RSYT}(1^n)$ be the preimage of $(\text{RSK}_P(T), \theta_{\text{sh}(T)}^{(1^n)\text{rev}}(\text{RSK}_Q(T)))$ under RSK then we see that $\Upsilon_\Omega \ni S$ as desired. \square

EXAMPLE 7.3. When $n = 4$, the sets Υ_Ω are as follows.

$$\begin{aligned} \Upsilon_{(1,2,3,4)} &= (((1, 2, 3, 4)), ((2, 3, 4), (1)), ((3, 4), (1, 2)), ((3, 4), (2), (1)), ((4), (3), (2), (1))) \\ \Upsilon_{(1,2,3,5)} &= (((1, 2, 3), (4)), ((2, 3), (1, 4)), ((2, 3), (4), (1)), ((3), (4), (2), (1))) \\ \Upsilon_{(1,2,4,5)} &= (((1, 2, 4), (3)), ((2, 4), (1, 3)), ((2, 4), (3), (1)), ((2), (4), (3), (1))) \\ \Upsilon_{(1,3,4,5)} &= (((1, 3, 4), (2)), ((1, 4), (2, 3)), ((1, 4), (3), (2)), ((1), (4), (3), (2))) \\ \Upsilon_{(1,3,5,6)} &= (((1, 3), (2, 4)), ((1, 3), (4), (2)), ((3), (1), (4), (2))) \\ \Upsilon_{(1,2,5,6)} &= (((1, 2), (3, 4)), ((1, 2), (4), (3)), ((2), (1), (4), (3))) \\ \Upsilon_{(1,4,5,7)} &= (((1, 4), (2), (3)), ((1), (2), (4), (3))) \\ \Upsilon_{(1,3,5,7)} &= (((1, 3), (2), (4)), ((1), (3), (4), (2))) \\ \Upsilon_{(1,2,5,7)} &= (((1, 2), (3), (4)), ((2), (3), (4), (1))) \\ \Upsilon_{(1,2,3,6)} &= (((2, 3), (1), (4)), ((3), (2), (4), (1))) \\ \Upsilon_{(1,2,4,6)} &= (((2, 4), (1), (3)), ((4), (2), (3), (1))) \\ \Upsilon_{(1,3,4,6)} &= (((3, 4), (1), (2)), ((4), (1), (3), (2))) \\ \Upsilon_{(1,4,7,9)} &= (((4), (1), (2), (3))) & \Upsilon_{(1,3,7,9)} &= (((3), (1), (2), (4))) \\ \Upsilon_{(1,2,7,9)} &= (((2), (1), (3), (4))) & \Upsilon_{(1,2,3,7)} &= (((3), (2), (1), (4))) \\ \Upsilon_{(1,2,4,7)} &= (((4), (2), (1), (3))) & \Upsilon_{(1,3,4,7)} &= (((4), (3), (1), (2))) \\ \Upsilon_{(1,3,6,8)} &= (((3), (4), (1), (2))) & \Upsilon_{(1,2,6,8)} &= (((2), (4), (1), (3))) \\ \Upsilon_{(1,4,5,8)} &= (((1), (4), (2), (3))) & \Upsilon_{(1,5,8,10)} &= (((1), (2), (3), (4))) \\ \Upsilon_{(1,3,5,8)} &= (((1), (3), (2), (4))) & \Upsilon_{(1,2,5,8)} &= (((2), (3), (1), (4))) \end{aligned}$$

If we apply the RSK to the elements in each set, then we get the following.

$$\begin{aligned} \text{RSK}_P(\Upsilon_{(1,2,3,4)}) &= \{((1, 2, 3, 4))\} & \text{RSK}_P(\Upsilon_{(1,2,3,5)}) &= \{((1, 2, 3), (4))\} \\ \text{RSK}_P(\Upsilon_{(1,2,4,5)}) &= \{((1, 2, 4), (3))\} & \text{RSK}_P(\Upsilon_{(1,3,4,5)}) &= \{((1, 3, 4), (2))\} \\ \text{RSK}_P(\Upsilon_{(1,3,5,6)}) &= \{((1, 3), (2, 4))\} & \text{RSK}_P(\Upsilon_{(1,2,5,6)}) &= \{((1, 2), (3, 4))\} \\ \text{RSK}_P(\Upsilon_{(1,4,5,7)}) &= \{((1, 4), (2), (3))\} & \text{RSK}_P(\Upsilon_{(1,3,5,7)}) &= \{((1, 3), (2), (4))\} \\ \text{RSK}_P(\Upsilon_{(1,2,5,7)}) &= \{((1, 2), (3), (4))\} & \text{RSK}_P(\Upsilon_{(1,2,3,6)}) &= \{((1, 2, 3), (4))\} \\ \text{RSK}_P(\Upsilon_{(1,2,4,6)}) &= \{((1, 2, 4), (3))\} & \text{RSK}_P(\Upsilon_{(1,3,4,6)}) &= \{((1, 3, 4), (2))\} \\ \text{RSK}_P(\Upsilon_{(1,4,7,9)}) &= \{((1, 4), (2), (3))\} & \text{RSK}_P(\Upsilon_{(1,3,7,9)}) &= \{((1, 3), (2), (4))\} \\ \text{RSK}_P(\Upsilon_{(1,2,7,9)}) &= \{((1, 2), (3), (4))\} & \text{RSK}_P(\Upsilon_{(1,2,3,7)}) &= \{((1, 2, 3), (4))\} \\ \text{RSK}_P(\Upsilon_{(1,2,4,7)}) &= \{((1, 2, 4), (3))\} & \text{RSK}_P(\Upsilon_{(1,3,4,7)}) &= \{((1, 3, 4), (2))\} \\ \text{RSK}_P(\Upsilon_{(1,3,6,8)}) &= \{((1, 3), (2, 4))\} & \text{RSK}_P(\Upsilon_{(1,2,6,8)}) &= \{((1, 2), (3, 4))\} \\ \text{RSK}_P(\Upsilon_{(1,4,5,8)}) &= \{((1, 4), (2), (3))\} & \text{RSK}_P(\Upsilon_{(1,5,8,10)}) &= \{((1), (2), (3), (4))\} \\ \text{RSK}_P(\Upsilon_{(1,3,5,8)}) &= \{((1, 3), (2), (4))\} & \text{RSK}_P(\Upsilon_{(1,2,5,8)}) &= \{((1, 2), (3), (4))\} \\ \\ \text{RSK}_Q(\Upsilon_{(1,2,3,4)}) &= \{((1, 1, 1, 1)), ((1, 2, 2, 2)), ((1, 1, 2, 2)), ((1, 2, 3, 3)), ((1, 2, 3, 4))\} \\ \text{RSK}_Q(\Upsilon_{(1,2,3,5)}) &= \{((1, 2, 2), (2)), ((1, 1, 2), (2)), ((1, 2, 3), (3)), ((1, 2, 3), (4))\} \\ \text{RSK}_Q(\Upsilon_{(1,2,4,5)}) &= \{((1, 2, 2), (2)), ((1, 1, 2), (2)), ((1, 2, 3), (3)), ((1, 2, 3), (4))\} \\ \text{RSK}_Q(\Upsilon_{(1,3,4,5)}) &= \{((1, 2, 2), (2)), ((1, 1, 2), (2)), ((1, 2, 3), (3)), ((1, 2, 3), (4))\} \end{aligned}$$

$$\begin{aligned}
 \text{RSK}_Q(\Upsilon_{(1,3,5,6)}) &= \{((1, 1), (2, 2)), ((1, 2), (3, 3)), ((1, 2), (3, 4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,2,5,6)}) &= \{((1, 1), (2, 2)), ((1, 2), (3, 3)), ((1, 2), (3, 4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,4,5,7)}) &= \{((1, 3), (2), (3)), ((1, 2), (3), (4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,3,5,7)}) &= \{((1, 3), (2), (3)), ((1, 2), (3), (4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,2,5,7)}) &= \{((1, 3), (2), (3)), ((1, 2), (3), (4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,2,3,6)}) &= \{((1, 3, 3), (2)), ((1, 2, 4), (3))\} \\
 \text{RSK}_Q(\Upsilon_{(1,2,4,6)}) &= \{((1, 3, 3), (2)), ((1, 2, 4), (3))\} \\
 \text{RSK}_Q(\Upsilon_{(1,3,4,6)}) &= \{((1, 3, 3), (2)), ((1, 2, 4), (3))\} \\
 \text{RSK}_Q(\Upsilon_{(1,4,7,9)}) &= \{((1, 4), (2), (3))\} & \text{RSK}_Q(\Upsilon_{(1,3,7,9)}) &= \{((1, 4), (2), (3))\} \\
 \text{RSK}_Q(\Upsilon_{(1,2,7,9)}) &= \{((1, 4), (2), (3))\} & \text{RSK}_Q(\Upsilon_{(1,2,3,7)}) &= \{((1, 3, 4), (2))\} \\
 \text{RSK}_Q(\Upsilon_{(1,2,4,7)}) &= \{((1, 3, 4), (2))\} & \text{RSK}_Q(\Upsilon_{(1,3,4,7)}) &= \{((1, 3, 4), (2))\} \\
 \text{RSK}_Q(\Upsilon_{(1,3,6,8)}) &= \{((1, 3), (2, 4))\} & \text{RSK}_Q(\Upsilon_{(1,2,6,8)}) &= \{((1, 3), (2, 4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,4,5,8)}) &= \{((1, 3), (2), (4))\} & \text{RSK}_Q(\Upsilon_{(1,5,8,10)}) &= \{((1), (2), (3), (4))\} \\
 \text{RSK}_Q(\Upsilon_{(1,3,5,8)}) &= \{((1, 3), (2), (4))\} & \text{RSK}_Q(\Upsilon_{(1,2,5,8)}) &= \{((1, 3), (2), (4))\}
 \end{aligned}$$

Here $\text{RSK}_P(\Upsilon_\Omega)$ is a singleton and the elements in each $\text{RSK}_Q(\Upsilon_\Omega)$ are connected by the standardization map θ . For example, for $\text{RSK}_Q(\Upsilon_{(1,2,3,6)})$ we have

$$\begin{array}{|c|c|c|} \hline 1 & 3 & 3 \\ \hline 2 & & \\ \hline \end{array} \xrightarrow{\theta_{(1,1,2)}^{(2,0,1,1)}} \begin{array}{|c|c|c|} \hline 1 & 1 & 4 \\ \hline 3 & & \\ \hline \end{array} \xrightarrow{\theta_{(2,1,1)}^{(1,1,1,1)}} \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & & \\ \hline \end{array}.$$

For the proof of Theorem 7.1, we generalize the definition of $w_{T,N}$ to the case when T is a row-standard Young tabloid that is not necessarily a tableau. The formula is identical; for a composition $\alpha = \text{sh}(T)$ of length $l = l(\alpha)$ let us write $L_i = \sum_{j=i+1}^l \alpha_j$ for $i \in [0, l]$ so that $L_{i-1} - L_i = \alpha_i$. Then for $N \in \mathbb{Z}$ we set $w_{T,N} \in \widetilde{\mathcal{S}}_n$ so that

$$(w_{T,N})_{T_i} = (Nn(l - i) + L_i + 1, Nn(l - i) + L_i + 2, \dots, Nn(l - i) + L_{i-1}).$$

We claim the following.

LEMMA 7.4. *Suppose that S and T are row-standard Young tabloids such that $S = R_t(T)$ for some combinatorial R -matrix R_t . If $N \gg 0$ then $w_{S,N}$ and $w_{T,N}$ are in the same left cell.*

Proof. We first claim that $w_{S,N}$ and $w_{T,N}$ are contained in the same two-sided cell parametrized by the rearrangement of $\text{sh}(S)$. By the construction of $w_{S,N}$, for $i, j \in [1, n]$ we have $i < j$ in the Shi poset of $w_{S,N}$ if and only if j is in the upper row of S than i . In this case, it is easy to show that the corresponding Greene–Kleitman partition is given by the rearrangement of $\text{sh}(S)$ as desired. Since the same result holds for $w_{T,N}$, we see that $w_{S,N}$ and $w_{T,N}$ are in the same two-sided cell.

Thus, by Theorem 4.5 it suffices to show that $\text{sgn}_\Omega(w_{S,N}) = \text{sgn}_\Omega(w_{T,N})$. Since $S_i = T_i$ for $1 \leq i < t$, from Lemma 6.1 it suffices to assume that $t = 1$. Moreover, if we again apply Lemma 6.1 to $\mathcal{R}(w_{S,N})$ and $\mathcal{R}(w_{T,N})$, respectively, then indeed it suffices to assume that S and T are two-row tabloids.

Without loss of generality we may assume that S_1 is longer than S_2 . Then $w_{S,N} \in \Gamma_S$ as before. It is also clear that $\mathcal{R}(w_{T,N}) \in \Gamma_{T^c}$ where T^c is the tableau satisfying $T_1^c = \{n + 1 - x \mid x \in T_2\}$ and $T_2^c = \{n + 1 - x \mid x \in T_1\}$ as sets. By [4, Theorem 3.18], it follows that $w_{T,N}$ is contained in the cell parametrized by $R_1(T) = S$ (i.e. the image of T^c under the affine evacuation) as desired. \square

The lemma below is another ingredient for the proof of Theorem 7.1.

LEMMA 7.5. Let $S = (S_1, \dots, S_l) \in \text{RSYT}(\alpha)$ where $\alpha = (\alpha_1, \dots, \alpha_l)$ is a composition of n of length l , $S_l = (x_1, \dots, x_{\alpha_l})$, and $\alpha_{l-1} = 0$. Let T be a row-standard Young tabloid where $T_i = S_i$ for $i \notin \{l-1, l\}$, $T_{l-1} = (x_{\alpha_l})$, and $T_l = (x_1, \dots, x_{\alpha_{l-1}})$. If $N \gg 0$ then we have $\text{sgn}_\Omega(w_{S,N}) = \text{sgn}_\Omega(w_{T,N})$.

Proof. We set $U = (U_1, \dots, U_l)$ to be the row-standard Young tabloid such that $U_i = T_i = S_i$ for $i \notin \{l-1, l\}$, $U_{l-1} = (x_2, \dots, x_{\alpha_l})$, and $U_l = (x_1)$. Note that $R_{l-1}(U) = T$ which means that $w_{T,N}$ and $w_{U,N}$ have an identical image under sgn_Ω by Theorem 7.4. Direct calculation shows that

$$w_{U,N} = w_{S,N} + \sum_{i=2}^{\alpha_l} Nn\#e_i = w_{S,N} + \sum_{i=1}^{\alpha_l} Nn\#e_i - Nn\#e_1$$

where $\#e_i$ is the standard vector with 1 on the i -th coordinate.

It follows from Lemma 6.1 that $\text{sgn}_\Omega(w_{S,N}) = \text{sgn}_\Omega(w_{S,N} + \sum_{i=1}^{\alpha_l} Nn\#e_i)$ if we set $X = [1, n] - S_l$. Now the element $w_{U,N}$ is obtained from $w_{S,N} + \sum_{i=1}^{\alpha_l} Nn\#e_i$ by replacing $1 + Nn$ on the first coordinate with 1. However, as $1 + Nn$ was already the smallest element of $w_{S,N} + \sum_{i=1}^{\alpha_l} Nn\#e_i$, lowering this entry does not affect the sign insertion process as it is inserted at the first step and not bumped by any other elements, which implies that $\text{sgn}_\Omega(w_{U,N}) = \text{sgn}_\Omega(w_{S,N} + \sum_{i=1}^{\alpha_l} Nn\#e_i)$. Thus the result follows. \square

Now we are ready to prove Theorem 7.1.

Proof of Theorem 7.1. Suppose that $T \in \Upsilon_\Omega \cap \text{RSYT}(\lambda)$ for some $\lambda \vdash n$ and assume that $(P, Q) = (\text{RSK}_P(T), \text{RSK}_Q(T))$. We regard λ as a composition of length n (with $n - l(\lambda)$ zeroes at the end) and let $Q' = \theta_{\lambda^{\text{rev}}}^{(1^n)}(Q)$. Then there exists a path $\lambda = \alpha_0, \alpha_1, \dots, \alpha_k = (1^n)$ where each α_i is a composition of n and $\alpha_i \mapsto \alpha_{i+1}$ is either rearrangement of parts or of the form $(0, \dots, 0, a, 0, \dots) \mapsto (0, \dots, 0, 1, a-1, \dots)$. (Here we allow rearrangement which permutes nonzero parts with zero parts with this does not cause any problem.) Then we may write $Q' = (\theta_{\alpha_{k-1}^{\text{rev}}}^{(1^n)} \circ \theta_{\alpha_{k-2}^{\text{rev}}}^{\alpha_{k-1}} \circ \dots \circ \theta_{\alpha_1^{\text{rev}}}^{\alpha_2} \circ \theta_{\lambda^{\text{rev}}}^{\alpha_1})(Q)$.

Let us write $T = T_0, T_1, \dots, T_k$ where $(\text{RSK}_P(T_i), \text{RSK}_Q(T_i)) = (P, (\theta_{\alpha_{i-1}^{\text{rev}}}^{\alpha_i} \circ \dots \circ \theta_{\alpha_1^{\text{rev}}}^{\alpha_2} \circ \theta_{\lambda^{\text{rev}}}^{\alpha_1})(Q))$. We claim that $T_k \in \Upsilon_\Omega$ by successively showing $\text{sgn}_\Omega(w_{T_i,N}) = \Omega$ for $N \gg 0$. The $i = 0$ case is trivial. Now if $\alpha_{i+1} = \sigma \cdot \alpha_i$ for some $\sigma \in \mathcal{S}_n$, then as we observed above we have $T_{i+1} = w_0 \sigma w_0 \cdot T_i$ where w_0 is the longest element of \mathcal{S}_n , thus the claim follows from Lemma 7.4. In the other case, first note that the shapes of T_{i+1} and T_i are $(\dots, 1, a-1, 0, \dots, 0)$ and $(\dots, 0, a, 0, \dots, 0)$ for some $a > 0$, respectively. Now direct calculation shows that T_{i+1} is obtained from T_i by the similar process to Lemma 7.5, i.e. changing $T_i = (\dots, \emptyset, (x_1, \dots, x_k), \emptyset, \dots, \emptyset)$ to $T_{i+1} = (\dots, (x_k), (x_1, \dots, x_{k-1}), \emptyset, \dots, \emptyset)$. Thus, by Lemma 7.5 the claim also follows.

As a result, for any $T \in \Upsilon_\Omega$ such that $(P, Q) = (\text{RSK}_P(T), \text{RSK}_Q(T))$ there exists $T' \in \Upsilon_\Omega \cap \text{RSYT}(1^n)$ such that $(P, \theta_{\text{sh}(T)^{\text{rev}}}^{(1^n)}(Q)) = (\text{RSK}_P(T'), \text{RSK}_Q(T'))$. Now suppose that $S \in \Upsilon_\Omega$ and set $S' \in \Upsilon_\Omega \cap \text{RSYT}(1^n)$ similarly. As $\Upsilon_\Omega \cap \text{RSYT}(1^n)$ is a singleton by Theorem 4.5, we should have $S' = T'$ which implies that $\text{RSK}_P(T) = \text{RSK}_P(T') = \text{RSK}_P(S') = \text{RSK}_P(S)$ and $\theta_{\text{sh}(T)^{\text{rev}}}^{(1^n)}(\text{RSK}_Q(T)) = \text{RSK}_Q(T') = \text{RSK}_Q(S') = \theta_{\text{sh}(S)^{\text{rev}}}^{(1^n)}(\text{RSK}_Q(S))$. Now if $\text{sh}(S) \geq \text{sh}(T)$ with respect to dominance order then $(\theta_{\text{sh}(T)^{\text{rev}}}^{(1^n)} \circ \theta_{\text{sh}(S)^{\text{rev}}}^{\text{sh}(T)})(\text{RSK}_Q(S)) = \theta_{\text{sh}(S)^{\text{rev}}}^{(1^n)}(\text{RSK}_Q(S)) = \theta_{\text{sh}(T)^{\text{rev}}}^{(1^n)}(\text{RSK}_Q(T))$ which implies that $\theta_{\text{sh}(S)^{\text{rev}}}^{\text{sh}(T)}(\text{RSK}_Q(S)) = \text{RSK}_Q(T)$ as $\theta_{\text{sh}(S)^{\text{rev}}}^{\text{sh}(T)}$ is an injection (it is an embedding of a poset).

Finally, suppose we are given $T \in \Upsilon_\Omega$ such that $(P, Q) = (\text{RSK}_P(T), \text{RSK}_Q(T))$ and $\text{sh}(T) \geq \lambda$ for some partition $\lambda \vdash n$. Let $S \in \text{RSYT}(n, \lambda^{\text{rev}})$ be such that

$(\text{RSK}_P(S), \text{RSK}_Q(S)) = (P, \theta_{\text{sh}(T)^{\text{rev}}}^{\lambda^{\text{rev}}}(Q))$ and suppose that $S \in \Upsilon_{\tilde{\Omega}}$ for some $\tilde{\Omega}$. We set $T' \in \Upsilon_{\Omega} \cap \text{RSYT}(1^n)$ and $S' \in \Upsilon_{\tilde{\Omega}} \cap \text{RSYT}(1^n)$ as above. Then $\text{RSK}_P(T') = \text{RSK}_P(T) = \text{RSK}_P(S) = \text{RSK}_P(S')$ and $\text{RSK}_Q(T') = \theta_{\text{sh}(T)^{\text{rev}}}^{(1^n)}(\text{RSK}_Q(T)) = (\theta_{\lambda^{\text{rev}}}^{(1^n)} \circ \theta_{\text{sh}(T)^{\text{rev}}}^{\lambda^{\text{rev}}})(\text{RSK}_Q(T)) = \theta_{\lambda^{\text{rev}}}^{(1^n)}(\text{RSK}_Q(S)) = \text{RSK}_Q(S')$. Since the RSK algorithm is injective it follows that $S' = T'$ which implies that $\Omega = \tilde{\Omega}$. Thus we have $S \in \Upsilon_{\Omega}$. It suffices for the proof. \square

Acknowledgements. The authors thank an anonymous referee for their useful comments.

REFERENCES

- [1] Susumu Ariki, *Robinson-Schensted correspondence and left cells*, in Combinatorial Methods in Representation Theory, Adv. Stud. Pure Math., vol. 28, American Mathematical Society, 2000, pp. 1–20.
- [2] Dan Barbasch and David Vogan, *Primitive ideals and orbital integrals in complex classical groups*, Math. Ann. **259** (1982), no. 2, 153–199.
- [3] Jonah Blasiak, *Cyclage, catabolism, and the affine Hecke algebra*, Adv. Math. **228** (2011), 2292–2351.
- [4] Michael Chmutov, Gabriel Frieden, Dongkwan Kim, Joel Brewster Lewis, and Elena Yudovina, *An affine generalization of evacuation*, Selecta Math. (N.S.) **28** (2022), no. 4, Paper No. 67, 40 pages.
- [5] Michael Chmutov, Joel Brewster Lewis, and Pavlo Pylyavskyy, *Monodromy in Kazhdan-Lusztig cells in affine type A*, Available at <https://arxiv.org/abs/1706.00471> (2017).
- [6] Michael Chmutov, Pavlo Pylyavskyy, and Elena Yudovina, *Matrix-ball construction of affine Robinson-Schensted correspondence*, Selecta Math. (N. S.) **24** (2018), no. 2, 667–750.
- [7] Adriano M. Garsia and Timothy J. McLarnan, *Relations between Young’s natural and the Kazhdan-Lusztig representations of S_n* , Adv. Math. **69** (1988), no. 1, 32–92.
- [8] David Kazhdan and George Lusztig, *Representations of Coxeter groups and Hecke algebras*, Invent. Math. **53** (1979), no. 2, 165–184.
- [9] Dongkwan Kim and Pavlo Pylyavskyy, *Asymptotic Hecke algebras and Lusztig-Vogan bijection via affine matrix-ball construction*, Available at <https://arxiv.org/abs/1902.06668> (2019), To appear in Int. Math. Res. Not. IMRN.
- [10] ———, *Two-row W -graphs in affine type A*, Adv. Math. **370** (2020).
- [11] Alain Lascoux, *Cyclic permutations on words, tableaux and harmonic polynomials*, Proceedings of the Hyderabad Conference on Algebraic Groups, 1991, pp. 323–347.
- [12] Alain Lascoux and Marcel P. Schützenberger, *Le monoïde plaxique*, Quaderni de ‘La ricerca scientifica’ **109** (1981), 129–156.
- [13] George Lusztig, *Cells in affine Weyl groups, IV*, J. Fac. Sci. Univ. Tokyo **36** (1989), 297–328.
- [14] Jian-Yi Shi, *The generalized Robinson-Schensted algorithm on the affine Weyl group of type \tilde{A}_{n-1}* , J. Algebra **139** (1991), 364–394.
- [15] Mark Shimozono, *Crystals for dummies*, Available at <https://www.aimath.org/WWN/kostka/crysdumb.pdf> (2005).
- [16] Mark Shimozono and Jerzy Weyman, *Graded characters of modules supported in the closure of a nilpotent conjugacy class*, European J. Combin. **21** (2000), 257–288.
- [17] Richard Peter Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge Studies in Advanced Mathematics, no. 62, Cambridge University Press, 1986.

DONGKWAN KIM, School of Mathematics, University of Minnesota Twin Cities, Minneapolis, MN 55455, U.S.A.
E-mail : d.kim@alum.mit.edu

PAVLO PYLYAVSKYY, School of Mathematics, University of Minnesota Twin Cities, Minneapolis, MN 55455, U.S.A.
E-mail : ppylyavs@umn.edu