



ALGEBRAIC COMBINATORICS


Christopher Ryba

Stable centres of wreath products

Volume 6, issue 2 (2023), p. 413-455.

<https://doi.org/10.5802/alco.264>

© The author(s), 2023.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Stable centres of wreath products

Christopher Ryba

ABSTRACT A result of Farahat and Higman shows that there is a “universal” algebra, FH, interpolating the centres of symmetric group algebras, $Z(ZS_n)$. We explain that this algebra is isomorphic to $\mathcal{R} \otimes \Lambda$, where \mathcal{R} is the ring of integer-valued polynomials and Λ is the ring of symmetric functions. Moreover, the isomorphism is via “evaluation at Jucys–Murphy elements”, which leads to character formulae for symmetric groups. Then, we generalise this result to wreath products $\Gamma \wr S_n$ of a fixed finite group Γ . This involves constructing wreath-product versions \mathcal{R}_Γ and $\Lambda(\Gamma_*)$ of \mathcal{R} and Λ , respectively, which are interesting in their own right (for example, both are Hopf algebras). We show that the universal algebra for wreath products, FH_Γ , is isomorphic to $\mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*)$ and use this to compute the p -blocks of wreath products.

1. INTRODUCTION

Let S_n be the n -th symmetric group, and $Z(ZS_n)$ the centre of its group ring over the integers. The centre has a basis consisting of conjugacy class sums. Farahat and Higman [6] showed that the structure constants of the multiplication with respect to this basis are integer-valued polynomials in n . For example, multiplying two transpositions can give either an element with two cycles of size 2, a 3-cycle, or the identity. If X'_μ is the sum of all elements of cycle type μ in S_n , so that $X'_{(2,1^{n-2})}$ is the sum of all transpositions, we have

$$(1) \quad (X'_{(2,1^{n-2})})^2 = 2X'_{(2^2,1^{n-4})} + 3X'_{(3,1^{n-3})} + \binom{n}{2} X'_{(1^n)}.$$

Here, we emphasise that $\binom{n}{2}$ is an integer-valued polynomial. This property of structure constants allowed Farahat and Higman to define an “interpolating” algebra, which we denote FH, with coefficients in the ring of integer-valued polynomials, \mathcal{R} . See Section 3 for a precise explanation. By construction, there are surjective “specialisation” homomorphisms $\text{FH} \rightarrow Z(ZS_n)$ for any n . The original motivation for defining the algebra FH was to give a new proof of Nakayama’s Conjecture about p -blocks of symmetric groups.

We begin by reviewing the construction of the algebra FH, in particular we explain that it is isomorphic to $\mathcal{R} \otimes \Lambda$, where Λ is the ring of symmetric functions. The Jucys–Murphy elements are key in describing this isomorphism, which leads to character formulae for S_n . We then fix a finite group Γ , and generalise the theory to the wreath products $\Gamma \wr S_n$. An algebra analogous to FH was defined by Wang in [24], and

Manuscript received 31st August 2021, revised 27th June 2022 and 26th August 2022, accepted 26th August 2022.

KEYWORDS. wreath products, Farahat–Higman algebra, Jucys–Murphy elements.

we denote it FH_Γ . (Wang mostly worked with an associated graded version of FH_Γ which he used to study Hilbert schemes of points on crepant resolutions of certain plane singularities.) The main result of this paper (Theorem 6.19) determines the algebra structure of FH_Γ via a ‘‘Jucys–Murphy evaluation’’ map that implies character formulae for wreath products of the same nature as those for the symmetric group.

We define two rings, \mathcal{R}_Γ and $\Lambda(\Gamma_*)$, which are, respectively, versions of \mathcal{R} and Λ weighted by the conjugacy classes of Γ . Then we show that $\text{FH}_\Gamma = \mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*)$, and mimic the approach of Farahat and Higman to classify the p -blocks of $\Gamma \wr S_n$.

In the appendix, we discuss some properties of \mathcal{R}_Γ and $\Lambda(\Gamma_*)$ that are not needed for the main theory, but may be of independent interest. In particular, both \mathcal{R}_Γ and $\Lambda(\Gamma_*)$ are Hopf algebras. The Hopf algebra structure on \mathcal{R}_Γ is a consequence of the fact that \mathcal{R}_Γ is a certain distribution algebra. We describe homomorphisms from \mathcal{R}_Γ to a field; similarly to the case of \mathcal{R} , such maps are parametrised by several p -adic numbers.


The paper is organised as follows. In Section 2, we review basic facts about symmetric groups, wreath products, and symmetric functions. We also briefly summarise the modular representation theory that is necessary for our applications. In Section 3, we review the theory of the Farahat–Higman algebra FH , Jucys–Murphy elements, and explain how FH is used to prove Nakayama’s Conjecture. In Section 4, we define $\Lambda(\Gamma_*)$, which we call the ring of Γ_* -weighted symmetric functions. We discuss the wreath-product Farahat–Higman algebra FH_Γ and introduce \mathcal{R}_Γ in Section 5. Then we prove the isomorphism $\text{FH}_\Gamma = \mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*)$ in Section 6, and use it to study p -blocks of wreath products in Section 7. In Appendix A we study maps from \mathcal{R}_Γ to a field, and show that both \mathcal{R}_Γ and $\Lambda(\Gamma_*)$ are Hopf algebras.

2. BACKGROUND

2.1. SYMMETRIC GROUPS AND WREATH PRODUCTS. We now introduce the notation and basic properties of symmetric groups and wreath products. Details may be found in [14, Chapter 1] (wreath products are discussed in Appendix B).

A *partition*, $\lambda = (\lambda_1, \dots, \lambda_r)$, is a finite non-increasing sequence of positive integers. The entries λ_i are called the *parts* of λ . It is also common to write $\lambda = (1^{m_1} 2^{m_2} \dots)$ to mean that λ has m_i parts of size i (this information uniquely determines λ). We will write $m_i(\lambda)$ for the number of parts of λ of size i . The size of a partition is the sum of its parts: $|\lambda| = \lambda_1 + \dots + \lambda_r$. If $|\lambda| = n$ it is common to say λ is a partition of n , and to write $\lambda \vdash n$. The length of a partition is the number of parts: $l(\lambda) = r$. We write \mathcal{P} for the set of partitions of any size.

We depict partitions with *Young diagrams*. The Young diagram of λ consists of $l(\lambda)$ rows, where the i -th row consists of λ_i boxes, left-justified. For example, the Young diagram of the partition $(5, 2, 1)$ is:




The *content* of the box in the i -th row from the top and j -th column from the left is defined to be $j - i \in \mathbb{Z}$. If the boxes of a Young diagram are labelled (usually with positive integers), we call it a *Young tableau*. If λ is a partition of n , and a Young diagram is labelled using the numbers $1, \dots, n$ (each number used once) such that numbers increase in each row from top to bottom, and increase in each column from left to right, then the corresponding Young tableau is called a *standard Young*

tableau. Suppose that $\lambda \vdash n$. For a standard Young tableau T of shape λ , an element $r \in \{1, \dots, n\}$ labels a unique box of T ; we write $c_T(r)$ for the content of the box in T labelled r . For example, if T is the standard Young tableau of shape $(3, 3, 1)$ below,

1	3	4
2	6	7
5		


then $c_T(5) = 1 - 3 = -2$, $c_T(6) = 2 - 2 = 0$, and $c_T(7) = 3 - 2 = 1$.

A *border strip* R of λ is a subset of the boxes of the Young diagram of λ satisfying the following conditions. Firstly, $\lambda \setminus R$ (i.e. the diagram of λ with the border strip R removed) should be the Young diagram of a partition. Secondly, the subset R should be contiguous (boxes sharing an edge are considered adjacent, but those sharing only a vertex are not). Finally, R should not contain any 2×2 square of boxes. Below are some examples of border strips R for the partition $(4, 2, 2)$, where in each case R is indicated by the shaded squares:



If p is a prime number, the p -core of λ is the partition obtained by successively removing border strips of size p from the diagram of λ until it is no longer possible to do so. It turns out that the resulting partition is independent of the choice of how to remove border strips of size p .

EXAMPLE 2.1. The 2-cores of the partitions (3) , $(2, 1)$, and $(1, 1, 1)$ are (1) , $(2, 1)$, and (1) respectively. The diagrams below illustrate this (grey boxes indicate a border strip of size 2 to be removed):



Consider a border strip R of size p , viewed as a sequence of adjacent boxes, starting at the bottom-left-most square, with each subsequent box either above or to the right of the previous one. Then the content of each subsequent box is 1 larger than the content of the preceding box as either the column index j increases or the row index i decreases. This means that the content of the boxes in R attain each congruence class in $\mathbb{Z}/p\mathbb{Z}$ exactly once. It follows that two partitions of the same size with the same p -core have the same multiset of content modulo p . In fact, two partitions of the same size have the same multiset of content modulo p if and only if the two partitions have the same p -core (e.g. in Example 2.1 above). Details can be found in [14, Examples 8 and 11, Section 1.1].

The n -th symmetric group, S_n , is the set of bijections from the set $\{1, \dots, n\}$ to itself, which is a group with the operation of function composition. The *cycle type* of an element $\sigma \in S_n$ is a partition λ such that $m_i(\lambda)$ is the number of cycles of size i in σ . Two elements of S_n are conjugate if and only if they have the same cycle type, so the conjugacy classes of S_n are in bijection with partitions of size n .

Let Γ be a finite group. We let $\Gamma_* = \{c_1, \dots, c_l\}$ be the set of conjugacy classes of Γ . Sometimes we will write 1 to denote the identity conjugacy class. We will need to consider the set Γ^* of irreducible representations of Γ over \mathbb{C} (we could work with any characteristic-zero splitting field of Γ , but we stick to the complex numbers for concreteness). By abuse of notation, we write $\chi \in \Gamma^*$ to mean both the irreducible representation and its character, for example we may write $\chi(1) = \dim(\chi)$. The centre of the group algebra $Z\Gamma$ is a free Z -module with a basis consisting of conjugacy-class sums. By further abuse of notation, we use the same symbol to denote a conjugacy class and the sum of its elements. This allows us to write $Z(Z\Gamma) = Z\Gamma_*$. In addition, we define the structure constants $A_{i,j}^k \in Z$ via the following equations in $Z\Gamma_*$:

$$(2) \quad c_i c_j = \sum_k A_{i,j}^k c_k.$$

By the Artin–Wedderburn theorem, $\mathbb{C}\Gamma$ is a product of matrix algebras indexed by Γ^* . Since the centre of a matrix algebra is one dimensional, we have an isomorphism of algebras $\mathbb{C}\Gamma_* = \mathbb{C}\Gamma^*$, where the implied basis of the latter space consists of orthogonal central idempotents in $\mathbb{C}\Gamma_*$.

The n -fold product of Γ with itself has an action of the symmetric group by permutation of factors. Concretely, if $(g_1, \dots, g_n) \in \Gamma^n$ and $\sigma \in S_n$, then

$$(3) \quad \sigma(g_1, \dots, g_n) = (g_{\sigma^{-1}(1)}, \dots, g_{\sigma^{-1}(n)}).$$

This defines an automorphism of Γ^n , and in fact we obtain a homomorphism $S_n \rightarrow \text{Aut}(\Gamma^n)$. The corresponding semidirect product $\Gamma^n \circ S_n$ is called the *wreath product* of Γ with S_n , and is denoted $\Gamma \wr S_n$. As a set, $\Gamma \wr S_n$ is equal to $\Gamma^n \times S_n$. We write an element $(g_1, \dots, g_n, \sigma)$ as (\mathbf{g}, σ) where $\mathbf{g} \in \Gamma^n$. Then the group operation is

$$(\mathbf{g}, \sigma)(\mathbf{h}, \rho) = (\mathbf{g}\sigma(\mathbf{h}), \sigma\rho),$$

where $\sigma(\mathbf{h})$ has the meaning in Equation (3). Both Γ^n and S_n are subgroups of $\Gamma \wr S_n$ in the obvious way. If $g \in \Gamma$, we write $g^{(i)} \in \Gamma^n$ for the element whose i -th component is g , and all other components are the identity. This notation extends linearly to give us an embedding of $Z\Gamma$ into $Z\Gamma^n$, so for example we may write $c^{(i)} = \sum_{g \in c} g^{(i)}$ for the conjugacy-class sum c embedded in the i -th component of $Z\Gamma^n$. We also let \mathbf{g}_i be the i -th entry in $\mathbf{g} \in \Gamma^n$, so we have the tautological equality

$$\mathbf{g} = \prod_{i=1}^n \mathbf{g}_i^{(i)}.$$

This also makes it simpler to write the S_n action:

$$\sigma(\mathbf{g}) = \prod_{i=1}^n \mathbf{g}_i^{(\sigma(i))}.$$

EXAMPLE 2.2. An important special case is $\Gamma = C_2$, the cyclic group of order 2. In that case, $C_2 \wr S_n$ is called *the n -th hyperoctahedral group*. It arises in Lie theory as the Weyl group of types B_n and C_n . We will use the case $\Gamma = C_2$ for examples throughout the paper.

We now describe the conjugacy classes in $\Gamma \wr S_n$. Suppose that $(\mathbf{g}, \sigma) \in \Gamma \wr S_n$, and that (i_1, \dots, i_r) is a cycle of $\sigma \in S_n$. We also say that it is a cycle of (\mathbf{g}, σ) , and we define its *type* to be the conjugacy class of $g_{i_r} \cdots g_{i_1}$. Note that $g_{i_1}(g_{i_r} \cdots g_{i_1})g_{i_1}^{-1} = g_{i_1}g_{i_r} \cdots g_{i_2}$, so that $g_{i_r} \cdots g_{i_1}$ and $g_{i_1}g_{i_r} \cdots g_{i_2}$ are conjugate. This means that the two (equal) cycles (i_1, \dots, i_r) and (i_2, \dots, i_r, i_1) have the same type, and iterating though all cyclic permutations we see that the type of a cycle is well defined. We record the sizes and types of an element of $\Gamma \wr S_n$ in a multipartition.

DEFINITION 2.3. If D is a set, we say that a multipartition indexed by D is a function from D to the set of partitions. We denote the set of multipartitions indexed by D by $\mathcal{P}(D)$. If D is not specified, we take $D = \Gamma_*$. The size of a multipartition λ is the sum of the sizes of its constituent partitions:

$$|\lambda| = \sum_{x \in D} |\lambda(x)|.$$

Similarly, the length of a multipartition is the sum of the lengths of its constituent partitions:

$$l(\lambda) = \sum_{x \in D} l(\lambda(x)).$$

To express a multipartition we write the juxtaposition of its constituent partitions with the corresponding element of D as a subscript (omitting empty partitions for brevity). If $\lambda(y)$ is the empty partition for $y \neq x$, we say that λ is concentrated in type x .

For example, $(2, 1)_{c_1}(1, 1, 1)_{c_2}$ is the multipartition in $\mathcal{P}(\Gamma_*)$ taking the value $(2, 1)$ at $c_1 \in \Gamma_*$ and taking the value $(1, 1, 1)$ at $c_2 \in \Gamma_*$, while $(3, 1)_{c_3}$ is concentrated in type c_3 .

DEFINITION 2.4. The cycle type of an element (\mathbf{g}, σ) of $\Gamma \wr S_n$ is the Γ_* -indexed multipartition μ such that for each $i \in \mathbb{Z}_{>0}$ and $c \in \Gamma_*$, $m_i(\mu(c))$ is equal to the number of cycles of type c and size i in (\mathbf{g}, σ) .

By construction, the size of the cycle type of an element of $\Gamma \wr S_n$ is n . Two elements of $\Gamma \wr S_n$ are conjugate if and only if they have the same cycle type. So the conjugacy classes of $\Gamma \wr S_n$ correspond to Γ_* -indexed multipartitions of n .

LEMMA 2.5. Let $\sigma = (i_1, \dots, i_r)$ be an r -cycle in S_n . Let $X_\sigma(c) \in \mathbb{Z}\Gamma \wr S_n$ be the sum of all elements of the form $(\mathbf{g}, \sigma) \in \Gamma \wr S_n$ where the type of the cycle σ is $c \in \Gamma_*$, and $\mathbf{g}_i = 1$ for all i not in the cycle. Then for any i_j in the cycle, we have

$$X_\sigma(c) = c^{(i_j)} X_\sigma(1) = X_\sigma(1) c^{(i_j)},$$

where $c^{(i)} = \sum_{g \in c} g^{(i)}$ is the conjugacy class sum c embedded in the i -th component of $\mathbb{Z}\Gamma^n$.

Proof. The second equality follows from the first because

$$(\mathbf{g}, \sigma) c^{(i_j)} = (\mathbf{g} c^{(i_{j+1})}, \sigma) = c^{(i_{j+1})} (\mathbf{g}, \sigma),$$

where i_{r+1} is taken to mean i_1 when $j = r$. In the second step we use the fact that c is a central element of $\mathbb{Z}\Gamma^n$. We write

$$X_\sigma(c) = \sum_{\mathbf{g}_{i_r} \cdots \mathbf{g}_{i_1} \in c} (\mathbf{g}, \sigma)$$

where $\mathbf{g}_i = 1$ for $i \neq i_1, \dots, i_r$. The condition $\mathbf{g}_{i_r} \cdots \mathbf{g}_{i_1} \in c$ may be written as

$$\mathbf{g}_{i_j} \in \mathbf{g}_{i_{j+1}}^{-1} \cdots \mathbf{g}_{i_r}^{-1} c \mathbf{g}_{i_1}^{-1} \cdots \mathbf{g}_{i_{j-1}}^{-1} = c(\mathbf{g}_{i_{j+1}}^{-1} \cdots \mathbf{g}_{i_r}^{-1} \mathbf{g}_{i_1}^{-1} \cdots \mathbf{g}_{i_{j-1}}^{-1}),$$

where we have used the fact that c is conjugation invariant to move it to the front of the product. In the case where $c = 1$, choosing all the elements other than \mathbf{g}_{i_j} arbitrarily uniquely determines \mathbf{g}_{i_j} . In the case of general c , we again may choose the elements other than \mathbf{g}_{i_j} arbitrarily, and then \mathbf{g}_{i_j} may be any element of c multiplied by the value of \mathbf{g}_{i_j} from the $c = 1$ case.

LEMMA 2.6. Let $r < s$ be positive integers and let $\sigma = (i_1, \dots, i_r)$ and $\rho = (i_r, \dots, i_s)$ be cycles in S_n of lengths r and $s - r + 1$ respectively that intersect at a single element i_r . Let $X_\sigma(c_i)$ and $X_\rho(c_j)$ be as in Lemma 2.5. Then

$$X_\sigma(c_i)X_\rho(c_j) = \sum_k A_{i,j}^k X_{\sigma\rho}(c_k),$$

where $A_{i,j}^k$ are the structure constants of $Z\Gamma_*$ appearing in Equation (2), and $\sigma\rho$ is the permutation obtained by first applying ρ and then applying σ .

Proof. We may assume $c_i = c_j = 1$ because

$$X_\sigma(c_i)X_\rho(c_j) = c_i^{(i_r)} X_\sigma(1)c_j^{(i_r)} X_\rho(1) = c_i^{(i_r)} c_j^{(i_r)} X_\sigma(1)X_\rho(1),$$

and

$$\sum_k A_{i,j}^k X_{\sigma\rho}(c_k) = \sum_k A_{i,j}^k c_k^{(i_r)} X_{\sigma\rho}(1) = c_i^{(i_r)} c_j^{(i_r)} X_{\sigma\rho}(1).$$

Now we suppose that (\mathbf{g}, σ) has all cycles of type 1, so in particular

$$\mathbf{g}_{i_r} \cdots \mathbf{g}_{i_1} = 1.$$

Similarly we consider (\mathbf{h}, ρ) with

$$\mathbf{h}_{i_s} \cdots \mathbf{h}_{i_r} = 1.$$

Then $(\mathbf{g}, \sigma)(\mathbf{h}, \rho) = (\mathbf{g}\sigma(\mathbf{h}), \sigma\rho)$, and we have

$$(\mathbf{g}\sigma(\mathbf{h}))_{i_p} = \begin{cases} \mathbf{g}_{i_1} \mathbf{h}_{i_r} & p = 1 \\ \mathbf{g}_{i_p} & 1 < p \leq r \\ \mathbf{h}_{i_p} & r < p \leq s \end{cases}$$

The type of this cycle is the conjugacy class of

$$\mathbf{h}_{i_s} \cdots \mathbf{h}_{i_{r+1}} \mathbf{g}_{i_r} \cdots \mathbf{g}_{i_1} \mathbf{h}_{i_r}.$$

But now, $\mathbf{g}_{i_r} \cdots \mathbf{g}_{i_1} = 1$, leaving $\mathbf{h}_{i_s} \cdots \mathbf{h}_{i_r}$ which equals 1. Moreover the elements $\mathbf{g}_{i_2}, \dots, \mathbf{g}_{i_r}$ and $\mathbf{h}_{i_{r+1}}, \dots, \mathbf{h}_{i_s}$ may be chosen arbitrarily, and then \mathbf{g}_{i_1} and \mathbf{h}_{i_r} are determined by the condition that the type of the cycle should be 1. As a result we get every term in the sum $X_{\sigma\rho}(1)$ exactly once.

We now turn our attention to describing the irreducible representations of $\Gamma \wr S_n$ over \mathbb{C} in terms of those of Γ and S_n . Recall that the irreducible representations S^λ of the symmetric group S_n in characteristic zero are called *Specht modules* and are labelled by partitions λ of size n . Analogously, the irreducible representations V^λ of $\Gamma \wr S_n$ over \mathbb{C} are in bijection with Γ^* -indexed multipartitions λ of size n . We explain how they can be constructed.

If $\tau = (\tau_1, \dots, \tau_l)$ is a sequence of non-negative integers adding to n , then the product of the symmetric groups S_{τ_i} is a subgroup of S_n , called a *Young subgroup*, and denoted S_τ . Here the factors S_{τ_i} are viewed as subgroups of S_n by permuting disjoint contiguous blocks of τ_i elements of $\{1, \dots, n\}$. We refer to the S_{τ_i} as the *factor groups* of the Young subgroup S_τ . The product

$$\prod_i (\Gamma \wr S_{\tau_i}) = \Gamma^n \circ S_\tau$$

is a subgroup of $\Gamma \wr S_n$ which we denote $\Gamma \wr S_\tau$. Secondly, if $\chi \in \Gamma^*$ and λ is a partition of τ_i , then

$$\chi^{\otimes \tau_i} \otimes S^\lambda$$

is a representation of $\Gamma \wr S_{\tau_i}$ where Γ^{τ_i} acts on the first factor in the natural way, while S_{τ_i} acts by permuting the tensor factors in the first term, and in the usual way on

the second term. Finally, the irreducible representation of $\Gamma \wr S_n$ indexed by λ is the following induced representation:

$$V^\lambda = \text{Ind}_{\Gamma \wr S_\tau}^{\Gamma \wr S_n} \left(\bigotimes_{\chi \in \Gamma^*} \chi^{\otimes \tau_\chi} \otimes S^{\lambda(\chi)} \right),$$

where τ is the composition of n whose parts are $\tau_\chi = |\lambda(\chi)|$, indexed by $\chi \in \Gamma^*$ in some order.

2.2. SYMMETRIC FUNCTIONS. We recall the construction of the ring of symmetric functions, Λ . Macdonald's book [14] is the authoritative reference on this topic. In Section 4, we will mimic this construction to define a related ring, $\Lambda(\Gamma_*)$.

The n -th tensor power of $Z[x]$ may be viewed as the polynomial ring $Z[x_1, \dots, x_n]$. The action of the symmetric group S_n by permutation of tensor factors may be understood as permutation of the variables. The S_n -action preserves the grading where each variable has degree 1, so the ring of S_n -invariants inherits the same grading. An important family of invariant elements are the *monomial symmetric polynomials*, m_λ , indexed by partitions λ . By definition, $m_\lambda(x_1, \dots, x_n)$ is the sum of all monomials $x_1^{a_1} x_2^{a_2} \dots x_n^{a_n}$ such that the number of exponents equal to $i \in Z_{>0}$ is $m_i(\lambda)$.

Let Λ_n be the S_n -invariants of $Z[x_1, \dots, x_n]$, and Λ_n^k be the degree k component of Λ_n . For $m > n$ we have a map $\rho_{m,n} : \Lambda_m^k \rightarrow \Lambda_n^k$ defined by setting the variables x_{n+1}, \dots, x_m to zero. For $m > k > n$, we have $\rho_{m,k} \circ \rho_{k,n} = \rho_{m,n}$, and so these $\rho_{m,n}$ define an inverse system on the Z -modules Λ_n^k (for fixed k and $n \in Z_{>0}$), and we take

$$\Lambda^k = \varprojlim \Lambda_n^k.$$

In explicit terms, Λ^k consists of sequences (q_1, q_2, \dots) , where $q_n \in \Lambda_n^k$, that satisfy the condition $\rho_{m,n}(q_m) = q_n$. It is well known that $\rho_{m,n}$ is an isomorphism when $m > n > k$, and in that case Λ_n^k has a basis of monomial symmetric functions m_λ indexed by partitions λ of size k . Moreover, the monomial symmetric functions with different numbers of variables are stable with respect to the maps $\rho_{m,n}$, i.e.

$$m_\lambda(x_1, \dots, x_n) = \rho_{m,n}(m_\lambda(x_1, \dots, x_m)) = m_\lambda(x_1, \dots, x_n, 0, \dots, 0).$$

Therefore the monomial symmetric polynomials define elements of the inverse limit Λ^k , which are called *monomial symmetric functions* and denoted m_λ . From this it follows that Λ^k is a free Z -module with basis m_λ indexed by partitions λ of size k . Finally, the ring of symmetric functions is

$$\Lambda = \bigoplus_{k=0}^{\infty} \Lambda^k,$$

which is free as a Z -module with basis consisting of the monomial symmetric functions. In fact, Λ is a graded ring because the operation of setting a variable x_m to zero respects degree and multiplication. By general properties of inverse limits, we have a canonical ring homomorphism $\Lambda \rightarrow \Lambda_n$ by sending (q_1, q_2, \dots) to q_n .

The *elementary symmetric functions* e_r are elements of Λ given by $e_r = m_{(1^r)}$. When evaluated in n variables they are

$$e_r(x_1, \dots, x_n) = \sum_{i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \dots x_{i_r},$$

namely, the sum of all products of r distinct variables. Often it is convenient to work with generating functions, in which case we have

$$\sum_{r>0} e_r(x_1, \dots, x_n) t^r = \prod_{i=1}^n (1 + x_i t).$$

A key fact is that the elementary symmetric functions generate Λ as a free polynomial algebra: $\Lambda = \mathbb{Z}[e_1, e_2, \dots]$.

2.3. CENTRAL CHARACTERS AND BLOCKS. In this subsection, we review the modular representation theory needed for our applications and for the appendix.

The notion of central character is a slight variation of a character.

DEFINITION 2.7. *Suppose that Γ is a finite group. The central character ω_c^χ of a conjugacy class $c \in \Gamma_*$ on an irreducible representation $\chi \in \Gamma^*$ is the scalar by which $c \in \mathbb{Z}\Gamma$ acts on χ (central elements always act by scalar multiplication on irreducible representations).*

There is a formula for the central characters in terms of the usual characters:

$$\omega_c^\chi = \frac{\sum_{g \in c} \chi(g)}{\chi(1)}.$$

This formula is obtained by taking the trace of c , viewed as a linear operator on the irreducible representation, and dividing by the dimension (which is the trace of the identity). Since characters are constant on conjugacy classes,

$$\omega_c^\chi = \frac{|c|\chi(g)}{\chi(1)},$$

where $|c|$ is the size of the conjugacy class c , and g is any element of c .

Central characters satisfy a variety of properties, for example they are always algebraic integers (see [5, Proposition 5.3.2]). We will use the fact that the central characters control the blocks of the modular representations of Γ . We briefly review some of the key definitions and properties of blocks.

For the rest of this section, F is an algebraically closed field of any characteristic. We still have $Z(F\Gamma) = F\Gamma_*$. Recall that the centre $F\Gamma_*$ acts on a simple $F\Gamma$ -module M by multiplication by scalars; multiplication by an element of the centre commutes with the module action, and hence defines an element of $\text{End}(M)$, which equals F by Schur's Lemma. This gives a homomorphism $F\Gamma_* \rightarrow F$, which is also called a central character. When $F = \mathbb{C}$ and $M = \chi \in \Gamma^*$, ω_c^χ is the value of this homomorphism on the conjugacy-class sum c .

DEFINITION 2.8. *The blocks of the group algebra $F\Gamma$ are the minimal indecomposable two-sided ideals B_i of $F\Gamma$.*

It is well known that $F\Gamma$ is the direct sum of its constituent blocks:

$$(4) \quad F\Gamma = \bigoplus_i B_i.$$

In particular, the intersection of distinct blocks is zero. Because the blocks are ideals, $B_i B_j \subseteq B_i \cap B_j = 0$ for $i \neq j$. This means that Equation (4) is a decomposition of F -algebras. So if we express the identity element of $F\Gamma$ as $1 = \sum_i e_i$ with $e_i \in B_i$, it follows that e_i is the identity element of B_i , and moreover $B_i = e_i F\Gamma e_i$. The upshot of this is that if M is any $F\Gamma$ -module, then

$$M = \bigoplus_i e_i M.$$

So if M is indecomposable, $e_i M$ is nonzero for exactly one value of i , and for that particular i , $e_i M = M$. This makes M into a module for some B_i . It is common to say that M belongs to the block B_i . As a result, we have a decomposition of module categories

$$F\Gamma - \text{mod} = \bigoplus_i (B_i - \text{mod}),$$

because any module M splits as a direct sum of modules $e_i M$ belonging to each block B_i . We can determine when two simple modules belong to the same block in terms of central characters.

PROPOSITION 2.9. *Two simple modules for $F\Gamma$ belong to the same block if and only if every element $Z(F\Gamma)$ acts on each of them by the same scalar (i.e. they have the same central character).*

Proof. Suppose that B_i is a block of $F\Gamma$. [19, Lemma 4.1] shows that $Z(B_i)/J(Z(B_i)) = F$, where J is the Jacobson radical. According to [19], a central character of a block B_i is defined as the composite homomorphism

$$Z(F\Gamma) \rightarrow Z(B_i) \rightarrow Z(B_i)/J(Z(B_i)) = F.$$

By definition, the Jacobson radical acts by zero on any simple module, so the action of $Z(F\Gamma)$ on a simple module factors through the above homomorphism. This shows that this definition of central character is consistent with ours. Moreover, two simple modules in the same block have the same central character. Simple modules in different blocks must have different central characters, because the central idempotents e_i act differently (e_i acts by the identity or zero depending on whether the module belongs to B_i or not).

There is a procedure for taking a representation of Γ in characteristic zero, and producing a representation in positive characteristic. The details are technical, so we only sketch the main idea, directing the interested reader to [19]. We begin with a *p-modular system* (here p is a prime), which is a triple (K, \mathcal{O}, k) defined as follows. Firstly, \mathcal{O} is a complete discrete valuation ring. Secondly, k is the residue field of \mathcal{O} which is required to be of characteristic p . Thirdly, K is the fraction field of \mathcal{O} , which is required to be of characteristic zero. We recall several facts:

- We may find a p -modular system such that every simple representation of Γ over either K or k is absolutely irreducible. For example, we may take this K to be a finite extension of the p -adic numbers, \mathbb{Q}_p , in which the polynomial $x^{|\Gamma|} - 1$ splits. Then we may also take \mathcal{O} to be the integral closure of \mathbb{Z} in K , which makes k a finite field. (See [19, Section 3.3].)
- The representation theory of Γ over \mathbb{C} and K is essentially the same. In either case, irreducible representations are defined over $\mathbb{Q}[\zeta]$, where ζ is a primitive $|\Gamma|$ -th root of unity. Then an identification of $\mathbb{Q}[\zeta]$ as a subring of \mathbb{C} with $\mathbb{Q}[\zeta]$ as a subring of K gives a correspondence of irreducible representations that respects central characters. (This follows from [19, Theorem 2.7A].)
- Given a $K\Gamma$ -module M , we may find a Γ -stable free \mathcal{O} -submodule $M_{\mathcal{O}}$ such that $KM_{\mathcal{O}} = M$ (i.e. $M_{\mathcal{O}}$ spans M over K). Taking the quotient by the maximal ideal of \mathcal{O} gives a $k\Gamma$ -module, $M_k = M_{\mathcal{O}} \otimes_{\mathcal{O}} k$. While the isomorphism class of M_k depends on the choice of integral form $M_{\mathcal{O}}$, the composition factors (with multiplicity) of M_k do not. (This is [19, Theorem 3.6].)

As a result, the operation $M \rightarrow M_k$ is well-defined on the level of Grothendieck groups:

$$d : K_0(K\Gamma - \text{mod}) \rightarrow K_0(k\Gamma - \text{mod}).$$

The map d is called the *decomposition matrix*, and its entries (with respect to the bases coming from simple modules) are called *decomposition numbers*. These definitions turn out to be independent of the choice of p -modular system. Decomposition numbers are poorly understood outside of a few special cases, and are an active area of research (one recent breakthrough was the disproof of the *James conjecture* [26] about decomposition numbers for the symmetric groups).

An easy observation from the form of the map d , is that since the centre $K\Gamma_*$ acts by scalars on a simple $K\Gamma$ -module M , any conjugacy class sum c acts by scalars on $M_{\mathcal{O}}$ and also M_k . This implies that every composition factor of M_k has the same central character. So if M is a simple $K\Gamma$ -module, M_k belongs to a single block B_i . The (necessarily disjoint) subsets of the simple (characteristic zero) modules mapping to a given (characteristic p) block are called p -blocks. The p -blocks determine a block-diagonal structure of the decomposition matrix. We will determine the p -blocks of $\Gamma \wr S_n$ in terms of the p -blocks of Γ in Theorem 7.9. This will use the following tool.

PROPOSITION 2.10 ([19, Theorem 4.2B]). *Two irreducible representations χ_1, χ_2 of $K\Gamma$ are in the same p -block if and only if*

$$\omega_c^{\chi_1} \equiv \omega_c^{\chi_2} \pmod{\pi},$$

for all $c \in \Gamma_*$, where π is a uniformiser for \mathcal{O} .

Proof. The p -block of an irreducible representation χ is determined by the central character of any composition factor of $(\chi)_k$ by Proposition 2.9. But the central character can be computed by taking the central character of χ and passing to $k = \mathcal{O}/(\pi)$. Since Γ_* is a basis for the group algebra of Γ , the equality of central characters is equivalent to the stated equations.

Our classification will use the following well-known result.

PROPOSITION 2.11. *Suppose that $|\Gamma| = p^r m$, where $p \nmid m$ and M is a simple $K\Gamma$ -module. Then the block B_i to which M_k belongs is semisimple (as a k -algebra) if and only if $p^r \mid \dim_K(M)$, in which case M_k is simple.*

Proof. The “if” direction is known as the Brauer–Nesbitt theorem (although there are also other results with that name). On the other hand, if B_i is semisimple, then the B_i -module M_k is projective for B_i and therefore for $k\Gamma$. If P is a p -Sylow subgroup of Γ , then M_k is a projective kP -module. But since p -groups only have one irreducible representation in characteristic p (the trivial representation), kP is a (not necessarily commutative) basic local algebra. This implies that a projective kP -module is free. In particular, $\dim_k(M_k)$ is a multiple of $\dim_k(kP) = p^r$.

3. THE FARAHAT–HIGMAN ALGEBRA

Let X'_μ be the sum of all elements of cycle type μ in S_n . These conjugacy-class sums define a \mathbb{Z} -basis of $Z(ZS_n)$ indexed by partitions of n . For example

$$X'_{(2,1^{n-2})} = \sum_{i < j} (i, j)$$

which is the sum of all elements of cycle type $(2, 1^{n-2})$ (i.e. transpositions) in S_n . Of course $Z(ZS_n)$ is a ring, so the product of two conjugacy-class sums should be a linear combination of such sums. Taking the example from the introduction,

$$(5) \quad (X'_{(2,1^{n-2})})^2 = 2X'_{(2^2,1^{n-4})} + 3X'_{(3,1^{n-3})} + \binom{n}{2} X'_{(1^n)},$$

which we now verify. Just as the structure constants $A^k_{i,j}$ (from Equation (2)) count the number of ways a fixed element of the conjugacy class c_k may be written as the product of an element of c_i and an element of c_j , we consider products of two transpositions, (i, j) and (k, l) . If i, j, k, l are distinct, we have

$$(i, j)(k, l) = (k, l)(i, j),$$

so each element of cycle type $(2^2, 1^{n-4})$ appears twice. If $\{i, j\}$ and $\{k, l\}$ have one element in common, $(i, j)(k, l)$ is a 3-cycle. Since

$$(i, j, k) = (i, j)(j, k) = (k, i)(i, j) = (j, k)(k, i),$$

$X'_{(3, 1^{n-3})}$ appears with multiplicity 3. Finally, if (i, j) and (k, l) move two elements in common, they are equal. Their product is therefore the identity element of S_n , and we obtain it once for each of the $\binom{n}{2}$ transpositions in S_n . The most important feature of Equation (5) is that the coefficients depend polynomially on n , the parameter of the symmetric group S_n .

DEFINITION 3.1. *The reduced cycle type of an element of a symmetric group is the partition obtained by subtracting 1 from each part of the cycle type, and ignoring any resulting parts of size zero.*

For example, the reduced cycle type of the identity is the empty partition, which we denote $?$. The reduced cycle type of a transposition is (1) . Note that the cycle type of an element of S_n can be recovered from the reduced cycle type, provided that n is known. The condition for S_n to have an element of reduced cycle type μ is $n > |\mu| + l(\mu)$. The upshot of working with reduced cycle types is that if we let X_μ be the sum of all elements of reduced cycle type μ , Equation (5) becomes

$$(6) \quad X_{(1)}^2 = 2X_{(1,1)} + 3X_{(2)} + \binom{n}{2}X_?$$

This equation is valid in $Z(ZS_n)$ for any $n \in \mathbb{Z}_{>0}$, provided X_μ is interpreted as zero if there are no elements of reduced cycle type μ in S_n (in that case X_μ as we have defined it would be an empty sum). It turns out that any product of the X_μ decomposes as a linear combination with coefficients that are polynomial in n .

DEFINITION 3.2. *The ring of integer-valued polynomials, \mathcal{R} , is the subring of $\mathbb{Q}[t]$ consisting of elements $p(t)$ such that $p(m) \in \mathbb{Z}$ for all $m \in \mathbb{Z}$.*

It is well known that \mathcal{R} is free as a \mathbb{Z} -module, with basis $\binom{t}{r}$ for $r \in \mathbb{Z}_{>0}$.

THEOREM 3.3 ([6, Theorem 2.2]). *For any partitions λ, μ, ν , there exists a unique integer-valued polynomial $\phi_{\mu, \nu}^\lambda(t)$ such that the equation*

$$X_\mu X_\nu = \sum_\lambda \phi_{\mu, \nu}^\lambda(n) X_\lambda$$

holds in $Z(ZS_n)$ for all $n \in \mathbb{Z}_{>0}$.

Using this theorem we can depart from the setting of $Z(ZS_n)$ for a specific n , and work with all n at once. This comes at the price of working over \mathcal{R} .

DEFINITION 3.4 ([6, Section 2]). *Let FH be the free \mathcal{R} -module with basis K_μ indexed by all partitions μ . Define a bilinear multiplication on FH by*

$$K_\mu K_\nu = \sum_\lambda \phi_{\mu, \nu}^\lambda(t) K_\lambda$$

where $\phi_{\mu, \nu}^\lambda(t)$ are the polynomials from Theorem 3.3, viewed as elements of the base ring \mathcal{R} . We call FH the Farahat–Higman algebra.

Because the equations in Theorem 3.3 hold in $Z(ZS_n)$ for any n , we can return from FH to $Z(ZS_n)$ via an appropriate specialisation homomorphism that sends K_μ to X_μ and evaluates the polynomial coefficients at n . It is then routine to verify that FH is a commutative, associative, unital \mathcal{R} -algebra; this reduces to considering the respective properties in $Z(ZS_n)$ for sufficiently large n .

THEOREM 3.5 ([6, Theorem 2.4]). *For each $n \in \mathbb{Z}_{>0}$ there is a surjective ring homomorphism $\Phi_n : \text{FH} \rightarrow Z(ZS_n)$ defined by*

$$\Phi_n \left(\sum_{\mu} a_{\mu}(t) K_{\mu} \right) = \sum_{\mu} a_{\mu}(n) X_{\mu},$$

where $a_{\mu}(t) \in \mathcal{R}$, and X_{μ} is the sum of all elements of S_n of reduced cycle type μ (or zero, if there are no such elements).

REMARK 3.6. Although we do not discuss it in this paper, there is a different construction and perspective of the algebra FH provided by Ivanov and Kerov [9]. Their approach transparently explains the above properties of FH. It was pointed out in [25] that this construction can be extended to wreath products, and details of this generalisation appear in [22].

The original motivation of Farahat and Higman for constructing FH was to provide a simple proof of Nakayama’s Conjecture (a theorem which describes the p -blocks of symmetric groups, originally proved by Brauer and Robinson [1]). In the remainder of this section, we show that $\text{FH} = \mathcal{R} \otimes \Lambda$, explain Jucys–Murphy elements, and discuss the theory of content evaluation character formulae for S_n . This will allow us to succinctly describe Farahat–Higman’s proof of Nakayama’s Conjecture.

3.1. ISOMORPHISM WITH SYMMETRIC FUNCTIONS. The main result of Farahat and Higman’s paper is that FH is generated by a particular set of elements. With some care, we can infer that FH is isomorphic to the ring of symmetric functions with coefficients in \mathcal{R} (Theorem 3.8).

THEOREM 3.7 ([6, Theorem 2.5]). *As an \mathcal{R} -algebra, FH is generated by the elements*

$$g_n = \sum_{\mu \vdash n} K_{\mu}$$

where $n \in \mathbb{Z}_{>0}$.

Recall that $\Lambda = \mathbb{Z}[e_1, e_2, \dots]$. It is graded:

$$\Lambda = \bigoplus_{k > 0} \Lambda^k$$

where e_r has degree r . In particular, if $\nu = (\nu_1, \nu_2, \dots, \nu_l)$ is a partition, then $e_{\nu} = e_{\nu_1} e_{\nu_2} \cdots e_{\nu_l}$ is in degree $|\nu|$, and a basis of Λ^n as a \mathbb{Z} -module is given by the set of e_{ν} as ν varies across partitions of n . The grading on Λ induces a filtration

$$\Lambda = \bigcup_{i > 0} \Lambda^{\triangleleft i},$$

where

$$\Lambda^{\triangleleft i} = \bigoplus_{k \triangleleft i} \Lambda^k,$$

and $\Lambda^{\triangleleft i}$ has \mathbb{Z} -basis consisting of e_{ν} where $|\nu| \triangleleft i$.

In fact FH is also a filtered \mathcal{R} -algebra,

$$\text{FH} = \bigcup_{i > 0} \mathcal{F}^i$$

where \mathcal{F}^i is the \mathcal{R} -submodule of FH spanned by K_{μ} with $|\mu| \triangleleft i$ (see [6, Lemma 3.9]). This filtration may be interpreted as follows. A k -cycle in a symmetric group may be written as a product of $k - 1$ transpositions (and no fewer), for example:

$$(i_1, i_2, i_3, \dots, i_k) = (i_1, i_2)(i_2, i_3) \cdots (i_{k-1}, i_k).$$

For an arbitrary permutation of cycle type $\nu = (\nu_1, \nu_2, \dots, \nu_l)$, the number of transpositions needed is

$$\sum_i (\nu_i - 1),$$

which is precisely the size of the corresponding reduced cycle type $(\nu_1 - 1, \nu_2 - 1, \dots, \nu_k - 1)$. So \mathcal{F}^i may be seen as filtering permutations according to how many transpositions are needed to construct them.

THEOREM 3.8. *There is an isomorphism $\Psi : \mathcal{R} \otimes_Z \Lambda \rightarrow \text{FH}$ of filtered \mathcal{R} -algebras defined by*

$$\Psi(e_n) = g_n = \sum_{\mu \vdash n} K_\mu.$$

We give two proofs. The first proof is intended to be in the spirit of the original work of Farahat and Higman. The second proof relies on the later work of Jucys, and it is this second proof that will generalise to the wreath-product setting.

First Proof of Theorem 3.8. The homomorphism Ψ is well defined because Λ is a free polynomial algebra, so there are no relations that need to be checked. Since $\Psi(e_i) = g_i \in \mathcal{F}^i$, it is immediate that Ψ respects the filtrations on both spaces. Additionally, Theorem 3.7 shows Ψ is surjective. However, the proof in [6] proceeds by showing that

$$\mathcal{F}^i = \mathcal{R}g_i + \sum_{\substack{j+k=i \\ j,k>1}} \mathcal{F}^j \cdot \mathcal{F}^k.$$

From this, it follows by induction on i that the restriction of Ψ to a map from the i -th filtered component of Λ to the i -th filtered component of FH is a surjection:

$$\begin{aligned} \mathcal{F}^i &= \mathcal{R}g_i + \sum_{\substack{j+k=i \\ j,k>1}} \mathcal{F}^j \cdot \mathcal{F}^k \\ &= \mathcal{R}\Psi(e_i) + \sum_{\substack{j+k=i \\ j,k>1}} \Psi(\Lambda^{\epsilon_j}) \cdot \Psi(\Lambda^{\epsilon_k}) \\ &\subseteq \Psi(\Lambda^{\epsilon_i}). \end{aligned}$$

We may pass to the fraction field of \mathcal{R} , namely the field of rational functions $\mathbb{Q}(t)$. Now Ψ restricts to a surjection of vector spaces:

$$\Psi|_{\Lambda^{\epsilon_i}} : \mathbb{Q}(t) \otimes_Z \Lambda^{\epsilon_i} \rightarrow \mathbb{Q}(t) \otimes_{\mathcal{R}} \mathcal{F}^i.$$

Since both spaces have the same dimension (each has a basis indexed by partitions of size at most i), standard linear algebra shows that since the linear map $\Psi|_{\Lambda^{\epsilon_i}}$ is surjective, it must also be injective. This in turn implies that Ψ is injective.

Theorem 3.8 is essentially a strengthening of [3, Theorem 3.1], the difference being that they work rationally (i.e. over $\mathbb{Q} \otimes_Z \mathcal{R} = \mathbb{Q}[t]$), while we work integrally (i.e. over \mathcal{R}). This will be essential when we consider modular representation theory in Section 7.

At this point it may be unclear why we choose to work with the ring of symmetric functions when the only property we have used is that it is a free polynomial algebra with one generator in each positive degree. In the next subsection, we will summarise the theory of Jucys–Murphy elements, which will allow us to interpret the isomorphism Ψ as “Jucys–Murphy evaluation” of symmetric functions. This will give us formulae for central characters. To that end, we will need the following definition.

DEFINITION 3.9. For a partition μ , the character symmetric function is $f_\mu = \Psi^{-1}(K_\mu)$ (it is an element of $\mathcal{R} \otimes_{\mathbb{Z}} \Lambda$).

EXAMPLE 3.10. By the definition of Ψ , $\Psi(e_1) = K_{(1)}$ and $\Psi(e_2) = K_{(1,1)} + K_{(2)}$. Additionally, Equation (6) shows that

$$\Psi(e_1^2) = 2K_{(1,1)} + 3K_{(2)} + \binom{n}{2}.$$

From this we conclude that $f_{(1)} = e_1$, $f_{(2)} = e_1^2 - 2e_2 - \binom{n}{2}$, and $f_{(1,1)} = 3e_2 - e_1^2 + \binom{n}{2}$.

One may consult [2, Section 5.4] for an alternative exposition of character symmetric functions. Note however, that our indexing variable μ is a partition corresponding to a reduced cycle type, while some of the literature uses partitions corresponding to cycle types, but with parts of size 1 removed.

3.2. JUCYS–MURPHY ELEMENTS. The Jucys–Murphy (“JM”) elements are a key part of what has come to be known as the Okounkov–Vershik approach to the representations of symmetric groups ([17] and [12] are both excellent references), some parts of which we briefly review before explaining the connection to the Farahat–Higman algebra.

The JM elements $L_1, \dots, L_n \in \mathbb{Z}S_n$ are sums of certain transpositions:

$$L_m = \sum_{1 \leq i < m} (i, m).$$

For example $L_1 = 0$, $L_2 = (1, 2)$, and $L_3 = (1, 3) + (2, 3)$. It is well known that the JM elements pairwise commute, so if $P(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$, the expression $P(L_1, \dots, L_n) \in \mathbb{Z}S_n$ is unambiguous. Furthermore, if $P(x_1, \dots, x_n)$ is symmetric in the x_i , then $P(L_1, \dots, L_n)$ is known to be a central element of $\mathbb{Z}S_n$.

PROPOSITION 3.11 ([16, Theorem 1.9]). If m_μ is the monomial symmetric function, then $m_\mu(L_1, \dots, L_n)$ is equal to X_μ plus a linear combination of X_ν such that either $|\nu| < |\mu|$ or $|\nu| = |\mu|$ and $l(\nu) < l(\mu)$.

Proof. Suppose that $g \in S_n$ has reduced cycle type μ . Then multiplying g by (i, j) either merges two (possibly trivial) cycles if i and j are in distinct cycles of g , or splits an individual cycle into two cycles if i and j are in the same cycle of g . The merging of two cycles has the effect of increasing the size of the reduced cycle type by 1, while splitting subtracts 1 from the size. Since each JM element is a sum of transpositions, $m_\mu(L_1, \dots, L_n)$ is a sum of products of $|\mu|$ transpositions. So to compute the leading order term, we only consider products that merge cycles at each step.

We consider L_j^r . This is a sum of products

$$(i_1, j) \cdots (i_r, j)$$

of r transpositions (i_k, j) where the i_k ($k = 1, \dots, r$) may be any numbers less than j . In order for the cycle to grow with each multiplication, it is necessary and sufficient that values of i_k must be distinct. The result of such a product will be the $(r + 1)$ -cycle

$$(i_r, i_{r-1}, \dots, i_1, j).$$

So we get every $(r + 1)$ cycle with largest element j exactly once because the elements of the cycle determine the transpositions involved in the product, and their order in the cycle determines the order of the transpositions in the product.

Finally, we note that $m_\mu(L_1, \dots, L_n)$ is a sum of products of L_j^r where the exponents r are the parts of μ in some order. By the previous paragraph, up to leading order, we get a product of $(r + 1)$ -cycles, and the length of the reduced cycle type is maximised when they do not intersect. In that case we get an element of reduced

cycle type μ , and moreover each such element arises exactly once because the length of the cycle whose largest element is j must have been the exponent of L_j in the monomial that gave rise to the cycle in consideration.

In the case where $\mu = (1^r)$ so that $m_\mu = e_r$, the above argument simplifies; it never happens that a transposition splits a cycle. This makes it possible to keep track of all the resulting permutations; we get the sum of all elements in S_n with $n - r$ cycles.

THEOREM 3.12 ([11, Section 3]). *If e_r is the r -th elementary symmetric function, then*

$$e_r(L_1, \dots, L_n) = \Phi_n(g_r) = \sum_{\mu \vdash r} X_\mu,$$

where X_μ is the sum of elements of reduced cycle type μ in S_n .

REMARK 3.13. The formulation of Theorem 3.12 appearing in [11, Section 3] states that

$$e_r(L_1, \dots, L_n) = \sum_{\substack{\mu \vdash n, \\ l(\mu) = n-r}} X'_\mu.$$

The partitions μ correspond to reduced cycle types ν of size r and length at most $n - r$. So while the sum in Theorem 3.12,

$$\sum_{\nu \vdash r} X_\nu,$$

features more summands than the first sum, the summands with $l(\nu) > n - r$ have $|\nu| + l(\nu) > n$ and therefore do not correspond to any permutations in S_n (so the corresponding terms X_ν are zero).

Now we can give the other proof of Theorem 3.8.

Second Proof of Theorem 3.8. By Theorem 3.12, $\Phi_n(\Psi(e_r)) \in Z(ZS_n)$ may be interpreted as the evaluation of the elementary symmetric function e_r at the Jucys–Murphy elements L_1, \dots, L_n . By Proposition 3.11, applying the same operation to the monomial symmetric functions m_μ , we get X_μ plus terms lower in a certain partial order. Since $\Phi_n(\Psi(m_\mu))$ is equal to X_μ plus lower-order terms, $\Psi(m_\mu)$ is equal to K_μ plus lower-order terms. Since the m_μ and K_μ are \mathcal{R} -bases of the respective spaces, this shows that Ψ is an isomorphism.

In view of Theorems 3.5 and 3.8, we have the following commutative diagram for each n ,

$$(7) \quad \begin{array}{ccc} \mathcal{R} \otimes_{\mathbb{Z}} \Lambda & \xrightarrow{\Psi} & \text{FH} \\ & \searrow^{ev_n} & \downarrow \Phi_n \\ & & Z(ZS_n) \end{array}$$

where $ev_n : \mathcal{R} \otimes_{\mathbb{Z}} \Lambda \rightarrow Z(ZS_n)$ evaluates an integer-valued polynomial at n and evaluates a symmetric function at L_1, \dots, L_n . To see that the diagram commutes, it suffices to check the generating elements e_r of Λ , which is the statement of Theorem 3.12. This suggests that FH can be thought of as being “the ring of symmetric functions evaluated at JM elements”.

Recall that the irreducible representations of CS_n are the Specht modules S^λ , indexed by partitions λ of size n . The Specht module S^λ possesses a Gelfand-Zetlin (“GZ”) basis v_T indexed by standard Young tableaux T of shape λ . A very special

property of the GZ basis is that it is diagonal for the action of the JM elements, and the eigenvalues are given by the content of boxes in T :

$$L_i v_T = c_T(i) v_T.$$

Now suppose that $P(x_1, \dots, x_n)$ is a symmetric polynomial. Evaluating at the JM elements and applying it to v_T , we obtain

$$P(L_1, \dots, L_n) v_T = P(c_T(1), \dots, c_T(n)) v_T.$$

Since P is symmetric, $P(c_T(1), \dots, c_T(n))$ does not depend on the order of the content of T . But for any T of shape λ , the multiset of $c_T(i)$ is precisely the content of the boxes in λ , in particular, it is the same for every T of shape λ . This implies that $P(L_1, \dots, L_n)$ acts by the same scalar on any v_T of fixed shape λ . This implies that $P(L_1, \dots, L_n) \in \mathbb{Z}S_n$ acts by scalar multiplication on each irreducible representation of the symmetric group, and is therefore a central element of $\mathbb{Z}S_n$ as stated earlier.

THEOREM 3.14. *The central character for S_n , ω_μ^λ (μ defines a conjugacy class via reduced cycle type), is equal to the character symmetric function f_μ evaluated at the content of the partition λ , with the integer-valued polynomial variable t evaluated at n .*

Proof. By definition ω_μ^λ is the scalar by which X_μ acts on S^λ . To calculate the scalar, we may choose an arbitrary GZ basis vector v_T and act on it by $ev_n(f_\mu) = X_\mu$. But ev_n evaluates elements of \mathcal{R} at $t = n$ and evaluates the symmetric function variables at JM elements, which act on v_T by the contents of λ .

EXAMPLE 3.15. We know from Example 3.10 that $f_{(1)} = e_1$. As (1) is the reduced cycle type of a transposition, we get that sum of all transpositions acts on the Specht module S^λ as multiplication by the sum of the contents of the boxes in the Young diagram λ . For a box in the diagram of λ , let $row(\)$ and $col(\)$ be the number of the row and column containing respectively. Then the sum of the contents of λ is

$$\sum_{\square \in \lambda} col(\square) - row(\square) = \sum_{\square \in \lambda} col(\square) - \sum_{\square \in \lambda} row(\square) = \sum_i \binom{\lambda_i + 1}{2} - \sum_j \binom{\lambda'_j + 1}{2},$$

where λ'_j are the lengths of the columns in the diagram of λ (equivalently, the parts of the partition dual to λ). Here we have used the identity $1 + 2 + \dots + m = \binom{m+1}{2}$ to sum each row/column. We have recovered the celebrated Frobenius formula [7], see also [14, Example 7, Section 1.7].

We are now able to give a proof of Nakayama’s Conjecture, which is a characterisation of the p -blocks of S_n . Although it was first proved by Brauer and Robinson [1], we take the simpler approach of Farahat and Higman.

THEOREM 3.16 ([6]). *Two irreducible representations of $\mathbb{C}S_n$ are in the same p -block if and only if they are labelled by partitions with the same p -core.*

Proof. The definition of p -block makes reference to a p -modular system (K, \mathcal{O}, k) . Proposition 2.10 provides a way to determine the blocks in terms of central characters. Note that since $\mathcal{O}/(\pi) = k$ has characteristic p , $p \in \mathcal{O}$ is contained in the ideal (π) . Suppose that $a, b \in \mathbb{Z} \subseteq \mathcal{O}$ are integers viewed as elements of \mathcal{O} . Then if $p|a - b$, certainly $\pi|a - b$. However, if $\pi|a - b$, then $a - b$ is an integer divisible by π . But if π divides an integer m coprime to p , then by Bézout’s identity, π divides $\gcd(m, p) = 1$, which is a contradiction as π is not a unit. Thus $\pi|a - b$ implies $p|a - b$, and we conclude that the a and b are congruent modulo π if and only if they are congruent modulo p .

Theorem 3.14 shows that the central characters of S_n are integers because the value of an element of \mathcal{R} at n is by definition an integer, and evaluating symmetric

polynomials with integer coefficients at integers (contents of λ) will give integers. So we are left to determine them modulo π , or equivalently, modulo p . The centre of the group algebra of S_n is $ev_n(\mathcal{R} \otimes \Lambda)$. Because $\mathcal{R} \otimes \Lambda$ is generated by \mathcal{R} and the elementary symmetric functions e_i , it is enough to consider the action of $ev_n(\mathcal{R})$ and $ev_n(e_i)$ on Specht modules S^λ . The action of $ev_n(\mathcal{R})$ is independent of the partition λ . To understand the action of the elementary symmetric polynomials, we assemble them into a generating function:

$$\sum_i ev_n(e_i)t^i = \sum_i e_i(L_1, \dots, L_n)t^i = \prod_{i=1}^n (1 + L_i t),$$

which acts on a GZ basis vector v_T (T is a standard Young tableau of shape λ) by the scalar

$$\prod_{i=1}^n (1 + c_T(i)t).$$

By the unique factorisation of polynomials in $k[t]$, this generating function determines, and is determined by, the content of λ viewed as elements of k , i.e. taken modulo $\text{char}(k) = p$. Two irreducibles are in the same p -block if and only if they have the same central characters, which holds if and only if they are labelled by partitions with the same content modulo p , which holds if and only if the partitions have the same p -core.

REMARK 3.17. Some authors (e.g. [24] and [14, Example 25, Section 1.7]) consider the associated graded algebra of FH (with respect to the filtration defined immediately before Theorem 3.8). One advantage of this is that the structure constants in the associated graded algebra are integers, rather than arbitrary elements of \mathcal{R} , so one may avoid working over \mathcal{R} altogether. One disadvantage is that the maps $ev_n : \text{FH} \rightarrow Z(ZS_n)$ become maps to an associated graded version of the centre of the group algebra (where X_μ is in degree $|\mu|$). Although this obstructs applications to modular representation theory, it turns out to be the right thing to do in the setting of Hilbert schemes. In fact, this associated graded version of $Z(ZS_n)$ is isomorphic to $H^*(\text{Hilb}^n(\mathbb{A}_{\mathbb{C}}^2), Z)$, the cohomology ring (with Z coefficients) of the Hilbert scheme of n points in the plane. This was shown in [13]. We will not discuss Hilbert schemes any further in this paper.

4. Γ_* -WEIGHTED SYMMETRIC FUNCTIONS

We now generalise the construction of the ring of symmetric functions in a way that incorporates Γ_* , which will define a ring $\Lambda(\Gamma_*)$ that will play a central role in this paper. Let Q be the subring of $Z\Gamma_*[x]$ consisting of polynomials whose constant term is a multiple of the identity (rather than an arbitrary element of $Z\Gamma_*$). Consider the n -th tensor power (over Z) of Q , which has an action of S_n by permutation of tensor factors. As shorthand, for $c \in Z\Gamma_*$ we write

$$x_i^r(c) = 1^{\otimes(i-1)} \otimes cx^r \otimes 1^{\otimes(n-i)}.$$

This means that $x_i^r(c)x_i^s(c') = x_i^{r+s}(cc')$, and that $x_i^r(c) + x_i^r(c') = x_i^r(c + c')$. Since Q has a grading (inherited from $Z\Gamma_*[x]$), there is a grading on the ring of S_n invariants of $Q^{\otimes n}$. We write $\Lambda_n(\Gamma_*)$ for the S_n -invariants of $Q^{\otimes n}$, and $\Lambda_n^k(\Gamma_*)$ for the degree k component of $\Lambda_n(\Gamma_*)$. As before, for $m > n$ there are homomorphisms $\rho_{m,n} : \Lambda_m^k(\Gamma_*) \rightarrow \Lambda_n^k(\Gamma_*)$ which evaluate the polynomial variables of all tensor factors past the n -th at zero. These define an inverse system, and we write

$$\Lambda^k(\Gamma_*) = \varprojlim \Lambda_n^k(\Gamma_*).$$

REMARK 4.1. It may seem unnatural to work with the ring Q rather than the full polynomial ring $Z\Gamma_*[x]$. The reason we do this is that evaluating elements of Q at zero yields an element of Z rather than $Z\Gamma_*$, so the codomain of $\rho_{m,n}$ is

$$\Lambda_n^k(\Gamma_*) \otimes Z^{\otimes(m-n)} = \Lambda_n^k(\Gamma_*),$$

rather than

$$\Lambda_n^k(\Gamma_*) \otimes (Z\Gamma_*)^{\otimes(m-n)},$$

which, being different from $\Lambda_n^k(\Gamma_*)$, would not allow us to construct an inverse system. Later we will account for these “missing” constant terms (see Theorem 5.18), using the ring \mathcal{R}_Γ which is defined in Section 5.

The ring Q has a basis consisting of elements of the form cx^j where $c \in \Gamma_*$ and $j \in Z_{>0}$ (we require $c = 1$ if $j = 0$). Thus the ring $Q^{\otimes n}$ has a basis consisting of pure tensors in this basis of Q . We refer to such a pure tensor as a Γ_* -weighted monomial and note that the S_n action sends Γ_* -weighted monomials to other Γ_* -weighted monomials.

DEFINITION 4.2. Let λ be a multipartition indexed by Γ_* . The Γ_* -weighted monomial symmetric polynomial, $m_\lambda(x_1, \dots, x_n) \in Q^{\otimes n}$ is the sum of all Γ_* -weighted monomials in $Q^{\otimes n}$ that contain cx^j with $j > 1$ as a tensor factor exactly $m_j(\lambda(c))$ times. There is no restriction on the number of times 1 may appear as a tensor factor.

It is immediate that the degree of $m_\lambda(x_1, \dots, x_n)$ is $|\lambda|$. Since the Γ_* -weighted monomial symmetric polynomials are orbit sums for the S_n action on our basis of $Q^{\otimes n}$, it follows that $m_\lambda(x_1, \dots, x_n)$ with $|\lambda| = k$ span $\Lambda_n^k(\Gamma_*)$, and the nonzero ones form a basis of this space. Moreover $m_\lambda(x_1, \dots, x_n)$ is nonzero as soon as there are enough tensor factors to accommodate all the basis vectors prescribed by λ , i.e. as soon as $n > l(\lambda)$. Finally, we observe that

$$\rho_{m,n}(m_\lambda(x_1, \dots, x_m)) = m_\lambda(x_1, \dots, x_n),$$

so these elements define an element of the inverse limit $\Lambda^k(\Gamma_*)$.

DEFINITION 4.3. The Γ_* -weighted monomial symmetric function m_λ is the element of $\Lambda^k(\Gamma_*)$ defined by the sequence of elements $m_\lambda(x_1, \dots, x_n)$ as n varies (here $k = |\lambda|$).

It now follows that $\Lambda^k(\Gamma_*)$ is free as a Z -module with basis m_λ indexed by all multipartitions λ of size k .

DEFINITION 4.4. The ring of Γ_* -weighted symmetric functions is

$$\Lambda(\Gamma_*) = \bigoplus_{k=0}^{\infty} \Lambda^k(\Gamma_*).$$

Similarly to the case of ordinary symmetric functions, the fact that evaluating a polynomial variable at zero is a ring homomorphism implies that $\Lambda(\Gamma_*)$ inherits the structure of a graded ring. In fact, if Γ is the trivial group, then $\Lambda(\Gamma_*) = \Lambda$. Just as for Λ , formal properties of inverse limits automatically give us ring homomorphisms

$$\Lambda(\Gamma_*) \rightarrow \Lambda_n(\Gamma_*).$$

EXAMPLE 4.5. We demonstrate how to multiply two degree 1 elements of $\Lambda(\Gamma_*)$. Suppose that $c_r \in \Gamma_*$ is a conjugacy class. Then $m_{(1)_{c_r}}$ is the element of $\Lambda^1(\Gamma)$ defined by the sequence of elements $\sum_{i=1}^n x_i(c_r) \in \Lambda_n^1(\Gamma_*)$. So we must express products of such elements in terms of Γ_* -weighted monomial symmetric polynomials $m_\lambda(x_1, \dots, x_n)$.

Recall that we place subscripts on partitions to indicate multipartitions. For any number of variables, we have

$$\begin{aligned} m_{(1)_{c_r}}^2 &= \left(\sum_i x_i(c_r) \right)^2 \\ &= \sum_i x_i^2(c_r) + \sum_{i \neq j} x_i(c_r)x_j(c_r) \\ &= \sum_i \sum_{c_s \in \Gamma_*} A_{r,r}^s x_i^2(c_s) + 2 \sum_{i < j} x_i(c_r)x_j(c_r) \\ &= \sum_{c_s \in \Gamma_*} A_{r,r}^s m_{(2)_{c_s}} + 2m_{(1,1)_{c_r}} \end{aligned}$$

(where $A_{i,j}^k$ are the structure constants of $Z\Gamma_*$), and therefore the equality between the first and last quantities may be interpreted as holding in $\Lambda^2(\Gamma_*)$. Similarly, if $c_r, c_s \in \Gamma_*$ are distinct conjugacy classes, then

$$\begin{aligned} m_{(1)_{c_r}} m_{(1)_{c_s}} &= \left(\sum_i x_i(c_r) \right) \left(\sum_j x_j(c_s) \right) \\ &= \sum_i x_i^2(c_r c_s) + \sum_{i \neq j} x_i(c_r)x_j(c_s) \\ &= \sum_{c_t \in \Gamma_*} A_{r,s}^t m_{(2)_{c_t}} + m_{(1)_{c_r}(1)_{c_s}}. \end{aligned}$$

Analogously to how Z -linear combinations of conjugacy-class sums and Z -linear combinations of central idempotents in $C\Gamma$ define different integral forms of $C\Gamma_*$, the ring $\Lambda(\Gamma_*)$ is a different integral form of the $|\Gamma^*|$ -th tensor power of Λ .

PROPOSITION 4.6. *We have*

$$\mathbb{C} \otimes \Lambda(\Gamma_*) = \mathbb{C} \otimes \Lambda^{\otimes |\Gamma^*|}.$$

Proof. We use the fact that $\mathbb{C} \otimes Z\Gamma_* = C\Gamma_* = C\Gamma^*$, where standard basis vectors in $C\Gamma^*$ are the orthogonal central idempotents e_χ in the group algebra $C\Gamma$ associated to irreducible representations $\chi \in \Gamma^*$. Then $e_\chi e_\psi = \delta_{\chi,\psi} e_\chi$ (where $\delta_{\chi,\psi}$ is the Kronecker delta). Working over \mathbb{C} allows us to define elements $m_{\lambda}^{irr} \in \Lambda(\Gamma_*)$ similar to the Γ_* -weighted monomial symmetric functions, but instead of taking sums of pure tensors in the basis Γ_* of $C\Gamma_*$, we use the basis e_χ . So $x_i^r(e_\chi)$ form a basis of (the positive degree part of) $\mathbb{C} \otimes Q$, and m_{λ}^{irr} is constructed by taking a sum of a S_n -orbit of a pure tensor of elements of this basis. Hence $\lambda \in \mathcal{P}(\Gamma^*)$ is indexed by Γ^* rather than Γ_* . The upshot of this is that since

$$x_i^r(e_\chi)x_i^s(e_\psi) = \delta_{\chi,\psi} x_i^{r+s}(e_\chi),$$

terms corresponding to different $\chi \in \Gamma^*$ do not interact. More precisely, suppose that λ is a multipartition and let $\lambda_\chi = \lambda(\chi)_\chi$ be the multipartition concentrated in type χ taking the value $\lambda(\chi)$. Then we have

$$m_{\lambda}^{irr} = \prod_{\chi \in \Gamma^*} m_{\lambda_\chi}^{irr}.$$

This is because each monomial in m_{λ}^{irr} arises uniquely as a product of monomials from each $m_{\lambda_\chi}^{irr}$, each involving distinct variables. Conversely, any product of monomials

from $m_{\lambda_\chi}^{irr}$ involving the same variable for different χ is zero because $x_i^r(e_\chi)x_i^s(e_\psi) = \delta_{\chi,\psi}x_i^{r+s}(e_\chi)$. This immediately shows that

$$\mathbb{C} \otimes \Lambda(\Gamma_*) = \bigotimes_{\chi \in \Gamma_*} \Lambda(\chi),$$

where $\Lambda(\chi)$ has \mathbb{C} -basis $m_{\lambda_\chi}^{irr}$, where λ is concentrated in type χ . Finally, it remains to observe that $\Lambda(\chi)$ is a ring isomorphic to $\mathbb{C} \otimes \Lambda$ via the \mathbb{C} -linear map taking $m_{\lambda_\chi}^{irr}$ to $m_{\lambda(\chi)}$. This is clearly a bijection. To see that it respects multiplication note that the multiplication in $\Lambda(\chi)$ is determined by the following rule for products in $\mathbb{C} \otimes Q$: $x_i^r(e_\chi)x_i^s(e_\chi) = x_i^{r+s}(e_\chi)$. In Λ the corresponding relation reads $x_i^r \cdot x_i^s = x_i^{r+s}$.

EXAMPLE 4.7. Suppose that $\Gamma = C_2 = \{1, \gamma\}$ is the cyclic group of order two (so $\gamma^2 = 1$). We show that $\Lambda(C_{2*})$ is not isomorphic to $\Lambda \otimes \Lambda$ as a graded ring by considering the module of indecomposables. Suppose that $A = \bigoplus_{i=0}^\infty A_i$ is a graded algebra and $I = \bigoplus_{i=1}^\infty A_i$ is the corresponding augmentation ideal. The *module of indecomposables* is defined to be I/I^2 , which is a (graded) module for A/I . We compute the degree two component of I/I^2 when A is either $\Lambda(C_{2*})$ or $\Lambda \otimes \Lambda$.

Recall that $\Lambda = \mathbb{Z}[e_1, e_2, \dots]$. If $A = \Lambda \otimes \Lambda$, then the degree 1 component of A has basis $e_1 \otimes 1, 1 \otimes e_1$, while the degree 2 component has basis $e_2 \otimes 1, e_1^2 \otimes 1, e_1 \otimes e_1, 1 \otimes e_2, 1 \otimes e_1^2$. Then the degree 2 component of I^2 has basis $e_1^2 \otimes 1, e_1 \otimes e_1, 1 \otimes e_1^2$. We conclude that the degree 2 component of the module of indecomposables of A is \mathbb{Z}^2 (spanned by the classes of $e_2 \otimes 1$ and $1 \otimes e_2$).

Analogously, for $A = \Lambda(C_{2*})$, the degree 1 component has basis $m_{(1)_1}, m_{(1)_\gamma}$, while the degree 2 component has basis $m_{(1,1)_1}, m_{(2)_1}, m_{(1)_1(1)_\gamma}, m_{(1,1)_\gamma}, m_{(2)_\gamma}$. The degree 2 component of I^2 is spanned by

$$\begin{aligned} m_{(1)_1}^2 &= m_{(2)_1} + 2m_{(1,1)_1} \\ m_{(1)_1}m_{(1)_\gamma} &= m_{(2)_\gamma} + m_{(1)_1(1)_\gamma} \\ m_{(1)_\gamma}^2 &= m_{(2)_1} + 2m_{(1,1)_\gamma}, \end{aligned}$$

which we computed in Example 4.5. From this we see that the degree 2 component of I/I^2 is isomorphic to $\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z}$. We conclude that $\Lambda(C_{2*}) \neq \Lambda^{\otimes 2}$. We discuss the algebraic structure of $\Lambda(\Gamma_*)$ further in the appendix.

5. WREATH-PRODUCT FARAHAT–HIGMAN ALGEBRAS

We now extend the results of Section 3 from the symmetric group S_n to the wreath products $\Gamma \wr S_n = \Gamma^n \circ S_n$, where Γ is a finite group. There is a wreath-product version of the Farahat–Higman algebra, which is also related to the ring of symmetric functions. The homomorphisms Φ_n, ev_n, Ψ all generalise to the wreath-product setting. We use the same notation for the wreath-product versions of these maps because taking Γ to be the trivial group (so that $\Gamma \wr S_n = S_n$), we recover the maps from Section 3.

Recall that the conjugacy classes of $\Gamma \wr S_n$ correspond to multipartitions of total size n indexed by Γ_* . We use the boldface Greek letters λ, μ, ν to indicate multipartitions. Thus $\mu(c)$ means the partition in μ indexed by $c \in \Gamma_*$. Similarly to symmetric groups, we have a notion of reduced cycle type.

DEFINITION 5.1 ([24, Subsection 2.3]). *The partially-reduced cycle type of an element of $\Gamma \wr S_n$ of cycle type μ is the multipartition obtained by subtracting 1 from each part of $\mu(1)$, and ignoring any resulting parts of size zero.*

This is the same as the case of symmetric groups, but applied only to the partition labelled by the identity conjugacy class. The group $\Gamma \wr S_n$ contains an element of

partially-reduced cycle type μ if and only if $n > |\mu| + l(\mu(1))$. Later in this section we will need to introduce the notion of *fully-reduced cycle type* which will involve reducing each partition in μ (this is why we do not use the terminology of “modified type” from [24]).

The centre of the integral group ring of $\Gamma \wr S_n$ has a \mathbb{Z} -basis consisting of conjugacy class sums. We let X_μ be the sum of all elements of partially-reduced cycle type μ (which is zero if there are no such elements). Then we have a direct analogue of Theorem 3.3.

THEOREM 5.2 ([24, Theorem 2.13]). *For any multipartitions λ, μ, ν , there exists a unique integer-valued polynomial $\phi_{\mu, \nu}^\lambda(t)$ such that the equation*

$$X_\mu X_\nu = \sum_{\lambda} \phi_{\mu, \nu}^\lambda(n) X_\lambda$$

holds in $Z(\mathbb{Z}\Gamma \wr S_n)$ for all $n \in \mathbb{Z}_{>0}$.

This allows us to define an algebra analogous to FH, but for wreath products.

DEFINITION 5.3. *Let FH_Γ be the free \mathcal{R} -module with basis K_μ indexed by all multipartitions $\mu \in \mathcal{P}(\Gamma_*)$. Define a bilinear multiplication on FH_Γ by*

$$K_\mu K_\nu = \sum_{\lambda} \phi_{\mu, \nu}^\lambda(t) K_\lambda$$

where $\phi_{\mu, \nu}^\lambda(t)$ are the polynomials from Theorem 5.2, viewed as elements of the base ring \mathcal{R} .

As in the case of FH, FH_Γ is a commutative, associative, unital \mathcal{R} -algebra, and has specialisation homomorphisms.

THEOREM 5.4 ([24, Section 2.5]). *For each $n \in \mathbb{Z}_{>0}$ there is a surjective ring homomorphism $\Phi_n : \text{FH}_\Gamma \rightarrow Z(\mathbb{Z}\Gamma \wr S_n)$ defined by*

$$\Phi_n \left(\sum_{\mu} a_\mu(t) K_\mu \right) = \sum_{\mu} a_\mu(n) X_\mu,$$

where $a_\mu(t) \in \mathcal{R}$, and X_μ is the sum of all elements of $\Gamma \wr S_n$ of partially-reduced cycle type μ (or zero, if there are no such elements).

As we now show, the family of homomorphisms Φ_n separates elements of FH_Γ . So to prove an identity in FH_Γ , it is enough to pass to $Z(\mathbb{Z}\Gamma \wr S_n)$ by applying Φ_n .

PROPOSITION 5.5. *Suppose $x, y \in \text{FH}_\Gamma$ satisfy $\Phi_n(x) = \Phi_n(y)$ for all sufficiently large positive integers n . Then $x = y$.*

Proof. Consider an element

$$z = \sum_{\mu} a_\mu(t) K_\mu.$$

Then

$$\Phi_n(z) = \sum_{\mu} a_\mu(n) X_\mu,$$

where X_μ is nonzero provided $n > |\mu| + l(\mu(1))$, and such nonzero elements form a basis of $Z(\mathbb{Z}\Gamma \wr S_n)$. This means that $\Phi_n(z)$ determines $a_\mu(n)$ for all n sufficiently large, which is a Zariski-dense subset of \mathbb{Z} . Hence $\Phi_n(z)$ determines each coefficient $a_\mu(t) \in \mathcal{R}$, and so $\Phi_n(z)$ determines z itself.

We have introduced FH_Γ as a \mathcal{R} -algebra. However, it will turn out that it is better to view it as an algebra over a Γ_* -weighted version of \mathcal{R} , which we now work towards defining.

DEFINITION 5.6. Consider formal power series in variables x_c indexed by $c \in \Gamma_*$. We encode monomials in the x_c using functions $\mathbf{N} : \Gamma_* \rightarrow \mathbb{Z}_{>0}$ that record the exponent of each variable:

$$x^{\mathbf{N}} = \prod_{c \in \Gamma_*} x_c^{\mathbf{N}(c)}.$$

Additionally, let $|\mathbf{N}| = \sum_c \mathbf{N}(c)$, so that $\deg(x^{\mathbf{N}}) = |\mathbf{N}|$.

Note that the set of such functions \mathbf{N} is $\mathbb{Z}_{>0}^{\Gamma_*}$.

DEFINITION 5.7. Let $\mathfrak{g} = \mathbb{Q}\Gamma_* = Z(\mathbb{Q}\Gamma)$, viewed as a Lie algebra via the associative multiplication (it is abelian). There is a canonical map $\iota : \mathfrak{g} \rightarrow \mathcal{U}(\mathfrak{g})$ from \mathfrak{g} to its universal enveloping algebra, which we extend to formal power series coefficientwise. We define

$$T : \mathfrak{g}[[x_{c_1}, \dots, x_{c_l}]] \rightarrow \mathcal{U}(\mathfrak{g})[[x_{c_1}, \dots, x_{c_l}]]$$

via

$$T \left(\sum_{\mathbf{N} \in \mathbb{Z}_{>0}^{\Gamma_*}} u_{\mathbf{N}} x^{\mathbf{N}} \right) = \sum_{\mathbf{N} \in \mathbb{Z}_{>0}^{\Gamma_*}} \iota(u_{\mathbf{N}}) x^{\mathbf{N}}.$$

where $u_{\mathbf{N}} \in \mathfrak{g}$.

REMARK 5.8. The map T allows us to distinguish between the associative multiplication in \mathfrak{g} , and the multiplication in the universal enveloping algebra. For example, if $u, v \in \mathfrak{g}$, then $uv \in \mathfrak{g}$, while $T(u)T(v)$ is an element of $\mathcal{U}(\mathfrak{g})$ that lies in degree two with respect to the Poincaré-Birkhoff-Witt (PBW) filtration. It may seem peculiar to consider the universal enveloping algebra of an abelian Lie algebra when the result is simply the symmetric algebra of the Lie algebra. The ring we are about to define, \mathcal{R}_Γ , is actually a distribution algebra, and so the universal enveloping algebra is actually the natural place to construct it. This is discussed further in the appendix. There is also similar situation when interpolating Grothendieck rings of wreath products (rather than centres of group algebras of wreath products) where \mathfrak{g} is the Grothendieck ring of a tensor category. In that case, \mathfrak{g} is not necessarily abelian and the universal enveloping algebra is the correct construction. The interested reader is directed to [20] and [21].

DEFINITION 5.9. Let $\Omega \in \mathcal{U}(\mathfrak{g})[[x_{c_1}, \dots, x_{c_l}]]$ be the generating function defined as follows,

$$\Omega(x_{c_1}, \dots, x_{c_l}) = \exp \left(T \left(\log \left(1 + \sum_{c \in \Gamma_*} cx_c \right) \right) \right).$$

Here $\log \left(1 + \sum_{c \in \Gamma_*} cx_c \right)$ is viewed as an element of $\mathfrak{g}[[x_{c_1}, x_{c_2}, \dots, x_{c_l}]]$ (note that expanding the power series involves the associative algebra structure of \mathfrak{g} , not just the Lie algebra structure). Writing Ω in terms of monomials, we let $B_{\mathbf{N}} \in U(\mathfrak{g})$ be the coefficients:

$$\Omega(x_{c_1}, \dots, x_{c_l}) = \sum_{\mathbf{N} \in \mathbb{Z}_{>0}^{\Gamma_*}} B_{\mathbf{N}} x^{\mathbf{N}}.$$

We let \mathcal{R}_Γ be the \mathbb{Z} -submodule of $\mathcal{U}(\mathfrak{g})$ spanned by all the $B_{\mathbf{N}}$, and we call \mathcal{R}_Γ the ring of Γ_* -weighted integer-valued polynomials.

We will shortly show that \mathcal{R}_Γ is indeed a ring, but first, the following example justifies the analogy to integer-valued polynomials.

EXAMPLE 5.10. Suppose that Γ is the trivial group, so that there is only one conjugacy class c , and it obeys $c^2 = c$. Then,

$$\begin{aligned} \log \left(1 + \sum_{c \in \Gamma_*} cx_c \right) &= \sum_{i>1} \frac{(-1)^{i-1}}{i} c^i x_c^i \\ &= c \sum_{i>1} \frac{(-1)^{i-1}}{i} x_c^i \\ &= c \log(1 + x_c). \end{aligned}$$

Hence Ω becomes

$$\begin{aligned} \Omega(x_c) &= \exp(T(c \log(1 + x_c))) \\ &= \exp(\log(1 + x_c)T(c)) \\ &= (1 + x_c)^{T(c)} \\ &= \sum_{N>0} \binom{T(c)}{N} x_c^N. \end{aligned}$$

So the $B_{\mathbf{N}}$ are simply binomial coefficients (with parameter $T(c)$) and $\mathcal{R}_\Gamma = \mathcal{R}$.

PROPOSITION 5.11. *We have the following relation:*

$$\Omega(x_{c_1}, \dots, x_{c_l}) \Omega(y_{c_1}, \dots, y_{c_l}) = \Omega(z_{c_1}, \dots, z_{c_l}),$$

where

$$z_{c_i} = x_{c_i} + y_{c_i} + \sum_{j,k} A_{j,k}^i x_{c_j} y_{c_k}.$$

Proof. Let $S(x) = 1 + \sum_{c \in \Gamma_*} cx_c$, and similarly define $S(y)$ and $S(z)$. Then

$$\begin{aligned} S(x)S(y) &= \left(1 + \sum_{c \in \Gamma_*} cx_c \right) \left(1 + \sum_{c \in \Gamma_*} cy_c \right) \\ &= 1 + \sum_{c_j \in \Gamma_*} c_j x_{c_j} + \sum_{c_k \in \Gamma_*} c_k y_{c_k} + \sum_{c_i} A_{j,k}^i c_i x_{c_j} y_{c_k} \\ &= S(z). \end{aligned}$$

Because we are working with commutative algebras, we may compute as follows:

$$\begin{aligned} \Omega(x_{c_1}, \dots, x_{c_l}) \Omega(y_{c_1}, \dots, y_{c_l}) &= \exp(T(\log(S(x)))) \exp(T(\log(S(y)))) \\ &= \exp(T(\log(S(x)) + T(\log(S(y)))) \\ &= \exp(T(\log(S(x)) + \log(S(y)))) \\ &= \exp(T(\log(S(x)S(y)))) \\ &= \exp(T(\log(S(z)))) \\ &= \Omega(z_{c_1}, \dots, z_{c_l}). \end{aligned}$$

LEMMA 5.12. *As an element of $\mathcal{U}(\mathfrak{g})$, $B_{\mathbf{N}}$ lies in PBW filtration degree $|\mathbf{N}|$ and has leading order term*

$$\prod_{c \in \Gamma_*} \frac{T(c)^{\mathbf{N}(c)}}{\mathbf{N}(c)!}.$$

Proof. The $B_{\mathbf{N}}$ are defined by the equation

$$\Omega(x_{c_1}, \dots, x_{c_l}) = \exp \left(T \left(\log \left(1 + \sum_{c \in \Gamma_*} cx_c \right) \right) \right) = \sum_{\mathbf{N} \in Z_{>0}^{\Gamma_*}} B_{\mathbf{N}} x^{\mathbf{N}}.$$

The expression

$$T \left(\log \left(1 + \sum_{c \in \Gamma_*} cx_c \right) \right)$$

is in PBW degree 1. When the logarithm is expanded as a power series in the x_c variables, the lowest order term is

$$\sum_{c \in \Gamma_*} T(c)x_c.$$

In particular, there is no constant term. This means that $\Omega(x_{c_1}, \dots, x_{c_l})$ is a sum of products of terms whose PBW degree is less than or equal to their degree in the x_c variables. Since $B_{\mathbf{N}}$ is the coefficient of $x^{\mathbf{N}}$, it is contained in PBW filtration degree $|\mathbf{N}|$. Moreover, to compute the leading term of the $B_{\mathbf{N}}$, we neglect all but the lowest degree monomials in x_c ; to leading order, Ω is approximated by

$$\exp \left(\sum_{c \in \Gamma_*} T(c)x_c \right) = \prod_{c \in \Gamma_*} \exp(T(c)x_c).$$

The leading term of $B_{\mathbf{N}}$ is found by taking the coefficient of $x^{\mathbf{N}}$.

PROPOSITION 5.13. *The $B_{\mathbf{N}}$ are a Z -basis of \mathcal{R}_{Γ} . Additionally, the multiplication in $\mathcal{U}(\mathfrak{g})$ induces a multiplication on \mathcal{R}_{Γ} , making it into a unital commutative ring.*

Proof. The $B_{\mathbf{N}}$ span \mathcal{R}_{Γ} by definition. It follows from Lemma 5.12 and the PBW theorem that the $B_{\mathbf{N}}$ form a \mathbb{Q} -basis of $\mathcal{U}(\mathfrak{g})$, so they are also linearly independent.

We note that when \mathbf{N} is zero, $B_{\mathbf{N}}$ is the constant term of Ω , namely the identity in $\mathcal{U}(\mathfrak{g})$, so \mathcal{R}_{Γ} has an identity element. To show that \mathcal{R}_{Γ} is a commutative ring, we must show that the product of two basis elements is a linear combination of basis elements. This amounts to showing that the coefficient of any monomial $x^{\mathbf{N}}y^{\mathbf{M}}$ in

$$\Omega(x_{c_1}, \dots, x_{c_l})\Omega(y_{c_1}, \dots, y_{c_l})$$

is contained in \mathcal{R}_{Γ} . We invoke Proposition 5.11, which expresses this product as

$$\Omega(z_{c_1}, \dots, z_{c_l}) = \sum_{\mathbf{K} \in Z_{>0}^{\Gamma_*}} B_{\mathbf{K}} z^{\mathbf{K}},$$

where $z_{c_i} = x_{c_i} + y_{c_i} + \sum_{j,k} A_{j,k}^i x_{c_j} y_{c_k}$, which is a Z -linear combination of monomials in the x and y variables. Thus the expansion in terms of $x^{\mathbf{N}}y^{\mathbf{M}}$ will have Z -linear combinations of the $B_{\mathbf{K}}$ as coefficients.

PROPOSITION 5.14. *The set of $B_{\mathbf{N}}$ in \mathcal{R}_{Γ} indexed by \mathbf{N} with $\mathbf{N}(c) = 0$ for $c \neq 1$ forms a subring of \mathcal{R}_{Γ} isomorphic to \mathcal{R} . As a module over \mathcal{R} , \mathcal{R}_{Γ} is free with basis $B_{\mathbf{M}}$ indexed by \mathbf{M} with $\mathbf{M}(1) = 0$.*

Proof. The $B_{\mathbf{N}}$ with $\mathbf{N}(c) = 0$ for $c \neq 1$ can be found by considering the generating function Ω and setting all x_c to zero other than x_1 . Then the calculation in Example 5.10 shows $B_{\mathbf{N}} = \binom{T(1)}{\mathbf{N}(1)}$, and binomial coefficients are a Z -basis of \mathcal{R} .

To show that \mathcal{R}_Γ is free over \mathcal{R} with the stated basis, we pass to the associated graded algebra with respect to the PBW filtration. Lemma 5.12 showed the leading order term of $B_{\mathbf{N}}$ is

$$\prod_{c \in \Gamma_*} \frac{T(c)^{\mathbf{N}(c)}}{\mathbf{N}(c)!} = \frac{T(1)^{\mathbf{N}(1)}}{\mathbf{N}(1)!} \prod_{c \neq 1} \frac{T(c)^{\mathbf{N}(c)}}{\mathbf{N}(c)!},$$

which we recognise as the leading term of $\binom{T(1)}{\mathbf{N}(1)} \in \mathcal{R}$ multiplied by the leading term of $B_{\mathbf{M}}$, where $\mathbf{M}(c) = \mathbf{N}(c)$ for $c \neq 1$ and $\mathbf{M}(1) = 0$. This shows that \mathcal{R}_Γ is spanned by the $B_{\mathbf{M}}$ over \mathcal{R} , and also shows the \mathcal{R} -linear independence of the $B_{\mathbf{M}}$.

DEFINITION 5.15. For any $\mathbf{N} \in Z_{>0}^{\Gamma_*}$ with $|\mathbf{N}| \triangleleft n$, we define $b_{\mathbf{N}} \in \mathcal{O}\Gamma \wr S_n$ as follows

$$b_{\mathbf{N}} = \frac{1}{(n - |\mathbf{N}|)! \prod_{c \in \Gamma_*} \mathbf{N}(c)!} \sum_{\sigma \in S_n} \sigma \left(1^{\otimes(n-|\mathbf{N}|)} \otimes \bigotimes_{c \in \Gamma_*} c^{\otimes \mathbf{N}(c)} \right) \sigma^{-1}.$$

The parenthesised tensor product is an element of $(\mathcal{O}\Gamma)^{\otimes n} = \mathcal{O}\Gamma^n \subseteq \mathcal{O}\Gamma \wr S_n$ (because of the averaging over S_n , the choice of order of the tensor factors does not matter).

LEMMA 5.16. The element $b_{\mathbf{N}}$ is actually contained in $Z(\mathcal{O}\Gamma \wr S_n)$.

Proof. Since $b_{\mathbf{N}}$ is defined by averaging over S_n , it commutes with $S_n \subseteq \Gamma \wr S_n$. As the element being averaged is a tensor product of central elements of $Z\Gamma$, $b_{\mathbf{N}}$ will commute with $\Gamma^n \subseteq \Gamma \wr S_n$. This shows that $b_{\mathbf{N}}$ is central. To see that $b_{\mathbf{N}}$ has integer coefficients, note that the conjugation action of the subgroup

$$S_{n-|\mathbf{N}|} \times \prod_{c \in \Gamma_*} S_{\mathbf{N}(c)} \subseteq S_n$$

is trivial. Any element of a fixed coset of this subgroup acts the same way, and the size of this coset cancels out the denominator.

LEMMA 5.17. We have the following expression for the generating function for the elements $b_{\mathbf{N}}$:

$$\sum_{|\mathbf{N}| \triangleleft n} b_{\mathbf{N}} x^{\mathbf{N}} = (1 + \sum_c cx_c)^{\otimes n}.$$

Proof. We observe that $b_{\mathbf{N}}$ is the sum of pure tensors with $\mathbf{N}(c)$ tensor factors set equal to c and all remaining tensor factors equal to 1. Here we distinguish tensor factors assigned $c = 1$ from unassigned tensor factors which also take the value 1. But the coefficient of $x^{\mathbf{N}}$ on the right hand side will be the sum of all pure tensors with $\mathbf{N}(c)$ factors equal to c .

THEOREM 5.18. There is a homomorphism $ev_n^{\mathcal{R}} : \mathcal{R}_\Gamma \rightarrow Z(\mathcal{O}\Gamma \wr S_n)$ defined by $ev_n^{\mathcal{R}}(B_{\mathbf{N}}) = b_{\mathbf{N}}$ if $|\mathbf{N}| \triangleleft n$, and $ev_n^{\mathcal{R}}(b_{\mathbf{N}}) = 0$ otherwise.

Proof. We apply $ev_n^{\mathcal{R}}$ to Ω coefficientwise. Using the generating function from Lemma 5.17,

$$\begin{aligned} ev_n^{\mathcal{R}}(\Omega(x_{c_1}, \dots, x_{c_l})) ev_n^{\mathcal{R}}(\Omega(y_{c_1}, \dots, y_{c_l})) &= (1 + \sum_c cx_c)^{\otimes n} (1 + \sum_c cy_c)^{\otimes n} \\ &= ((1 + \sum_c cx_c)(1 + \sum_c cy_c))^{\otimes n} \\ &= (1 + \sum_c cz_c)^{\otimes n} \\ &= ev_n^{\mathcal{R}}(\Omega(z_{c_1}, \dots, z_{c_l})) \\ &= ev_n^{\mathcal{R}}(\Omega(x_{c_1}, \dots, x_{c_l}) \Omega(y_{c_1}, \dots, y_{c_l})) \end{aligned}$$

where $z_{c_i} = x_{c_i} + y_{c_i} + \sum_{j,k} A_{j,k}^i x_{c_j} y_{c_k}$ as in Proposition 5.13. Considering the coefficients of monomials $x^{\mathbf{N}} y^{\mathbf{M}}$ shows $ev_n^{\mathcal{R}}(B_{\mathbf{N}}) ev_n^{\mathcal{R}}(B_{\mathbf{M}}) = ev_n^{\mathcal{R}}(B_{\mathbf{N}} B_{\mathbf{M}})$, i.e. that $ev_n^{\mathcal{R}}$ is a homomorphism.

The elements $B_{\mathbf{N}}$ can be thought of as interpolating the $b_{\mathbf{N}}$. To make this precise, we show that $ev_n^{\mathcal{R}}$ factors through FH_{Γ} .

COROLLARY 5.19. *We obtain a canonical homomorphism $\Psi : \mathcal{R}_{\Gamma} \rightarrow \text{FH}_{\Gamma}$ via*

$$\Psi(B_{\mathbf{N}}) = \binom{t - |\mathbf{N}| + \mathbf{N}(1)}{\mathbf{N}(1)} K_{\boldsymbol{\mu}},$$

where t is the polynomial variable of \mathcal{R} , $\boldsymbol{\mu}(c) = (1^{\mathbf{N}(c)})$ for $c \neq 1$, and $\boldsymbol{\mu}(1)$ is the empty partition. This homomorphism obeys $\Phi_n \circ \Psi = ev_n^{\mathcal{R}}$.

Proof. We first show that $\Phi_n \circ \Psi = ev_n^{\mathcal{R}}$. This amounts to checking that

$$b_{\mathbf{N}} = \binom{n - |\mathbf{N}| + \mathbf{N}(1)}{\mathbf{N}(1)} X_{\boldsymbol{\mu}}.$$

Each pure tensor in $b_{\mathbf{N}}$ is an element of generalised cycle type $\boldsymbol{\mu}$. However, such a pure tensor has $\mathbf{N}(1)$ assigned factors equal to 1 and $n - |\mathbf{N}|$ unassigned factors equal to 1. So each such pure tensor arises in $\binom{n - |\mathbf{N}| + \mathbf{N}(1)}{\mathbf{N}(1)}$ ways according to how the $\mathbf{N}(1)$ assigned factors are chosen. This accounts for the multiplicative factor. To check that Ψ respects multiplication, let $x, y \in \mathcal{R}_{\Gamma}$. Because $ev_n^{\mathcal{R}}$ and Φ_n are homomorphisms,

$$\begin{aligned} \Phi_n(\Psi(x)\Psi(y)) &= \Phi_n(\Psi(x))\Phi_n(\Psi(y)) \\ &= ev_n^{\mathcal{R}}(x)ev_n^{\mathcal{R}}(y) \\ &= ev_n^{\mathcal{R}}(xy) \\ &= \Phi_n(\Psi(xy)). \end{aligned}$$

We now use Proposition 5.5 to conclude that $\Psi(x)\Psi(y) = \Psi(xy)$.

The upshot is that we may now view FH_{Γ} as an \mathcal{R}_{Γ} -algebra, not merely an \mathcal{R} -algebra. But the fact that our ground ring now incorporates all conjugacy classes of Γ means that rather than considering partially reduced cycle types, we will have to reduce all parts of a multipartition.

DEFINITION 5.20. *If an element of $\Gamma \wr S_n$ has cycle type $\boldsymbol{\mu}$, we say it has fully-reduced cycle type $\boldsymbol{\nu}$, where for each $c \in \Gamma_*$, $\boldsymbol{\nu}(c)$ is obtained from $\boldsymbol{\mu}(c)$ by subtracting 1 from each part, and ignoring any resulting parts of size zero. We let $\hat{\boldsymbol{\mu}}$ be the multipartition obtained from $\boldsymbol{\mu}$ by adding 1 to each nonzero part of $\boldsymbol{\mu}(c)$ for each $c \neq 1$, and leaving $\boldsymbol{\mu}(1)$ unchanged.*

For an element of $\Gamma \wr S_n$, passing from the cycle type to the fully-reduced cycle type is generally not reversible, even if n is known. While we may recover the number of 1-cycles, their types may be arbitrary conjugacy classes. The notation has been set up so that $X_{\hat{\boldsymbol{\mu}}}$ is the sum of all elements of $\Gamma \wr S_n$ with all 1-cycles labelled by the trivial conjugacy class and having fully-reduced cycle type $\boldsymbol{\mu}$.

THEOREM 5.21. *As \mathcal{R}_{Γ} -module, FH_{Γ} is free with basis $K_{\hat{\boldsymbol{\mu}}}$ where $\boldsymbol{\mu} \in \mathcal{P}(\Gamma_*)$.*

Proof. By Proposition 5.14, \mathcal{R}_{Γ} is free over \mathcal{R} with basis $B_{\mathbf{M}}$ indexed by \mathbf{M} with $\mathbf{M}(1) = 0$. So it is enough to show that $K_{\hat{\boldsymbol{\mu}}} B_{\mathbf{M}}$ is a \mathcal{R} -basis of FH_{Γ} . To accomplish this, we introduce a filtration of FH_{Γ} by \mathcal{R} -modules by placing $K_{\boldsymbol{\mu}}$ in filtration degree $|\boldsymbol{\mu}| + l(\boldsymbol{\mu}(1))$. We now show that this respects multiplication, by passing to $Z(\Gamma \wr S_n)$.

Let us say that $(\mathbf{g}, \sigma) \in \Gamma \wr S_n$ affects $i \in \{1, \dots, n\}$ if either $\sigma(i) \neq i$, or $\mathbf{g}_i \neq 1$. The number of elements of $\{1, \dots, n\}$ that are affected by an element of partially-reduced

cycle type μ is precisely $|\mu| + l(\mu(1))$. Consider two elements $(\mathbf{g}, \sigma), (\mathbf{h}, \rho) \in \Gamma \wr S_n$, and suppose that their product $(\mathbf{g}\sigma(\mathbf{h}), \sigma\rho)$ affects i . Then either $\rho(\sigma(i)) \neq i$, or $\mathbf{g}_i\sigma(\mathbf{h})_i \neq 1$. If $\rho(\sigma(i)) \neq i$, then either $\sigma(i) \neq i$, or $\rho(i) \neq i$, which implies that one of $(\mathbf{g}, \sigma), (\mathbf{h}, \rho)$ affects i . If instead $\mathbf{g}_i\sigma(\mathbf{h})_i \neq 1$, either $\mathbf{g}_i \neq 1$ or $\sigma(\mathbf{h})_i \neq 1$. In the first of these cases, (\mathbf{g}, σ) affects i . In the second case, either $\sigma(i) \neq i$, or $\mathbf{h}_i \neq 1$. Hence we conclude that if $(\mathbf{g}\sigma(\mathbf{h}), \sigma\rho)$ affects i , then one of (\mathbf{g}, σ) and (\mathbf{h}, ρ) affects i . This implies that the set of elements in $\{1, \dots, n\}$ affected by the product of two group elements is a subset of the union of the elements affected by each factor. This proves that we have a filtration, and shows that only products of elements affecting disjoint subsets of $\{1, \dots, n\}$ contribute to the leading term in the associated graded ring.

To compute $K_{\hat{\mu}}B_{\mathbf{M}}$, we again pass to $Z\Gamma \wr S_n$. This gives $X_{\hat{\mu}}b_{\mathbf{M}}$. The leading order term arises from $X_{\hat{\mu}}$ and $b_{\mathbf{M}}$ affecting disjoint subsets of $\{1, \dots, n\}$. In that case, we obtain X_{ν} , where $\nu(c) = \hat{\mu}(c) \cup (1^{\mathbf{M}(c)})$ for $c \in \Gamma_*$. Any multipartition ν arises from exactly one pair $(\hat{\mu}, \mathbf{M})$ in this way. In particular μ is the fully-reduced cycle type corresponding to the partially-reduced cycle type ν , while \mathbf{M} records how many parts of ν of size 1 have a given label in $\Gamma_* \setminus \{1\}$. We conclude that $K_{\hat{\mu}}B_{\mathbf{M}}$ equals K_{ν} plus lower order terms. In particular the $K_{\hat{\mu}}B_{\mathbf{M}}$ form an \mathcal{R} -basis of FH_{Γ} and the theorem follows.

6. ISOMORPHISM WITH Γ_* -WEIGHTED SYMMETRIC FUNCTIONS

This section is dedicated to proving that FH_{Γ} is isomorphic to $\mathcal{R}_{\Gamma} \otimes \Lambda(\Gamma_*)$ via Jucys–Murphy evaluation, analogously to Theorem 3.8.

We introduce two filtrations. Let us say that $\sigma \in S_n$ moves an element $m \in \{1, \dots, n\}$ if $\sigma(m) \neq m$ (i.e. m is not a fixed point of σ). The first filtration is obtained by placing an element $(\mathbf{g}, \sigma) \in \Gamma \wr S_n$ in degree i , where i is the number of elements of $\{1, \dots, n\}$ which are moved by σ .

DEFINITION 6.1. *The moving filtration, \mathcal{F}_{mov}^i , is the family of \mathcal{R}_{Γ} -submodules of FH_{Γ} spanned by $K_{\hat{\mu}}$ with $|\mu| + l(\mu) \leq i$.*

PROPOSITION 6.2. *The moving filtration is an algebra filtration. In the associated graded algebra, we have the following equation:*

$$K_{\hat{\mu}}K_{\hat{\nu}} = \left(\prod_{c \in \Gamma_*} \prod_{i > 1} \frac{m_i(\lambda(c))!}{m_i(\mu(c))!m_i(\nu(c))!} \right) K_{\hat{\lambda}},$$

where $m_i(\lambda(c)) = m_i(\mu(c)) + m_i(\nu(c))$.

Proof. If $\sigma \in S_n$ has reduced cycle type μ , the number of elements moved by σ is $|\mu| + l(\mu)$. Similarly if $(\mathbf{g}, \sigma) \in \Gamma \wr S_n$ has fully-reduced cycle type μ , then the number of elements moved by σ is $|\mu| + l(\mu)$. Note that this corresponds to the condition for membership in \mathcal{F}_{mov}^i , where $i = |\mu| + l(\mu)$.

Fix μ and ν with $|\mu| + l(\mu) = i$ and $|\nu| + l(\nu) = j$. Consider $K_{\hat{\mu}}K_{\hat{\nu}}$. To compute this product, we pass to $Z(Z\Gamma \wr S_n)$ for large n . A term in $X_{\hat{\mu}}$ is of the form (\mathbf{g}, σ) , where σ moves i elements. Similarly a term in $X_{\hat{\nu}}$ is of the form (\mathbf{h}, ρ) , where ρ moves j elements. Now, $(\mathbf{g}, \sigma) \cdot (\mathbf{h}, \rho) = (\mathbf{g}\sigma(\mathbf{h}), \sigma\rho)$. The permutation $\sigma\rho$ is obtained by moving j elements, and then moving i elements. This means that $\sigma\rho$ moves at most $i + j$ elements, with equality if and only if the elements moved by σ and ρ are disjoint. This proves the assertion that \mathcal{F}_{mov}^i defines a filtration.

If σ and ρ move disjoint elements, then $\sigma(\mathbf{h}) = \mathbf{h}$ because the m -th factor in \mathbf{h} is equal to 1 if m is not moved by ρ (so in particular if m is moved by σ). Hence the cycles (of size larger than 1) of $(\mathbf{g}, \sigma) \cdot (\mathbf{h}, \rho)$ are the constituent cycles of (\mathbf{g}, σ) and (\mathbf{h}, ρ) . This shows that in the associated graded algebra, $K_{\hat{\mu}}K_{\hat{\nu}}$ is equal to a multiple

of K_{λ} . The multiple is equal to the number of ways to split the cycles of λ among μ and ν .

The second filtration is obtained by placing an element $(\mathbf{g}, \sigma) \in \Gamma \wr S_n$ in degree i , where i is the smallest number of factors required to express σ as a product of transpositions.

DEFINITION 6.3. *The transposition filtration, \mathcal{F}_{tsp}^i , is the family of \mathcal{R}_Γ -submodules of FH_Γ spanned by $K_{\hat{\mu}}$ with $|\mu| \subset i$.*

PROPOSITION 6.4. *The transposition filtration is an algebra filtration.*

Proof. Similarly to the other filtration, we pass to $Z(Z\Gamma \wr S_n)$ for n sufficiently large. Fix μ and ν with $|\mu| = i$ and $|\nu| = j$. A term in $X_{\hat{\mu}}$ is of the form (\mathbf{g}, σ) , where σ may be written as the product of i transpositions. Similarly a term in $X_{\hat{\nu}}$ is of the form (\mathbf{h}, ρ) , where ρ is a product of j transpositions. Then, $(\mathbf{g}, \sigma) \cdot (\mathbf{h}, \rho) = (\mathbf{g}\sigma(\mathbf{h}), \sigma\rho)$. The permutation $\sigma\rho$ may be written as a product of $i + j$ transpositions (although possibly fewer). We conclude that \mathcal{F}_{tsp}^i defines a filtration.

REMARK 6.5. Using [6, Lemma 3.2] and [24, Proposition 2.9], one can show that the structure constants of the $K_{\hat{\mu}}$ elements in the associated graded algebra are integers rather than arbitrary elements of \mathcal{R}_Γ . We will not need this fact.

6.1. JUCYS–MURPHY ELEMENTS. Analogously to the JM elements associated to symmetric groups, Pushkarev [18] and Wang [25] independently introduced the following elements of $Z\Gamma \wr S_n$:

$$L_j = \sum_{i < j} \sum_{g \in \Gamma} g^{(i)}(g^{-1})^{(j)}(i, j).$$

In the case where Γ is the trivial group, we recover the JM elements for symmetric groups. However, these elements are not sufficient for our purposes, and we must introduce further generalisations.

DEFINITION 6.6. *For $c \in \Gamma_*$ and $j \subset n$, the wreath-product Jucys–Murphy elements are*

$$L_j(c) = \sum_{i < j} \sum_{\substack{g_1, g_2 \in \Gamma \\ g_2 g_1 \in c}} g_1^{(i)} g_2^{(j)}(i, j),$$

which are elements of $Z\Gamma \wr S_n$. We extend this notation by linearity in c : if $c = \sum_i n_i c_i$ for some $n_i \in \mathbb{Z}$, we let

$$L_j(c) = \sum_{i=1}^l n_i L_j(c_i).$$

In the case where $c = 1$, the condition $g_2 g_1 \in c$ becomes $g_2 = g_1^{-1}$, so $L_j(1)$ recovers the definition of Pushkarev and Wang.

PROPOSITION 6.7. *Let $c \in Z\Gamma_*$. We have*

$$L_j(c) = c^{(j)} L_j(1) = L_j(1) c^{(j)}.$$

Proof. By linearity, it suffices to consider $c \in \Gamma_*$. To show the first equality, it suffices that for $i < j$,

$$c^{(j)} \sum_{\substack{g_1, g_2 \in \Gamma \\ g_2 g_1 = 1}} g_1^{(i)} g_2^{(j)} = \sum_{\substack{h_1, h_2 \in \Gamma \\ h_2 h_1 \in c}} h_1^{(i)} h_2^{(j)},$$

as we may right-multiply by (i, j) and sum over i less than j . We may set $g_2 = g_1^{-1}$ and use the fact that $c^{(j)}$ commutes with $g_1^{(i)}$ and $g_2^{(j)}$. The left hand side becomes

$$\sum_{g_1 \in \Gamma} g_1^{(i)} (g_1^{-1})^{(j)} c^{(j)} = \sum_{g_1 \in \Gamma, k \in c} g_1^{(i)} (g_1^{-1}k)^{(j)},$$

which agrees with the right hand side upon identifying $h_1 = g_1$ and $h_2 = g_1^{-1}k$. The second equality is similar, except that moving $c^{(j)}$ past the transposition (i, j) turns it into $c^{(i)}$, which means that the manipulations take place in the i -index, rather than the j -index.

DEFINITION 6.8. For $c \in \Gamma_*$, let

$$M_c^{(i, i+1)} = \sum_{\substack{g, h \in \Gamma \\ hg \in c}} g^{(i)} h^{(i+1)}.$$

The following lemmas have routine proofs, so we omit them.

LEMMA 6.9. We have $M_c^{(i, i+1)} = c^{(i)} M_1^{(i, i+1)} = c^{(i+1)} M_1^{(i, i+1)}$, and $M_c^{(i, i+1)}$ commutes with the transposition $s_i = (i, i + 1)$.

LEMMA 6.10 ([18, Proposition 5]). We have that the $L_j(1)$ ($j = 1, \dots, n$) commute with each other. The $L_j(1)$ commute with Γ^n . Furthermore

$$s_i L_i(1) s_i + M_1^{(i, i+1)} s_i = L_{i+1}(1).$$

The following lemma is easily proven by induction.

LEMMA 6.11. We have the two identities

$$s_i L_i(1)^r = L_{i+1}(1)^r s_i - \sum_{p=0}^{r-1} L_{i+1}(1)^p M_1^{(i, i+1)} L_i(1)^{r-1-p}$$

and

$$s_i L_{i+1}(1)^r = L_i(1)^r s_i + \sum_{p=0}^{r-1} L_i(1)^p M_1^{(i, i+1)} L_{i+1}(1)^{r-1-p}.$$

PROPOSITION 6.12. Suppose $c_a, c_b \in \Gamma_*$. The following elements both commute with s_i :

$$L_i(1)^r c_a^{(i)} c_b^{(i+1)} + L_{i+1}(1)^r c_b^{(i)} c_a^{(i+1)} \text{ and } L_i(1)^r L_{i+1}(1)^r.$$

Proof. For the first part, we add the first equation in Lemma 6.11 right-multiplied by $c_a^{(i)} c_b^{(i+1)}$ to the second equation right-multiplied by $c_b^{(i)} c_a^{(i+1)}$. Because $L_i(1)$ and $L_{i+1}(1)$ commute with Γ^n , they commute with $M_1^{(i, i+1)}$. So since $M_1^{(i, i+1)} c_a^{(i)} c_b^{(i+1)} = M_1^{(i, i+1)} c_b^{(i)} c_a^{(i+1)}$, the sums over the variable p cancel out, leaving the required identity. For the second part it suffices to consider the case $r = 1$ as the general case is recovered by raising $L_i(1)L_{i+1}(1)$ to a suitable power. Then applying Lemma 6.11 (specifically, the second equation, with $r = 1$), we have

$$\begin{aligned} s_i L_i(1) L_{i+1}(1) &= (L_{i+1}(1) s_i - M_1^{(i, i+1)}) L_{i+1}(1) \\ &= L_{i+1}(1) (L_i(1) s_i + M_1^{(i, i+1)}) - M_1^{(i, i+1)} L_{i+1}(1) \\ &= L_{i+1}(1) L_i(1) s_i. \end{aligned}$$

Since $L_i(1)$ and $L_{i+1}(1)$ commute, $s_i L_i(1) L_{i+1}(1) = L_i(1) L_{i+1}(1) s_i$ and we are done.

PROPOSITION 6.13. The wreath-product JM elements commute with each other: $L_{j_1}(c_1) L_{j_2}(c_2) = L_{j_2}(c_2) L_{j_1}(c_1)$ for any $c_1, c_2 \in \Gamma_*$ and $j_1, j_2 \in \{1, \dots, n\}$.

Proof. We have that the elements $L_{j_1}(c_1) = L_{j_1}(1)c_1^{(j_1)}$ and $L_{j_2}(c_2) = L_{j_2}(1)c_2^{(j_2)}$ and $L_{j_1}(1), c_1^{(j_1)}, L_{j_2}(1), c_2^{(j_2)}$ commute pairwise.

Because the wreath-product JM elements commute, it makes sense to evaluate a polynomial in them.

PROPOSITION 6.14. *There is a ring homomorphism*

$$ev_n^Q : Q^{\otimes n} \rightarrow Z\Gamma \wr S_n,$$

defined by

$$ev_n^Q(x_d^r(c)) = L_d(1)^r c^{(d)}.$$

Moreover, the image of $\Lambda_n(\Gamma_*)$ (the S_n -invariants of $Q^{\otimes n}$) is contained in the centre of $Z\Gamma \wr S_n$.

Proof. Since the wreath-product JM elements commute pairwise, it suffices to show that ev_n^Q is well-defined on each tensor factor of $Q^{\otimes n}$. The d -th factor of Q has a basis $x_d^r(c)$ for $r > 0$ and $c \in \Gamma_*$ (where we require $c = 1$ if $r = 0$). This basis obeys the multiplication rule mentioned at the start of Section 4 (where we expand the multiplication in $Z\Gamma_*$ in terms of the structure tensor $A_{i,j}^k$)

$$x_d^r(c_i)x_d^s(c_j) = x_d^{r+s}(c_i c_j) = \sum_k A_{i,j}^k x_d^{r+s}(c_k).$$

Correspondingly, using Proposition 6.13,

$$\begin{aligned} ev_n^Q(x_d^r(c_i))ev_n^Q(x_d^s(c_j)) &= L_d(1)^r c_i^{(d)} L_d(1)^s c_j^{(d)} \\ &= L_d(1)^{r+s} (c_i c_j)^{(d)} \\ &= L_d(1)^{r+s} \sum_k A_{i,j}^k c_k^{(d)} \\ &= \sum_k A_{i,j}^k ev_n^Q(x_d^{r+s}(c_k)). \end{aligned}$$

We conclude that ev_n^Q is well defined.

Now suppose that $P(x_1, \dots, x_n) \in (Q^{\otimes n})^{S_n}$. To show $ev_n^Q(P)$ is central, it suffices to show that it commutes with a generating set of $\Gamma \wr S_n$. In particular, we may take Γ^n together with the adjacent transpositions $s_i = (i, i + 1)$ for $i \in \{1, \dots, n - 1\}$. Since $L_j(1)$ commutes with Γ^n , the same is true for $L_j(c) = c^{(j)}L_j(1)$, from which it follows that the entire image of ev_n^Q commutes with Γ^n . Now we show that $ev_n^Q(P)$ commutes with s_i . Since $P \in (Q^{\otimes n})^{S_n}$, P is in particular an element of

$$Q^{\otimes(i-1)} \otimes (Q \otimes Q)^{S_2} \otimes Q^{\otimes(n-i-1)},$$

and $(Q \otimes Q)^{S_2}$ is spanned by elements of the form $x_i^r(c)x_{i+1}^r(c)$ and $x_i^r(c_a)x_{i+1}^s(c_b) + x_i^s(c_b)x_{i+1}^r(c_a)$. So it is enough that applying ev_n to these elements gives something that commutes with s_i . In the first case,

$$ev_n^Q(x_i^r(c)x_{i+1}^r(c)) = L_i(1)^r c^{(i)} L_{i+1}(1)^r c^{(i+1)},$$

which commutes with s_i because $L_i(1)L_{i+1}(1)$ and $c^{(i)}c^{(i+1)}$ do. For the second case, without loss of generality, we assume $s \subsetneq r$. Now,

$$\begin{aligned} ev_n^Q(x_i^r(c_a)x_{i+1}^s(c_b) + x_i^s(c_b)x_{i+1}^r(c_a)) &= L_i(1)^r c_a^{(i)} L_{i+1}(1)^s c_b^{(i+1)} + L_i(1)^r c_b^{(i)} L_{i+1}(1)^s c_a^{(i+1)} \\ &= L_i(1)^s L_{i+1}(1)^s (L_i(1)^{r-s} c_a^{(i)} c_b^{(i+1)} + L_{i+1}(1)^{r-s} c_b^{(i)} c_a^{(i+1)}) \end{aligned}$$

is a product of elements that commute with s_i from Proposition 6.12.

DEFINITION 6.15. *Let $ev_n^\Lambda : \Lambda(\Gamma_*) \rightarrow Z(Z\Gamma \wr S_n)$ be the composition of the canonical map $\Lambda(\Gamma_*) \rightarrow \Lambda_n(\Gamma_*) \subseteq Q^{\otimes n}$ with $ev_n^Q : Q^{\otimes n} \rightarrow Z(Z\Gamma \wr S_n)$.*

Finally, we combine our evaluation maps into a single map.

DEFINITION 6.16. For each $n \in \mathbb{Z}_{>0}$ there is a ring homomorphism

$$ev_n : \mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*) \rightarrow Z(Z\Gamma \wr S_n)$$

defined as

$$ev_n(a \otimes f) = ev_n^{\mathcal{R}}(a)ev_n^\Lambda(f),$$

using the maps $ev_n^{\mathcal{R}}$ and ev_n^Λ from Theorem 5.18 and Definition 6.15, respectively.

THEOREM 6.17. There is a unique homomorphism $\Psi : \mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*) \rightarrow \text{FH}_\Gamma$ extending the map in Corollary 5.19 such that the following diagram commutes.

$$\begin{array}{ccc} \mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*) & \xrightarrow{\Psi} & \text{FH}_\Gamma \\ & \searrow ev_n & \downarrow \Phi_n \\ & & Z(ZS_n) \end{array}$$

Proof. Corollary 5.19 constructs Ψ on \mathcal{R}_Γ , so it remains to construct Ψ on $\Lambda(\Gamma_*)$. To do this, we pass to complex coefficients, and then show that the map is defined integrally. Proposition 4.6 shows

$$\mathbb{C} \otimes \Lambda(\Gamma_*) = \mathbb{C} \otimes \Lambda^{|\Gamma^*|}$$

by changing basis in $\mathbb{C}\Gamma_*$ from conjugacy-class sums to primitive central idempotents e_χ . To compute ev_n on the tensor factor of Λ corresponding to $\chi \in \Gamma^*$, we must evaluate symmetric functions in the variables

$$L_j(e_\chi) = L_j(1)e_\chi^{(j)}.$$

Since Λ is freely generated by the elementary symmetric functions e_r , it suffices to consider the evaluation of e_r at the $L_j(e_\chi)$. Momentarily ignoring the Γ^n parts in $L_j(e_\chi)$, we see that the transpositions involved in $L_j(e_\chi)$ are the same as in the JM elements L_j for the symmetric group. So the S_n parts are the same as in Theorem 3.12. Lemma 2.6 tells us that the labels of intersecting cycles multiply. But each label is the idempotent e_χ . So we get a sum of elements whose reduced cycle type has size r , and all of whose cycles are labelled by e_χ (which is a linear combination of the usual labels in Γ_*). We could write this

$$e_r(L_1(e_\chi), \dots, L_n(e_\chi)) = \sum_{|\mu|=r} X_{\mu_\chi}^{irr},$$

where the meaning of $X_{\mu_\chi}^{irr}$ is analogous to m_{λ_χ} from Proposition 4.6. Changing bases of $\mathbb{C}\Gamma_*$ back from e_χ to conjugacy-class sums changes the labels back to conjugacy classes. This is a linear operation that is independent of n . Hence we obtain some element

$$e_r(L_1(e_\chi), \dots, L_n(e_\chi)) = \sum_{|\nu|=r} R_{r,\nu}^\chi X_{\nu},$$

where $R_{r,\nu}^\chi$ is independent of n , and the multipartition ν is indexed by Γ_* (the sizes of the cycles of the elements are unchanged by changing the type so we still sum over multipartitions of size r). Thus for $e_r \in \Lambda(\chi)$, we may take

$$\Psi(e_r) = \sum_{|\nu|=r} R_{r,\nu}^\chi K_{\nu},$$

which guarantees that $ev_n = \Phi_n \circ \Psi$.

To see that the image of Ψ on $\Lambda(\Gamma_*)$ is contained in FH_Γ rather than $\mathbb{C} \otimes \text{FH}_\Gamma$, note that we can tell whether an element $f \in \mathbb{C} \otimes \text{FH}_\Gamma$ is contained in FH_Γ by applying Φ_n for sufficiently large n and checking whether the coefficients of the conjugacy class

sums X_{μ} are integers. This would imply that the coefficient of K_{μ} in f , which is an element of $\mathbb{C} \otimes \mathcal{R}$, is integer-valued, i.e. an element of \mathcal{R} . On the other hand, we already know the image of ev_n on $\Lambda(\Gamma_*)$ is contained in $Z(Z\Gamma \wr S_n)$ and hence integral.

PROPOSITION 6.18. *Recall that FH_{Γ} is equipped with two filtrations (the moving and transposition filtrations). If we take the leading order term of $\Psi(m_{\mu})$ with respect to the transposition filtration, and then the leading order term with respect to the moving filtration, we get $X_{\hat{\mu}}$.*

Proof. Consider $ev_n(m_{\mu})$. In the notation of Lemmas 2.5 and 2.6,

$$L_j(c) = \sum_{i < j} X_{(i,j)}(c).$$

Analogously to the proof of Proposition 3.11, suppose that σ is a cycle containing the index j . Multiplying $X_{\sigma}(1)$ by $X_{(i,j)}(c)$ inserts i into the cycle if i is not already in σ , or removes it if it is. The size with respect to the transposition filtration is increased by 1 in the first case, and reduced by 1 in the second case. To find the leading order term of $L_j(1)^r$, we view it as a sum of products of $X_{(i_k,j)}(1)$ ($k = 1, \dots, r$) where each i_k is less than j . The maximal degree with respect to the transposition filtration is achieved when the i_k are distinct. In that case, Lemmas 2.5 and 2.6 imply that the product of the $X_{(i_k,j)}(1)$ is $X_{\sigma}(1)$, where $\sigma = (i_r, i_{r-1}, \dots, i_1, j)$ is an $(r + 1)$ -cycle, and furthermore the leading term of $L_j(1)^r c^{(j)}$ in the transposition filtration is the sum of $X_{\sigma}(c)$ across all $(r + 1)$ -cycles σ whose largest element is j . Now, $ev_n(m_{\mu})$ is a sum of products of $L_j(1)^r c^{(j)}$ where the pair of parameters (r, c) occurs $m_r(\mu(c))$ times. Each such term gives us an $(r + 1)$ -cycle of type c .

We maximise the degree in the moving filtration when all these cycles are disjoint. In that case, the result is an element of fully-reduced cycle type μ . Every such element arises exactly once, because we may reconstruct the monomial which gave rise to it: an $(r + 1)$ cycle whose largest element is j and has type c must have come from $ev_n(x_j^r(c))$.

THEOREM 6.19. *The map Ψ from Theorem 6.17 is an isomorphism and in particular $\text{FH}_{\Gamma} = \mathcal{R}_{\Gamma} \otimes \Lambda(\Gamma_*)$.*

Proof. We see that up to lower order terms, Ψ sends a \mathcal{R}_{Γ} -basis of $\mathcal{R}_{\Gamma} \otimes \Lambda(\Gamma_*)$, namely m_{λ} , to a \mathcal{R}_{Γ} -basis of FH_{Γ} , namely $K_{\hat{\lambda}}$. This shows that the homomorphism Ψ is a bijection.

7. APPLICATIONS AND FURTHER DIRECTIONS

We prove a wreath-product version of Nakayama’s Conjecture. The first step is to understand the action of the wreath-product JM elements on the irreducible (characteristic zero) representations V^{λ} of $\Gamma \wr S_n$.

Analogously to the GZ basis for irreducible representations of symmetric groups, there is a similar construction for $\Gamma \wr S_n$. It is not a basis, but rather a decomposition into subspaces. What is important for us is that the wreath-product JM elements act by scalars on the GZ subspaces of an irreducible representation of $\Gamma \wr S_n$.

DEFINITION 7.1 ([15, Section 1]). *Let λ be a Γ^* -indexed multipartition of size n . A Young Γ -tableau T of shape λ is a labelling of the boxes of all partitions in λ with numbers. Suppose that the labelling uses the numbers $\{1, \dots, n\}$ (with each being used exactly once), such that for each partition on λ , the labels increase from top to bottom in a column and increase from left to right in a row; in this case we say T is standard. If T is a standard Young Γ -tableau, then we write $\chi_{T(i)}$ for the irreducible*

representation $\chi \in \Gamma^*$ such that $T(\chi)$ contains the box with label i , and $c_T(i)$ for the content of the box containing i in the partition corresponding to $\chi_{T(i)}$.

EXAMPLE 7.2. If $\Gamma = C_2$, the following pair of partitions, corresponding to the trivial and sign representations of C_2 , define a standard Young C_2 -tableau.

1	3	6
7		

2	5
4	

In particular, the constituent tableaux of a standard Young Γ -tableau need not be standard Young tableaux because the labels of the tableau corresponding to $\chi \in \Gamma^*$ are allowed to come from the set $\{1, \dots, |\lambda|\}$, which may be larger than $\{1, \dots, |\lambda(\chi)|\}$.

DEFINITION 7.3. For $m \leq n$, let $H_{m,n}$ be the subgroup of $\Gamma \wr S_n$ consisting of elements of the form (\mathbf{g}, σ) where $\mathbf{g} \in \Gamma^n$ is arbitrary and $\sigma \in S_m$, where S_m is viewed as the subgroup of S_n fixing all numbers larger than m . Equivalently, $H_{m,n} = \Gamma^n \circ S_m$ with the same embedding of S_m into S_n .

PROPOSITION 7.4 ([15, Section 4]). The restriction of an irreducible representation of $H_{m,n}$ to $H_{m-1,n}$ is multiplicity free (i.e. has no repeated irreducible summands).

The significance of this multiplicity-free statement is that the decomposition of the restricted representation into irreducibles is unique (since the irreducible summands coincide with isotypic components).

DEFINITION 7.5 ([15, Section 3]). The GZ subspaces of an irreducible representation V of $H_{n,n} = \Gamma \wr S_n$ are defined inductively as follows. An irreducible representation of $H_{m,n}$ splits uniquely as a sum of irreducible representations of $H_{m-1,n}$, each of which may be again restricted. Iterating this for $m = n, n - 1, \dots, 0$, we are left with a canonical decomposition of V into irreducible representations V_T of $H_{0,n} = \Gamma^n$.

PROPOSITION 7.6 ([15, Section 4]). The GZ subspaces V_T of the irreducible representation V^λ are indexed by standard Young Γ -tableaux T of shape λ . We have

$$V^\lambda = \bigoplus_T V_T,$$

where the sum is across standard Young Γ -tableaux of shape λ . As a representation of Γ^n ,

$$V_T = \chi_{T(1)} \otimes \cdots \otimes \chi_{T(n)}.$$

EXAMPLE 7.7. For the standard Young C_2 -tableau T in Example 7.2, the corresponding GZ subspace is isomorphic to

$$C_+ \otimes C_- \otimes C_+ \otimes C_- \otimes C_- \otimes C_+ \otimes C_+,$$

where C_+ is the trivial representation and C_- is the sign representation.

PROPOSITION 7.8. The wreath-product JM element $L_j(c)$ acts on the GZ subspace V_T via scalar multiplication by

$$\frac{|\Gamma|}{\dim(\chi_{T(j)})} c_T(i) \omega_c^{\chi_{T(j)}}.$$

Proof. Theorem 6.5 of [15] asserts that $L_j(1)$ acts by the scalar $\frac{|\Gamma|}{\dim(\chi_{T(j)})} c_T(i)$. But $L_j(c) = L_j(1)c^{(j)}$, and $c^{(j)}$ acts on $\chi_{T(1)} \otimes \cdots \otimes \chi_{T(n)}$ as multiplication by $\omega_c^{\chi_{T(j)}}$.

THEOREM 7.9. *Suppose that p is a prime number, and suppose that B_i are the p -blocks of Γ (the subsets of Γ^* with a given central character in characteristic p). Let λ and μ be two Γ^* -indexed multipartitions of size n . Then the irreducible representations V^λ, V^μ of $\Gamma \wr S_n$ are in the same p -block if and only if both of the following conditions hold:*

- (1) For all p -blocks B_i of Γ ,

$$\sum_{\chi \in B_i} |\lambda(\chi)| = \sum_{\chi \in B_i} |\mu(\chi)|.$$

- (2) For each B_i containing only one irreducible χ , the partitions $\lambda(\chi)$ and $\mu(\chi)$ have the same p -core.

Proof. Analogously to the proof of Nakayama’s conjecture (Theorem 3.16), we fix a p -modular system (K, \mathcal{O}, k) , and let π be a uniformiser of \mathcal{O} . We determine when two central characters are equal modulo (π) , where we use the surjection $ev_n : \mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*) \rightarrow Z(Z\Gamma \wr S_n)$ to compute the action of the centre. The action of $\mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*)$ is determined by the individual actions of \mathcal{R}_Γ and $\Lambda(\Gamma_*)$.

First we consider the action of \mathcal{R}_Γ , whose image in $Z(Z\Gamma \wr S_n)$ is contained in $Z(Z\Gamma^n)$ (see Theorem 5.18). So to compute the action by scalars, we may restrict the representation V^λ from $\Gamma \wr S_n$ to Γ^n . This gives the sum of the GZ subspaces V_T , each of which is of the form

$$\chi_{T(1)} \otimes \cdots \otimes \chi_{T(n)},$$

where the number times a given irreducible $\chi \in \Gamma^*$ appears is $|\lambda(\chi)|$. We compute the action of \mathcal{R}_Γ on a GZ subspace V_T . By Lemma 5.17,

$$\sum_{\mathbf{N}} ev_n(B_{\mathbf{N}})x^{\mathbf{N}} = (1 + \sum_c cx_c)^{\otimes n}.$$

Now, the action of $(1 + \sum_c cx_c)$ on χ is by multiplication by

$$(1 + \sum_c \omega_c^\chi x_c),$$

where ω_c^χ is the central character of Γ corresponding to the action of c on χ . Hence the action of $\sum_{\mathbf{N}} ev_n(B_{\mathbf{N}})x^{\mathbf{N}}$ is

$$\prod_{\chi \in \Gamma^*} (1 + \sum_c \omega_c^\chi x_c)^{|\lambda(\chi)|}.$$

This result remains true after modular reduction (i.e. passing to $k = \mathcal{O}/(\pi)$), although this has the effect of making central characters within a single p -block coincide. Since the $B_{\mathbf{N}}$ are a basis of \mathcal{R}_Γ , the action of \mathcal{R}_Γ is determined by this generating function which is actually a polynomial. Polynomials over a field form a unique factorisation domain, so two such polynomials agree if and only if they have the same factors (note that all our factors are normalised to have constant term 1, so two factors are associates if and only if they are equal). But two factors for χ_1 and χ_2 , coincide precisely when

$$(1 + \sum_c \omega_c^{\chi_1} x_c) = (1 + \sum_c \omega_c^{\chi_2} x_c),$$

i.e. when $\omega_c^{\chi_1} = \omega_c^{\chi_2}$, which is to say that the central characters of χ_1 and χ_2 agree. This in turn happens when χ_1 and χ_2 are in the same p -block. By factoring the generating function we determine the multiplicity of the χ from each p -block of Γ . We conclude that the \mathcal{R}_Γ acts by the same scalars (in k) on V^λ and V^μ if and only if condition (1) holds.

To understand the action of $\Lambda(\Gamma_*)$, we again consider a GZ subspace V_T . By Proposition 7.8, the wreath-product JM element $L_j(c)$ acts on V_T by the scalar

$$\frac{|\Gamma|}{\dim(\chi_{T(j)})} c_T(j) \omega_c^{\chi_{T(j)}}.$$

If $\frac{|\Gamma|}{\dim(\chi_{T(j)})}$ is a multiple of p , then this scalar is zero in k . Now suppose that conditions (1) and (2) are both satisfied for λ and μ . For each $\chi \in \Gamma_*$ and $c \in \Gamma_*$, the multiset of scalars

$$\frac{|\Gamma|}{\dim(\chi)} c_T(j) \omega_c^\chi$$

for j such that $\chi_{T(j)} = \chi$ coincides for T of shape λ and μ . To check this, we consider whether $\frac{|\Gamma|}{\dim(\chi)}$ is a multiple of p . If it is a multiple of p , then all these scalars are zero. If $\frac{|\Gamma|}{\dim(\chi)}$ is not a multiple of p , then the contents $c_T(j)$ coincide because $\lambda(\chi)$ and $\mu(\chi)$ have the same p -core. As a result, any element of $\Lambda(\Gamma_*)$ acts by the same scalar modulo (π) on V^λ and V^μ . We conclude that the “if” implication of the theorem holds.

For χ such that $|G|/\dim(\chi)$ is not divisible by p , the idempotent

$$e_\chi = \frac{\dim(\chi)}{|G|} \sum_{g \in \Gamma} \chi(g) g^{-1}$$

is well defined in k . We may therefore consider the elements $L_j(e_\chi)$ in $k\Gamma \wr S_n$. By Proposition 7.8, the action of $L_j(e_\chi)$ on V_T is

$$\frac{|\Gamma|}{\dim(\chi)} c_T(j)$$

if $\chi_{T(j)} = \chi$, and zero otherwise. This is because χ is in a p -block by itself (see Proposition 2.11), and e_χ acts by zero on irreducibles in other p -blocks, i.e. on all other irreducibles.

Just as in the proof of Theorem 6.17, the elementary symmetric functions e_r evaluated in $L_j(e_\chi)$,

$$e_r(L_1(e_\chi), \dots, L_n(e_\chi)),$$

are well defined elements of $k \otimes \Lambda(\Gamma_*)$. So the action of $\Lambda(\Gamma_*)$ on V_T determines the action of the generating function

$$\sum_{r>0} t^r e_r(L_1(e_\chi), \dots, L_n(e_\chi)) = \prod_{j=1}^n (1 + t L_j(e_\chi)).$$

But we have determined the action of $L_j(e_\chi)$, so we know that this generating function acts by

$$\prod_j (1 + t \frac{|\Gamma|}{\dim(\chi)} c_T(j))$$

where the product is only over those j such that $\chi_{T(j)} = \chi$. Appealing to the unique factorisation of polynomials over k , the action of $\Lambda(\Gamma_*)$ determines the multiset of scalars $\frac{|\Gamma|}{\dim(\chi)} c_T(j)$ (viewed as elements of k). Now, $\frac{|\Gamma|}{\dim(\chi)}$ is invertible in k , so we have determined the multiset $c_T(j)$ for j such that $\chi_{T(j)} = \chi$, i.e. the contents of $\lambda(\chi)$, viewed as elements of k . Together with $|\lambda(\chi)|$, which was determined by the action of \mathcal{R}_Γ , this determines the p -core of $\lambda(\chi)$, which completes the proof of the “only if” direction.

REMARK 7.10. [14, Appendix B, Chapter 1] computes the characters of $\Gamma \wr S_n$ in terms of the characters of Γ and S_n . This can be used to give an expression for the central characters, and in turn to prove Theorem 7.9.

EXAMPLE 7.11. As pointed out on [4, Page 1], over a field of characteristic 2, every irreducible representation of $C_2 \wr S_n$ can be obtained by pulling back an irreducible representation of S_n via the quotient map $C_2 \wr S_n \rightarrow S_n$. This does not, however, mean that the representation categories of $C_2 \wr S_n$ and S_n are equivalent. Note that there is only one 2-block B_i of C_2 , and this block is not semisimple. So in Theorem 7.9, condition (2) is vacuous, and condition (1) holds automatically since $\sum_{\chi \in B_i} |\lambda(\chi)| = n$ as there is only one block. As a result $C_2 \wr S_n$ has only one 2-block for any n . By comparison, Example 2.1 shows S_3 has two 2-blocks. In general, this argument shows that when Γ is a non-trivial p -group, $\Gamma \wr S_n$ will have only one p -block.

REMARK 7.12. It is straightforward to generalise the character symmetric function from Definition 3.9 to the case of wreath products. For a partially-reduced conjugacy class μ , we have an element of $\mathcal{R}_\Gamma \otimes \Lambda(\Gamma_*)$ given by $\psi^{-1}(K_\mu)$. Its “content evaluation” as in the proof of Theorem 7.9 will yield central characters of wreath products $\Gamma \wr S_n$.

APPENDIX A. PROPERTIES OF \mathcal{R}_Γ AND $\Lambda(\Gamma_*)$

In this appendix we show that both \mathcal{R}_Γ and $\Lambda(\Gamma_*)$ are Hopf algebras. We also explain that \mathcal{R}_Γ is the algebra of distributions on $Z\Gamma_*$ (viewed as an affine monoid scheme). Furthering the analogy with integer-valued polynomials, we classify homomorphisms from \mathcal{R}_Γ to a field; in positive characteristic, they are indexed by several p -adic parameters, one for each p -block of Γ .

First we show that $\Lambda(\Gamma_*)$ is a Hopf algebra.

THEOREM A.1. *There is a Hopf algebra structure on $\Lambda(\Gamma_*)$ with comultiplication*

$$\Delta(m_\lambda) = \sum_{\mu, \nu} m_\mu \otimes m_\nu,$$

where the sum is over all pairs of multipartitions μ, ν such that $m_i(\mu(c)) + m_i(\nu(c)) = m_i(\lambda(c))$ for all $i \in \mathbb{Z}_{>0}$ and $c \in \Gamma_*$. The counit ε obeys $\varepsilon(m_\mu) = 0$ when $|\mu| > 0$ (and $\varepsilon(1) = 1$).

Proof. We have that $(\Delta \otimes 1) \circ \Delta(m_\lambda) = (1 \otimes \Delta) \circ \Delta(m_\lambda)$, both being equal to

$$\sum_{\mu, \nu, \rho} m_\mu \otimes m_\nu \otimes m_\rho,$$

where the sum is over all multipartitions obeying $m_i(\mu(c)) + m_i(\nu(c)) + m_i(\rho(c)) = m_i(\lambda(c))$. This verifies that Δ is coassociative. To see that the ε defines the counit of a bialgebra, we note that

$$(\varepsilon \otimes 1) \circ \Delta(m_\lambda) = \sum_{\mu, \nu} \varepsilon(m_\mu) \otimes m_\nu,$$

and $\varepsilon(m_\mu)$ is nonzero only if $m_i(\mu(c)) = 0$ for all i and c , in which case $m_\nu = m_\lambda$. The case of $(1 \otimes \varepsilon) \circ \Delta$ is identical. Finally, we must show there is an antipode S . We do not give an explicit formula, but we prove it exists by induction on $|\lambda|$. We take $S(1) = 1$, which serves as the base case $|\lambda| = 0$. For $|\lambda| > 1$, we write

$$\Delta(m_\lambda) = m_\lambda \otimes 1 + \sum_{\mu, \nu} m_\mu \otimes m_\nu,$$

where we have written the term where ν is the empty partition (and $\mu = \lambda$) separately. The antipode axiom becomes

$$S(m_\lambda) + \sum_{\mu, \nu} S(m_\mu)m_\nu = 0.$$

But every term in the sum has $|\mu| < |\lambda|$, so this serves to define $S(m_\lambda)$ inductively. By construction, the resulting map S obeys the antipode axiom.

REMARK A.2. The Hopf algebra structure on $\Lambda(\Gamma_*)$ agrees with the usual Hopf algebra structure on Λ (see [14, Section 1.5, Example 25]) when we set Γ to be the trivial group. In fact, it can be shown that the isomorphism

$$\mathbb{C} \otimes \Lambda(\Gamma_*) = \mathbb{C} \otimes \Lambda^{\otimes |\Gamma^*|}$$

from Proposition 4.6 is an isomorphism of Hopf algebras, where $\Lambda^{\otimes |\Gamma^*|}$ is a tensor product of Hopf algebras, and is therefore a Hopf algebra.

REMARK A.3. One of the most important tools in the theory of symmetric functions is the Cauchy identity:

$$\prod_{i,j} \frac{1}{1 - x_i y_j} = \sum_{\lambda} s_\lambda(x) s_\lambda(y).$$

It would be desirable to have a similar identity for $\Lambda(\Gamma_*)$. However, the most natural formulation of the Cauchy identity is as a formula for the series

$$\sum_i b_i \otimes b_i^*,$$

where b_i is a basis of Λ and b_i is the dual basis with respect to the Hall inner product on Λ . The equation above takes both b_i and b_i^* to be the Schur functions s_λ , which are self dual. For this notion to make sense in $\Lambda(\Gamma_*)$, we need an inner product on $\Lambda(\Gamma_*)$. The natural choice, $\langle -, - \rangle$, comes from the isomorphism

$$\mathbb{C} \otimes \Lambda(\Gamma_*) = \mathbb{C} \otimes \Lambda^{\otimes |\Gamma^*|},$$

where each tensor factor of Λ is given the Hall inner product. The reason for this choice is that the Hall inner product defines a Hopf pairing on Λ , and since the Hopf algebra structure on $\Lambda(\Gamma_*)$ comes from the tensor product of Hopf algebra structures on Λ , the tensor product of Hall inner products defines a Hopf pairing on $\mathbb{C} \otimes \Lambda(\Gamma_*)$. Now we encounter a problem: $\Lambda(\Gamma_*)$ is not self dual with respect to this inner product: if b_i is a basis of $\Lambda(\Gamma_*)$, then b_i^* will be contained in

$$\Lambda(\Gamma_*)^* = \{f \in \mathbb{C} \otimes \Lambda^{\Gamma^*} \mid \langle f, g \rangle \in \mathbb{Z} \text{ for all } g \in \Lambda(\Gamma_*)\},$$

which is not a subset of $\Lambda(\Gamma_*)$; roughly speaking it is the “dual” of $\Lambda(\Gamma_*)$, so $\Lambda(\Gamma_*)^*$ inherits the structure of a Hopf algebra. It would be interesting to give an independent description of $\Lambda(\Gamma_*)^*$.

It would also be interesting to have a presentation of $\Lambda(\Gamma_*)$ by generators and relations. The argument from [23, Corollary 2.3] (filtration/induction on $l(\lambda)$) shows that $\Lambda(\Gamma_*)$ is generated by $m_{(r^n)_c}$ for $r, n \in \mathbb{Z}_{>0}$ and $c \in \Gamma_*$. However the argument from [23, Proposition 2.5] can be adapted to show that there are redundancies; the generator $m_{(r^n)_c}$ may be omitted if c is an n -th power in the ring $\mathbb{Z}\Gamma_*$.

Now we turn our attention to \mathcal{R}_Γ . Let R be an arbitrary commutative ring, and let us view $R\Gamma_* = \text{hom}_{\text{alg}}(A, R)$ as the affine scheme represented by $A = \mathbb{Z}[t_{c_1}, \dots, t_{c_l}]$, where the polynomial variables are indexed by elements of Γ_* . Now, $R\Gamma_*$ is a ring whose addition and multiplication maps are algebraic, i.e. the corresponding maps

$$R\Gamma_* \times R\Gamma_* \rightarrow R\Gamma_*$$

are morphisms of affine schemes. This amounts to saying that $R\Gamma_*$ is a (commutative) ring object in the category of affine schemes. On the level of coordinate rings, we get homomorphisms $\Delta^{(+)}, \Delta^{(\times)}$ from A to $A \otimes A$. Explicitly, they are defined by

$$\Delta^{(+)}(t_c) = t_c \otimes 1 + 1 \otimes t_c$$

and

$$\Delta^{(\times)}(t_c) = \sum_{d,e \in \Gamma_*} A_{d,e}^c t_d \otimes t_e.$$

We call these two maps the *coaddition* and *comultiplication*, respectively. They satisfy relations coming from the associative and distributive laws in a ring. Because of these two maps, a commutative ring object in the category of affine schemes is sometimes called a *birring*, although this is not the same thing as a bialgebra.

Now we let I be the ideal of A vanishing at the multiplicative identity; I is generated by $t_1 - 1$ and t_c for $c \neq 1$. We have $\Delta^{(\times)}(I) \subseteq I \otimes A + A \otimes I$ (this follows from the fact that I is the ideal corresponding to the identity element of $R\Gamma_*$, but can also be checked directly). We use I to define the distribution algebra of $R\Gamma_*$ (see [10, Chapter 7 of Part I]).

DEFINITION A.4. *The distribution algebra of $R\Gamma_*$ is*

$$D(R\Gamma_*) = \bigcup_{n>0} (A/I^n)^*,$$

where $(A/I^n)^*$ is viewed as a subspace of $A^* = \text{hom}_{\mathbb{Z}}(A, \mathbb{Z})$.

So $D(R\Gamma_*)$ is the set of linear functionals on A that vanish on some power of I . We obtain a multiplication on $D(R\Gamma_*)$ by dualising $\Delta^{(\times)}$. If $f_1, f_2 \in D(R\Gamma_*)$, then we take

$$(f_1 f_2)(a) = (f_1 \otimes f_2)(\Delta^{(\times)}(a))$$

for $a \in A$. The product $f_1 f_2$ vanishes on some power of I because we have

$$\Delta^{(\times)}(I^n) \subseteq \sum_{m=0}^n I^m \otimes I^{n-m},$$

and taking n sufficiently large we guarantee that each summand is annihilated by $f_1 \otimes f_2$. Finally, we see that the coassociativity of $\Delta^{(\times)}$ implies the associativity of the multiplication, and the canonical map $A \rightarrow A/I = \mathbb{Z}$ is the multiplicative identity. Thus $D(R\Gamma_*)$ is an associative ring.

THEOREM A.5. *We have that $D(R\Gamma_*) = \mathcal{R}_{\Gamma}$.*

Proof. Define the variables $z_1 = t_1 - 1$ and $z_c = t_c$ for $c \neq 1$. The coordinate ring A has a basis consisting of monomials $\prod_{c \in \Gamma_*} z_c^{\mathbf{N}(c)}$, indexed by $\mathbf{N} \in \mathbb{Z}_{>0}^{\Gamma_*}$. Such a monomial is contained in $I^{|\mathbf{N}|}$. Hence $D(R\Gamma_*)$ has \mathbb{Z} -basis consisting of the dual basis, which we denote $\beta_{\mathbf{N}}$. The structure constants of the multiplication with respect to the basis $\beta_{\mathbf{N}}$ are precisely the structure constants of $\Delta^{(\times)}$ with respect to the monomials $\prod_c z_c^{\mathbf{N}(c)}$. To compute the multiplication, note that for any $c \in \Gamma_*$,

$$\Delta^{(\times)}(z_c) = z_c \otimes 1 + 1 \otimes z_c + \sum_{d,e} A_{d,e}^c z_d \otimes z_e.$$

We recognise this as the change of variables appearing in Proposition 5.13 (subject to the relabelling $z_i \otimes 1 = x_i$ and $1 \otimes z_i = y_i$) that describes the structure constants of the multiplication in \mathcal{R}_{Γ} . This implies that the linear map $\mathcal{R}_{\Gamma} \rightarrow D(R\Gamma_*)$ given by $B_{\mathbf{N}} \mapsto \beta_{\mathbf{N}}$ is a ring isomorphism.

It is well known that over a field of characteristic zero, the distribution algebra of an algebraic group G coincides with the universal enveloping algebra of the Lie algebra of G . Proposition 5.13 witnesses this fact; the proof explains that $\mathbb{Q} \otimes \mathcal{R}_\Gamma = \mathcal{U}(\mathbb{Q}\Gamma_*)$. Although $R\Gamma_*$ is a monoid rather than a group, we may consider the group of units, which has same distribution algebra. Since distribution algebras of algebraic groups are Hopf algebras, the same is true of \mathcal{R}_Γ .

PROPOSITION A.6. *We have that $\mathcal{R}_\Gamma \subseteq \mathcal{U}(\mathbb{Q}\Gamma_*)$ is a Hopf subalgebra.*

Now we turn our attention to classifying maps into a field. For context and background, we direct the reader to [8]. We write \mathbb{Z}_p for the set of p -adic integers. The motivating example is the following.

EXAMPLE A.7 ([8, Section 8]). Homomorphisms from the ring \mathcal{R} of integer-valued polynomials to a field F are given by the following. If F has characteristic zero, they are given by evaluating the polynomial variable t at an arbitrary element of F . If F has characteristic $p > 0$, then they are given by evaluating the polynomial variable t at a p -adic integer. Evaluation of an integer-valued polynomial at a p -adic integer yields another p -adic integer, which may be viewed as an element of F via the canonical map

$$\mathbb{Z}_p \rightarrow \mathbb{Z}_p/(p) = \mathbb{Z}/p\mathbb{Z} \subseteq F.$$

In fact, if $t \in \mathbb{Z}_p$, $\binom{t}{p^r}$ is the r -th p -adic digit of t (viewed as an element of F).

First we get the characteristic-zero case out of the way.

PROPOSITION A.8. *Suppose that F is a field of characteristic zero. Homomorphisms $\mathcal{R}_\Gamma \rightarrow F$ are given by evaluating each $T(c)$ at an arbitrary element of F (and are hence parametrised by $F^{\Gamma_*} = F\Gamma_*$).*

Proof. We have $\mathbb{Q} \subseteq F$, and $\mathbb{Q} \otimes \mathcal{R}_\Gamma = \mathcal{U}(\mathbb{Q}\Gamma_*)$ is the free polynomial algebra in variables $T(c)$ for $c \in \Gamma_*$. Homomorphisms from a polynomial algebra to F amount to evaluating each variable at an element of F .

Now we move on to the positive characteristic case. Fix a prime p , and let F be a field of characteristic p . Since any field embeds in its algebraic closure, we may as well take F to be algebraically closed.

DEFINITION A.9. *Let $q \in F\Gamma_*$ be $q = \sum_{c \in \Gamma_*} m_c c$. We define*

$$B_{q,r} = \sum_{|\mathbf{M}|=p^r} \prod_{c \in \Gamma_*} m_c^{\mathbf{M}(c)} B_{\mathbf{M}}.$$

For example, if $q = c \in \Gamma_*$, we have

$$B_{c,r} = B_{\mathbf{M}}$$

where $\mathbf{M}(c) = p^r$ and $\mathbf{M}(c') = 0$ for $c' \neq c$. Also, $B_{0,r} = 0$ for all r .

LEMMA A.10. *The algebra $F \otimes \mathcal{R}_\Gamma$ is generated by the elements $B_{q,r}$, where q varies across a basis of $F\Gamma_*$ and $r \in \mathbb{Z}_{>0}$,*

Proof. Lemma 5.12 asserts that $B_{\mathbf{M}}$ has leading order term

$$\prod_{c \in \Gamma_*} \frac{T(c)^{\mathbf{M}(c)}}{\mathbf{M}(c)!}.$$

Writing $q = \sum_{c \in \Gamma_*} m_c c$ and using the multinomial theorem, we see that the leading order term of $B_{q,r}$ is

$$\sum_{|\mathbf{M}|=p^r} \prod_{c \in \Gamma_*} m_c^{\mathbf{M}(c)} \frac{T(c)^{\mathbf{M}(c)}}{\mathbf{M}(c)!} = \frac{T(q)^{p^r}}{p^r!}.$$

If the p -adic decomposition of n is $\sum_r n_r p^r$, then over F we have

$$\prod_r \left(\frac{T(q)^{p^r}}{p^{r!}} \right)^{n_r} = \frac{n!}{\prod_r (p^{r!})^{n_r}} \cdot \frac{T(q)^n}{n!}.$$

Here, the multinomial coefficient

$$\frac{n!}{\prod_r (p^{r!})^{n_r}}$$

is invertible in F . This implies that the leading order term of any B_M is a (scalar multiple of a) product of leading order terms of the $B_{c,r}$, where we are using the particular basis of conjugacy class sums, proving the lemma for that particular basis. To prove the lemma for an arbitrary basis q_i , we express $B_{c,r}$ in terms of the $B_{q_i,r}$. Let $c = \sum_i k_i q_i$, and observe that to leading order, $B_{c,r}$ may be written

$$\frac{T(\sum_i k_i q_i)^{p^r}}{p^{r!}} = \sum_{|\mathbf{W}|=p^r} \prod_i k_i^{\mathbf{W}(q_i)} \frac{T(q_i)^{\mathbf{W}(q_i)}}{\mathbf{W}(q_i)!},$$

where the sum is over functions \mathbf{W} from the basis q_i to $Z_{>0}$, such that $\sum_i \mathbf{W}(q_i) = p^r$. We observe that this is equal to

$$\sum_i k_i^{p^r} \frac{T(q_i)^{p^r}}{p^{r!}}$$

plus “mixed” terms whose exponents (values of $\mathbf{W}(q)$) are less than p^r and may therefore be written in terms of $B_{q,s}$ for $s < r$. By induction on r , the generating set $B_{c,r}$ is contained in the subring generated by all elements of the form $B_{q,r}$, and this completes the proof.

LEMMA A.11. *View the map $ev_n^{\mathcal{R}}$ from Theorem 5.18 as having domain $F \otimes \mathcal{R}_\Gamma$ and codomain $(F\Gamma_*)^{\otimes n}$. We have that $ev_n^{\mathcal{R}}(B_{q,r})$ is equal to the sum of all pure tensors with p^r entries set to q , and the remainder set to 1.*

Proof. Expanding in terms of the basis Γ_* ,

$$q^{\otimes p^r} = \left(\sum_c m_c c \right)^{\otimes p^r},$$

we get each tensor monomial with $\mathbf{M}(c)$ factors equal to c with multiplicity in the conjugacy class sums c with multiplicity $\prod_{c \in \Gamma_*} m_c^{\mathbf{M}(c)}$. Inserting $n - p^r$ tensor factors of 1 in all possible ways completes the proof.

LEMMA A.12. *For any $q \in F\Gamma_*$, we have the following equation in $F \otimes \mathcal{R}_\Gamma$:*

$$B_{q,r}^p = B_{q^p,r},$$

where q^p is viewed as an element of $F\Gamma_*$.

Proof. We use Theorem 5.18 and the fact that the b_N are linearly independent. So it is enough to show that

$$ev_n^{\mathcal{R}}(B_{q,r})^p = ev_n^{\mathcal{R}}(B_{q^p,r})$$

over F for all n . We note that $ev_n^{\mathcal{R}}(B_{q,p^r})$ is the sum of all pure tensors (with respect to the basis Γ_*) with p^r factors set to q , and the remainder set to 1 (see Lemma 5.17). Raising this sum to the p -th power is the same as raising each pure tensor to the p -th power because the binomial coefficients from the cross-terms vanish in characteristic p . So we have the sum of pure tensors with p^r factors set to q^p , which agrees with $ev_n^{\mathcal{R}}(B_{q^p,r})$.

PROPOSITION A.13. Let $\varphi : F \otimes \mathcal{R}_\Gamma \rightarrow F$ be a ring homomorphism. If $q \in J(F\Gamma_*)$, then $\varphi(B_{q,r}) = 0$, where J is the Jacobson radical. If $e \in F\Gamma_*$ is idempotent, then $\varphi(B_{e,r})$ is an element of the prime subfield of F (isomorphic to Z/pZ).

Proof. Because $F\Gamma_*$ is a finite-dimensional algebra, its Jacobson radical is nilpotent. This means that $J(F\Gamma_*)^{p^k} = 0$ for some k . Hence for q in the Jacobson radical,

$$\begin{aligned} \varphi(B_{q,r})^{p^k} &= \varphi(B_{q,r}^{p^k}) \\ &= \varphi(B_{q^{p^k},r}) \\ &= \varphi(B_{0,r}) \\ &= 0. \end{aligned}$$

Since the only nilpotent element of a field is zero, $\varphi(B_{q,r}) = 0$. If e is idempotent,

$$\varphi(B_{e,r})^p = \varphi(B_{e^p,r}) = \varphi(B_{e,r}).$$

This means $\varphi(B_{q,r})$ solves the equation $x^p = x$ (i.e. is Frobenius-fixed), but such solutions are precisely elements of the prime subfield of F .

To avoid conflicts of notation, instead of writing B_i for the blocks of $F\Gamma$, we instead write C_u . Correspondingly, we write $\omega_c^u \in F$ for the central character of the block C_u evaluated at the conjugacy class sum c .

THEOREM A.14. Homomorphisms $\varphi : \mathcal{R}_\Gamma \rightarrow F$ are given by evaluating

$$T(c) \mapsto \sum_u \omega_c^u t_u,$$

where the sum is over the blocks C_u of $F\Gamma$, and $t_u \in Z_p$ are arbitrary p -adic integers (one for each block C_u).

Proof. For each block idempotent $e_u \in C_u$ (e_u is the identity element of C_u , viewed as an element of $F\Gamma_*$), $\varphi(B_{e_u,r})$ is an element of Z/pZ . Thus

$$t_u = \sum_r \varphi(B_{e_u,r}) p^r$$

are well-defined p -adic integers. First of all, we show that the t_u determine φ . It is enough to show that the t_u determine φ on the generating set from Lemma A.10. As discussed in the proof of Proposition 2.9, $Z(C_u)/J(Z(C_u)) = F$, and since $J(F\Gamma_*) = \bigoplus_u J(Z(C_u))$, any basis of $J(F\Gamma_*)$ together with the identity elements e_u of the blocks C_u form a basis of $F\Gamma_*$. Let q be an element of this basis of $F\Gamma_*$. Lemma A.10 implies that $B_{q,r}$ generates $F \otimes \mathcal{R}_\Gamma$, but Proposition A.13 implies that $\varphi(B_{q,r}) = 0$ for q in $J(F\Gamma_*)$. So φ is determined by the values on $B_{e_u,r}$, and this is precisely what is encoded in the t_u .

Now we show that any choice of the t_u gives rise to a homomorphism φ . We describe it via an equality of generating functions:

$$\sum_{\mathbf{N}} \varphi(B_{\mathbf{N}}) x^{\mathbf{N}} = \prod_u \left(1 + \sum_c \omega_c^u x_c \right)^{t_u}.$$

Here we are using the series expansion

$$(1+x)^t = \sum_r x^r \binom{t}{r}.$$

Because an integer-valued polynomial (e.g. $\binom{t}{r}$) evaluated at a p -adic integer is again a p -adic integer, $\varphi(B_{\mathbf{N}})$ is a well-defined element of F . It remains to check that φ defines a homomorphism, and that it actually corresponds to the parameters t_u . To check φ

is a homomorphism, we first observe that because central characters are themselves homomorphisms $F\Gamma_* \rightarrow F$,

$$\sum_i \omega_{c_i}^u x_{c_i} \sum_j \omega_{c_j}^u y_{c_j} = \sum_k \sum_{i,j} A_{i,j}^k \omega_{c_k}^u x_{c_i} y_{c_j}.$$

Now we compute

$$\begin{aligned} \sum_{\mathbf{N}} \varphi(B_{\mathbf{N}}) x^{\mathbf{N}} \sum_{\mathbf{M}} \varphi(B_{\mathbf{M}}) y^{\mathbf{M}} &= \prod_i (1 + \sum_i \omega_{c_i}^u x_{c_i})^{t_u} \prod_j (1 + \sum_j \omega_{c_j}^u y_{c_j})^{t_u} \\ &= \prod_u (1 + \sum_i \omega_{c_i}^u x_{c_i} + \sum_j \omega_{c_j}^u y_{c_j} + \sum_k \sum_{i,j} A_{i,j}^k \omega_{c_k}^u x_{c_i} y_{c_j})^{t_u} \\ &= \prod_u (1 + \sum_i \omega_{c_i}^u (x_{c_i} + y_{c_i} + \sum_{j,k} A_{j,k}^i x_j y_k))^{t_u} \\ &= \prod_u (1 + \sum_i \omega_{c_i}^u z_{c_i})^{t_u} \\ &= \sum_{\mathbf{K}} \varphi(B_{\mathbf{K}}) z^{\mathbf{K}}, \end{aligned}$$

where $z_{c_i} = x_{c_i} + y_{c_i} + \sum_{j,k} A_{j,k}^i x_j y_k$, as in Section 5. Proposition 5.11 now implies that φ is a homomorphism. It remains to verify that $\varphi(B_{e_u,r})$ is equal to the r -th p -adic digit of t_u . We fix a block C_v and write $e_v = \sum m_c c$. Now, to compute

$$ev(B_{e_v,r}) = \sum_{|\mathbf{M}|=p^r} \prod_{c \in \Gamma_*} m_c^{\mathbf{M}(c)} ev(B_{\mathbf{M}}),$$

we take the generating function we used to define φ and evaluate at $x_c = \epsilon m_c$, where ϵ is a formal variable. This gives

$$\sum_{\mathbf{N}} \varphi(B_{\mathbf{N}}) \prod_c m_c^{\mathbf{N}(c)} \epsilon^{|\mathbf{N}|} = \prod_u (1 + \sum_c \omega_c^u m_c \epsilon)^{t_u}.$$

We observe that $\sum_c \omega_c^u m_c$ is the central character of C_u evaluated at e_v , and therefore this sum is equal to one if $u = v$, and zero otherwise. This shows that $\varphi(B_{e_v,r})$ is equal to the coefficient of ϵ^{p^r} in

$$(1 + \epsilon)^{t_v}.$$

But this is precisely the r -th p -adic digit of t_v by Example A.7.

REFERENCES

- [1] R. Brauer and G de B Robinson, *On a conjecture by Nakayama*, Trans. Roy. Soc. Canada Sect. III **41** (1947), 20–25.
- [2] Tullio Ceccherini-Silberstein, Fabio Scarabotti, and Filippo Tolli, *Representation theory of the symmetric groups: the okounkov-vershik approach, character formulas, and partition algebras*, vol. 121, Cambridge University Press, 2010.
- [3] Sylvie Corteel, Alain Goupil, and Gilles Schaeffer, *Content evaluation and class symmetric functions*, Adv. Math. **188** (2004), no. 2, 315–336.
- [4] Richard Dipper and Gordon James, *Representations of Hecke algebras of type B_n* , J. Algebra **146** (1992), no. 2, 454–481.
- [5] Pavel I Etingof, Oleg Golberg, Sebastian Hensel, Tiankai Liu, Alex Schwendner, Dmitry Vaintrob, and Elena Yudovina, *Introduction to representation theory*, vol. 59, American Mathematical Soc., 2011.
- [6] Hanafi K Farahat and Graham Higman, *The centres of symmetric group rings*, Proc. R. Soc. Lond. Ser. A. Phys. Eng. Sci. **250** (1959), no. 1261, 212–221.
- [7] Georg Ferdinand Frobenius, *Über die charaktere der symmetrischen gruppe*, Königliche Akademie der Wissenschaften, 1900.
- [8] Nate Harman and Sam Hopkins, *Quantum integer-valued polynomials*, J. Algebraic Combin. **45** (2017), no. 2, 601–628.

- [9] Vladimir N Ivanov and Sergei V Kerov, *The algebra of conjugacy classes in symmetric groups and partial permutations*, J. Math. Sci. (N.Y.) **107** (2001), no. 5, 4212–4230.
- [10] Jens Carsten Jantzen, *Representations of algebraic groups*, vol. 107, American Mathematical Soc., 2003.
- [11] A.-A. A. Jucys, *Symmetric polynomials and the center of the symmetric group ring*, Rep. Mathematical Phys. **5** (1974), no. 1, 107–112.
- [12] Alexander Kleshchev, *Linear and projective representations of symmetric groups*, no. 163, Cambridge University Press, 2005.
- [13] Manfred Lehn and Christoph Sorger, *Symmetric groups and the cup product on the cohomology of Hilbert schemes*, Duke Math. J. **110** (2001), no. 2, 345–357.
- [14] Ian G Macdonald, *Symmetric functions and hall polynomials*, second ed., Oxford mathematical monographs, Clarendon Press New York, Oxford, 1995.
- [15] Ashish Mishra and Murali K Srinivasan, *The Okounkov–Vershik approach to the representation theory of $G \wr S_n$* , J. Algebraic Combin. **44** (2016), no. 3, 519–560.
- [16] G E Murphy, *The idempotents of the symmetric group and Nakayama’s conjecture*, J. Algebra **81** (1983), no. 1, 258–265.
- [17] Andrei Okounkov and Anatoly Vershik, *A new approach to representation theory of symmetric groups*, Selecta Math. (N.S.) **2** (1996), no. 4, 581–606.
- [18] Igor A Pushkarev, *On the representation theory of wreath products of finite groups and symmetric groups*, J. Math. Sci. (N.Y.) **96** (1999), no. 5, 3590–3599.
- [19] Bannikuppe M Puttaswamaiah and John D Dixon, *Modular representations of finite groups*, Academic press, 1977.
- [20] Christopher Ryba, *Stable Grothendieck rings of wreath product categories*, J. Algebraic Combin. **49** (2019), no. 3, 267–307.
- [21] ———, *The structure of the grothendieck rings of wreath product deligne categories and their generalisations*, Int. Math. Res. Not. IMRN **2021** (2019), no. 16, 12420–12462.
- [22] Omar Tout, *The algebra of conjugacy classes of the wreath product of a finite group with the symmetric group*, 2021, <https://arxiv.org/abs/math/2107.05024>.
- [23] Francesco Vaccarino, *The ring of multisymmetric functions*, Ann. Inst. Fourier (Grenoble), vol. 55, 2005, pp. 717–731.
- [24] Weiqiang Wang, *The Farahat–Higman ring of wreath products and Hilbert schemes*, Adv. Math. **187** (2004), no. 2, 417–446.
- [25] ———, *Vertex algebras and the class algebras of wreath products*, Proc. Lond. Math. Soc. (3) **88** (2004), no. 2, 381–404.
- [26] Geordie Williamson, *Schubert calculus and torsion explosion*, J. Amer. Math. Soc. **30** (2017), no. 4, 1023–1046.

CHRISTOPHER RYBA, Department of Mathematics, University of California, Berkeley, CA 94720, USA
E-mail : ryba@math.berkeley.edu