



ALGEBRAIC COMBINATORICS

Nate Harman & Christopher Ryba

A tensor-cube version of the Saxl conjecture

Volume 6, issue 2 (2023), p. 507-511.

<https://doi.org/10.5802/alco.267>

© The author(s), 2023.



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





A tensor-cube version of the Saxl conjecture

Nate Harman & Christopher Ryba

ABSTRACT Let n be a positive integer, and let $\rho_n = (n, n-1, n-2, \dots, 1)$ be the “staircase” partition of size $N = \binom{n+1}{2}$. The Saxl conjecture asserts that every irreducible representation S^λ of the symmetric group S_N appears as a subrepresentation of the tensor square $S^{\rho_n} \otimes S^{\rho_n}$. In this short note we give two proofs that every irreducible representation of S_N appears in the tensor cube $S^{\rho_n} \otimes S^{\rho_n} \otimes S^{\rho_n}$.

1. INTRODUCTION

Let n , ρ_n , and N be as defined in the abstract. In 2012, Jan Saxl conjectured that $S^{\rho_n} \otimes S^{\rho_n}$ contains every irreducible representation of S_N as a subrepresentation. This was motivated by a similar phenomenon for finite simple groups of Lie type [3]. Despite a great amount of effort, tensor products of symmetric group representations are poorly understood from a combinatorial point of view. The Saxl conjecture serves as one benchmark for results in this area. Progress has been made using a wide variety of approaches, for example, using ideas from combinatorics [9, 6], modular representation theory [1, 2], or probability [10, 7]. In this last paper mentioned, Luo and Sellke proved a weaker version of the Saxl conjecture, namely that for n sufficiently large, $(S^{\rho_n})^{\otimes 4}$ contains every irreducible representation of S_N as a subrepresentation. Our main theorem is the following improvement:

THEOREM 1.1. $(S^{\rho_n})^{\otimes 3}$ contains all irreducible representations of S_N .

We note that the results of Luo and Sellke extend to N not of the form $\binom{n+1}{2}$, while ours rely on special properties of staircase partitions which do not seem to generalize. We provide two proofs, the first using combinatorics, and the second using modular representation theory.

Manuscript received 11th July 2022, revised 31st August 2022, accepted 3rd September 2022.

KEYWORDS. Saxl conjecture, symmetric groups.

ACKNOWLEDGEMENTS. The first author would like to thank Joshua Mundinger for helpful conversations. The second author would like to thank Mark Sellke for helpful comments on a previous version of this paper.

2. PROOF USING COMBINATORICS

For now, let N be an arbitrary positive integer, which we will specialize to $\binom{n+1}{2}$ at the end of this section. We use the following result of Luo and Sellke (although for our final conclusion the prior result of Ikenmeyer [4] is sufficient).

LEMMA 2.1 (Luo-Sellke [7, Appendix B]). *Let μ, ν be partitions of size N such that μ has distinct parts, and ν is greater than or equal to μ in the dominance order. Then $S^\mu \otimes S^\mu$ contains S^ν as a subrepresentation.*

If $\mu = (\mu_1, \mu_2, \dots, \mu_r)$ is a partition of N , let M^μ be the permutation representation on cosets of the Young subgroup $S_\mu = S_{\mu_1} \times S_{\mu_2} \times \dots \times S_{\mu_r}$ of S_N . Equivalently, $M^\mu = \text{Ind}_{S_\mu}^{S_N}(\mathbf{1})$, where $\mathbf{1}$ is the trivial representation.

DEFINITION 2.2. *Given a partition μ , let $C(\mu)$ be the partition obtained as follows. Start off with the empty partition. For each $i \geq 1$, add i parts of size $\mu_i - \mu_{i+1}$ (if i is greater than the length of μ , we define $\mu_i = 0$).*

EXAMPLE 2.3. Suppose that $\mu = \rho_n$ is the staircase partition of size $\binom{n+1}{2}$. Then we have $\mu_i - \mu_{i+1} = 1$ for $i = 1, 2, \dots, n$. Thus $C(\mu)$ consists of the part 1 repeated $1 + 2 + \dots + n = \binom{n+1}{2}$ times.

THEOREM 2.4. *Let μ be any partition. Then $M^\mu \otimes S^\mu$ contains $M^{C(\mu)}$ as a subrepresentation.*


Proof. It is convenient to take Frobenius characteristics to turn representations into symmetric functions. The Frobenius characteristics of M^μ and S^μ are h_μ and s_μ respectively. Their tensor product is described by the internal product of symmetric functions, which admits the following description (see [8, §1.7, Example 23(c)]):

$$h_\mu * s_\mu = \sum \prod_{i \geq 1} s_{\lambda^{(i)}/\lambda^{(i+1)}}$$

where the sum ranges over all nested families $\mu = \lambda^{(1)} \supset \lambda^{(2)} \supset \dots \supset \emptyset$ of partitions $\lambda^{(i)}$ for which $|\lambda^{(i)}| - |\lambda^{(i+1)}| = \mu_i$.

Each summand is the Frobenius characteristic of a genuine (rather than virtual) representation, so to show that $M^{C(\mu)}$ is a subrepresentation of $M^\mu \otimes S^\mu$, it suffices to show that one of the summands is equal to $h_{C(\mu)}$. We do this by considering the term where $\lambda^{(i)}$ is obtained from μ by removing the first $(i - 1)$ rows (so $\lambda_j^{(i)} = \mu_{j+i-1}$). Then it is not difficult to see that the skew-diagram $\lambda^{(i)}/\lambda^{(i+1)}$ consists of the disjoint union of rows of lengths $\mu_i - \mu_{i+1}, \mu_{i+1} - \mu_{i+2}, \dots$. We illustrate this with an example.

Suppose that $\mu = (10, 6, 4, 1)$. Then we have $\lambda^{(1)} = (10, 6, 4, 1)$, $\lambda^{(2)} = (6, 4, 1)$, $\lambda^{(3)} = (4, 1)$, $\lambda^{(4)} = (1)$. In the following diagram, $\lambda^{(1)}/\lambda^{(2)}$ is depicted in red, $\lambda^{(2)}/\lambda^{(3)}$ is depicted in orange, $\lambda^{(3)}/\lambda^{(4)}$ is depicted in green, and $\lambda^{(4)}$ is depicted in blue:



In particular a horizontal row of size $\mu_1 - \mu_2 = 4$ appears once, a horizontal row of size $\mu_2 - \mu_3 = 2$ appears twice, a horizontal row of size $\mu_3 - \mu_4 = 3$ appears three times, and a horizontal row of size $\mu_4 = 1$ appears 4 times.

Since a skew-Schur function for a skew-diagram D is the product of the skew-Schur functions associated to the connected components of D , and the skew Schur function associated to a horizontal row of size r is $s_{(r)} = h_r$, we conclude that $s_{\lambda^{(i)}/\lambda^{(i+1)}} = h_{\mu_i - \mu_{i+1}} \cdot h_{\mu_{i+1} - \mu_{i+2}} \cdot \dots$. Thus

$$\prod_i s_{\lambda^{(i)}/\lambda^{(i+1)}} = h_{C(\mu)},$$

and so we have found the required summand. □

COROLLARY 2.5. *Let μ be a partition of size N with distinct parts. Then for each λ greater than or equal to $C(\mu)$ in the dominance order, $S^\mu \otimes S^\mu \otimes S^\mu$ contains S^λ as a summand.*

Proof. By Lemma 2.1 $S^\mu \otimes S^\mu$ contains S^ν for every ν greater than or equal to μ in the dominance order. In particular this means $S^\mu \otimes S^\mu$ contains a copy of every irreducible constituent of M^μ . Therefore $S^\mu \otimes S^\mu \otimes S^\mu$ contains all of the irreducible constituents of $M^\mu \otimes S^\mu$, which by Theorem 2.4 includes all S^λ with λ greater than or equal to $C(\mu)$ in the dominance order. □

First Proof of Theorem 1.1: By Corollary 2.5, $S^{\rho_n} \otimes S^{\rho_n} \otimes S^{\rho_n}$ contains S^λ for every partition λ greater than or equal to $C(\rho_n) = (1^N)$ (see Example 2.3). However, every partition of N is greater than or equal to (1^N) in the dominance order. □

3. PROOF USING MODULAR REPRESENTATION THEORY

We will now give a second proof of the main theorem, this time following the approach laid out in [1] to study the Saxl conjecture via the 2-modular representation theory of symmetric groups. The proof itself will be quite short, but in the interest of self-containment and readability we will first state the facts from modular representation theory we will be using.

First we will recall some general facts about projective objects in the modular representation theory of a finite group. These are all fairly standard results in modular representation theory and can be found in many places, but as a standard reference we will refer to [11].

LEMMA 3.1 (See [11, Chapters 14 and 15]). *Let G be a finite group, $k = \bar{k}$ be an algebraically closed field of characteristic $p > 0$, and let $\text{Rep}_p(G)$ denote the category of finite dimensional representations of G over k .*

- (1) *The number of isomorphism classes of projective indecomposable objects in $\text{Rep}_p(G)$ is equal to the number of simple objects $\text{Rep}_p(G)$, which is equal to the number of conjugacy classes of elements of G with order prime to p . Moreover, each simple object occurs uniquely as an irreducible quotient of a projective indecomposable object.*
- (2) *If P is a projective object of $\text{Rep}_p(G)$, and V is any representation then $P \otimes V$ is again a projective object. Moreover if P is a fixed projective object, then every indecomposable projective object P' arises as a direct summand in a module of the form $P \otimes X$ where X is simple.*
- (3) *If P is a projective object in $\text{Rep}_p(G)$, then P lifts to characteristic zero – more precisely P is isomorphic to $\tilde{P} \otimes_{\mathcal{O}} \mathcal{O}/\mathfrak{m}\mathcal{O}$ where \mathcal{O} denotes the ring of Witt vectors for k , \mathfrak{m} is the unique maximal ideal in \mathcal{O} , and \tilde{P} is projective in the category of representations of G over \mathcal{O} .*
- (4) *Every ordinary irreducible representation of G over $K = \text{frac}(\mathcal{O})$ occurs inside $\tilde{P} \otimes_{\mathcal{O}} K$ for some projective indecomposable representation \tilde{P} of G over \mathcal{O} .*

Next we will recall some basic facts and notation specific to the modular representation theory of symmetric groups.

LEMMA 3.2 (See [5, Chapter 6]).

- (1) *Simple objects in $\text{Rep}_p(S_n)$ are indexed by p -regular partitions λ (that is, partitions where there are at most $p - 1$ parts of any given size) and are denoted by D^λ .*
- (2) *D^λ appears with multiplicity one as a composition factor inside any reduction mod p of the corresponding ordinary irreducible representation S^λ .*
- (3) *If λ is a p -core (meaning all of the hook lengths of λ are relatively prime to p) then the reduction of S^λ modulo p is simple and projective.*

In the previous section we referred to partitions with distinct parts, and here we will refer to 2-regular partitions. These are the same notion, but the former terminology is more common in the combinatorics literature whereas the latter is more common in the modular representation theory literature so we feel the inconsistency in terminology is warranted.

Second Proof of Theorem 1.1: By Lemma 2.1 (or its predecessor in [4]) the tensor square $S^{\rho_n} \otimes S^{\rho_n}$ contains every S^λ with $\lambda \geq \rho_n$ in the dominance order. In particular this includes all 2-regular partitions λ . Therefore any reduction mod 2 of $S^{\rho_n} \otimes S^{\rho_n}$ contains every irreducible 2-modular representation D^λ as a composition factor by Lemma 3.2 part (2).

By Lemma 3.2 part (3), S^{ρ_n} is projective in characteristic 2 since ρ_n is a 2-core. So by Lemma 3.1 part (2) when we tensor with the third copy of S^{ρ_n} it splits all extensions and we see that $S^{\rho_n} \otimes S^{\rho_n} \otimes S^{\rho_n}$ contains a copy of $D^\lambda \otimes S^{\rho_n}$ as a direct summand for every 2-regular partition λ .

By part (2) of Lemma 3.1 this means that every projective indecomposable object in $\text{Rep}_2(S_N)$ is a summand of $S^{\rho_n} \otimes S^{\rho_n} \otimes S^{\rho_n}$, which lifting back to characteristic zero implies every ordinary irreducible representation occurs in $S^{\rho_n} \otimes S^{\rho_n} \otimes S^{\rho_n}$ by part (4) of Lemma 3.1. \square

REMARK 3.3. A 2-modular strengthened Saxl’s conjecture was made in [1] saying that every projective indecomposable object in $\text{Rep}_2(S_N)$ should occur as a direct summand of the Saxl square $S^{\rho_n} \otimes S^{\rho_n}$. We will note that this proof proves the tensor cube version of this conjecture along the way.

REFERENCES

- [1] C. Bessenrodt, C. Bowman, and L. Sutton, *Kronecker positivity and 2-modular representation theory*, Trans. Amer. Math. Soc. Ser. B **8** (2021), 1024–1055.
- [2] Christine Bessenrodt and Chris Bowman, *Splitting Kronecker squares, 2-decomposition numbers, Catalan combinatorics, and the Saxl conjecture*, preprint, 2022, <https://arxiv.org/abs/2202.03066>.
- [3] Gerhard Heide, Jan Saxl, Pham Huu Tiep, and Alexandre E. Zalesski, *Conjugacy action, induced representations and the Steinberg square for simple groups of Lie type*, Proc. Lond. Math. Soc. (3) **106** (2013), no. 4, 908–930.
- [4] Christian Ikenmeyer, *The Saxl conjecture and the dominance order*, Discrete Math. **338** (2015), no. 11, 1970–1975.
- [5] Gordon James and Adalbert Kerber, *The representation theory of the symmetric group*, Encyclopedia of Mathematics and its Applications, vol. 16, Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [6] Xin Li, *Saxl conjecture for triple hooks*, Discrete Math. **344** (2021), no. 6, article no. 112340 (19 pages).
- [7] Sammy Luo and Mark Sellke, *The Saxl conjecture for fourth powers via the semigroup property*, J. Algebraic Combin. **45** (2017), no. 1, 33–80.

- [8] I. G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Classic Texts in the Physical Sciences, The Clarendon Press, Oxford University Press, New York, 2015.
- [9] Igor Pak, Greta Panova, and Ernesto Vallejo, *Kronecker products, characters, partitions, and the tensor square conjectures*, Adv. Math. **288** (2016), 702–731.
- [10] Mark Sellke, *Covering $\text{Irrep}(S_n)$ with tensor products and powers*, preprint, 2020, <https://arxiv.org/abs/2004.05283>.
- [11] Jean-Pierre Serre, *Linear representations of finite groups*, Graduate Texts in Mathematics, Vol. 42, Springer-Verlag, New York-Heidelberg, 1977, Translated from the second French edition by Leonard L. Scott.

NATE HARMAN, Department of Mathematics, University of Michigan, Ann Arbor, MI 48101 (USA)
E-mail : nharman@umich.edu

CHRISTOPHER RYBA, Department of Mathematics, University of California, Berkeley, Berkeley, CA 94720 (USA)
E-mail : ryba@math.berkeley.edu