



ALGEBRAIC COMBINATORICS

Malte Gerhold & Philipp Varšo

Towards a classification of multi-faced independences: a combinatorial approach

Volume 7, issue 3 (2024), p. 679-711.

<https://doi.org/10.5802/alco.356>

© The author(s), 2024.




This article is licensed under the
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Towards a classification of multi-faced independences: a combinatorial approach

Malte Gerhold & Philipp Varšo

ABSTRACT We determine a set of necessary conditions on a partition-indexed family of complex numbers to be the “highest coefficients” of a positive and symmetric multi-faced universal product, i.e. the product associated with a multi-faced version of noncommutative stochastic independence, such as bifreeness. The highest coefficients of a universal product are the weights of the moment-cumulant relation for its associated independence. We show that these conditions are *almost* sufficient, in the sense that whenever the conditions are satisfied, one can associate a (automatically unique) symmetric universal product with the prescribed highest coefficients. Furthermore, we give a quite explicit description of such families of coefficients, thereby producing a list of candidates that must contain all positive symmetric universal products. We discover in this way four (three up to trivial face-swapping) previously unknown moment-cumulant relations that give rise to symmetric universal products; to decide whether they are positive, and thus give rise to independences which can be used in an operator algebraic framework, remains an open problem.

1. INTRODUCTION

At the latest with Voiculescu’s invention of *freeness* [31], it became apparent that the “obvious” extension of classical stochastic independence, *tensor independence*, is not the only and not always the most suitable concept in inherently noncommutative situations. In fact, *Boolean independence* (not yet under this name) has already featured much earlier in the work of von Waldenfels [33, 34]. Those “noncommutative independences” share many properties with classical stochastic independence and tensor independence. In particular, under the assumption of independence, mixed moments are uniquely determined and can be calculated from marginal moments (also giving rise to an associated convolution product for probability measures on the real line). Another interesting independence is *monotone independence*, which was discovered by Muraki [20]; this is a non-symmetric independence relation.

An extremely useful tool when dealing with random variables which have all moments are the corresponding *cumulants*. The theory of free cumulants, linearizing free

Manuscript received 3rd February 2023, revised 3rd May 2023 and 17th October 2023, accepted 16th January 2024.

KEYWORDS. noncommutative probability, multi-faced independence, cumulants, set partitions.

ACKNOWLEDGEMENTS. The work of both authors was supported by German Research Foundation (DFG) grant no. 397960675, the work of MG also by grant no. 465189426. The work of MG was carried out as a postdoctoral researcher at Saarland University, during the tenure of an ERCIM ‘Alain Bensoussan’ Fellowship Programme at NTNU Trondheim, as a guest researcher at Saarland University in the scope of the SFB-TRR 195, and as a postdoctoral scientific employee at University of Greifswald. The work of PV was partially carried out as a PhD student and scientific employee at University of Greifswald.

additive convolution, was developed by Speicher [26], see also the book by Nica and Speicher [24].⁽¹⁾ Boolean cumulants were formalized by Speicher and Woroudi [28]. Understanding the monotone cumulants took a bit longer, many questions were answered by Hasebe and Saigo [15]. The problem in the monotone case is that independence is not in general characterized by vanishing of mixed cumulants. This is directly related to the non-symmetric nature, as becomes apparent when interpreting moment-cumulant relations via exponential and logarithm maps, as is done in related but different settings by Manzel and Schürmann [18] (Hopf algebraic) or Ebrahimi-Fard and Patras [5] (shuffle-algebraic); non-zero mixed cumulants can appear in the Campbell-Baker-Hausdorff series.

Since the work of Speicher [27], Ben Ghorbal and Schürmann [1], and Muraki [21, 22], we know that the five independence relations for noncommutative random variables, *tensor*, *free*, *Boolean*, *monotone* and *antimonotone* independence, are indeed very special. For these independences, the joint distribution of independent random variables is obtained from the marginal distributions by means of a “universal product”, i.e. a product operation which fulfills a number of natural conditions, including associativity and *universality* (i.e. in a specific sense not dependent on the concrete realization of the noncommutative random variables) and a “factorization for length 2”-condition; and they are the only ones with this property.⁽²⁾ Replacing that “factorization for length 2”-condition by a positivity condition, a decade later, Muraki [23] proved a similar result with a much simpler proof, while at the same time using a much better motivated assumption, namely that the product operation restricts to a product operation for states on augmented \ast -algebras.⁽³⁾ This kind of positivity is also the right condition to study quantum Lévy processes on dual groups in the sense of Ben Ghorbal and Schürmann [2], see also [25], where *Schoenberg correspondence* between convolution semigroups of states and conditionally positive generators is proved in this context. In 2014, Voiculescu [32] introduced a new nontrivial extension of free independence, *bifreeness*, for sequences of pairs of random variables, or *pairs of faces* as Voiculescu called the general underlying framework. Taking up on this idea, more examples of *2-faced* or, more generally, *multi-faced* independences have been discovered [17, 16, 14, 13, 9]. The general theory of *multi-faced universal products* from which those independences can be obtained was established by Manzel and Schürmann [18]. It turned out that not all of the examples fulfill the natural positivity condition. Positivity is still enough to assure Schoenberg correspondence in this generalized setting, see [8]. In an effort to classify positive multi-faced universal products, two routes have been taken. In [10], Gerhold, Hasebe, Ulrich completely classified 2-faced universal products which have a natural representation on the tensor product or the free product Hilbert space of the GNS spaces of the factors. In Varšo’s PhD thesis [30], he proved that there are at most 12 two-faced universal products which fulfill additional assumptions of symmetry and a “combinatorial” moment cumulant relation (i.e. determined by a subset of all two-faced partitions, where more

⁽¹⁾For a single variable, Voiculescu defined free cumulants and proved their uniqueness already in his seminal paper [31].

⁽²⁾Speicher [27] proved that there are only three *universal calculation rules for mixed moments* in the symmetric case. Ben Ghorbal and Schürmann [1] axiomatized independences via universal products and showed equivalence to universal calculation rules. Muraki [21, 22] extended the results to the non-symmetric setting.

⁽³⁾In the purely algebraic context, i.e. without positivity, Muraki’s classification was slightly extended by Gerhold and Lachs in [11], showing that there is a non-symmetric deformation of Boolean independence.

generally weights on two-faced partitions can appear).⁽⁴⁾ In this article we present, simplify, and extend those results of [30].

A single-faced independence can trivially be regarded as a two-faced independence, and every two-faced independence is a certain kind of mixture of two single-faced independences. However, neither do those two single-faced independences determine the two-faced independence, nor is it obvious that any combination of single-faced independences can be combined in any way to form a two-faced independence.⁽⁵⁾ The main result of this article is to present a family of two-faced symmetric universal products such that every positive symmetric two-faced universal product must belong to that family, we call them *candidates*. This is achieved in three steps. First, we prove necessary conditions for a family of weights on ordered partitions to be the highest coefficients of a positive multi-faced universal product (Theorem 5.3); second, we determine all permutation invariant weights (= weights on non-ordered partitions) which fulfill those properties (Corollary 6.11), we call such weights here *admissible*; third, we prove that admissible weights are always the highest coefficients of a (uniquely determined) symmetric multi-faced universal product (Theorem 8.2). The family of candidates consists of (identifying an independence with its underlying universal product, and disregarding the difference between a 2-faced independence and its image under swapping the faces)

- 2-faced continuous 1-parameter deformations of free, tensor and bifree independence (positivity is proved in [10]),
- a tensor-free independence (positivity is not known),
- a new free-free and a new tensor-tensor independence, different from the trivial ones, bifreeness, and their deformations (positivity is not known),
- tensor-Boolean, free-Boolean and Boolean independence; positivity for those is also covered in [10], for free-Boolean it was first shown by Liu [17] and for Boolean independence positivity is of course well-known.

We call the independences which are not realized in [10], i.e. those whose positivity is yet unknown, *exceptional*.

We prove many of the preliminary results for the general symmetric multi-faced case. Theorem 5.3, where we find necessary conditions on weights to arise as highest coefficients of a universal product, is even formulated for not necessarily symmetric products and could be used as a starting point for a more general classification including multi-faced universal products based on monotone independence, such as for example bimonotone independence (of type II) as defined in [9, 13].

It easily follows from the main result that there are no non-trivial positive and symmetric trace preserving universal products (Remark 6.12) and that tensor independence and bifreeness are the only two positive symmetric 2-faced independences which allow to define a convolution of probability measures on \mathbb{R}^2 (Remark 6.13).

Among our additional results, we characterize when a positive symmetric multi-faced universal product is unit preserving (Theorem 9.7), i.e. when it can be defined consistently for arbitrary unital algebras (in the other cases, the product operation is only defined for linear functionals on augmented algebras). This is indeed the case for

⁽⁴⁾In [30], it was also noticed for the first time the possibility that the moment cumulant relation of a positive universal product might not need to be of combinatorial form, which was indeed confirmed in [10] (cumulants are not discussed explicitly in [10], but it is apparent that the universal products obtained as deformations can have non-0-1 highest coefficients).

⁽⁵⁾Note that the study of another kind of *mixture* of single-faced independences was initiated by Młotkowski [19] and received again more attention after work Speicher and Wysocki [29] and Ebrahimi-Fard, Patras and Speicher [6] on the corresponding cumulants; this approach is closely related to *graph products* of groups and the corresponding universal products are not associative binary operations.

the three continuous families and the four (three up to swapping the faces) exceptional cases. Furthermore, we establish a simplified mixed moment formula for the special *combinatorial* case where the highest coefficients are only 0 or 1, so that the moment cumulant relation is simply governed by a specific set of partitions (Theorem 8.4).

The outline of the article is as follows. In Sections 2 to 4, we introduce the basic concepts, in particular multi-faced universal products and multi-faced partitions. In Section 5 we prove the necessary conditions for a family of weights to be the highest coefficients of a positive multi-faced universal product (symmetric or not). In Section 6 we show that those necessary conditions allow us to obtain a concrete list of candidates for symmetric and positive two-faced universal products. In Section 7 we give an introduction to Manzel and Schürmann's cumulant theory, adapted to the relevant special case of symmetric multi-faced independences. In Section 8 we prove, using cumulants, that in the symmetric case the conditions exhibited in Section 5 are sufficient to reconstruct a universal product in the algebraic sense (with a simplified formula in the combinatorial case), but it remains open whether these universal products are automatically positive. Finally, we characterize in Section 9 which universal products in our list are unit preserving. In Section 10 we name four tasks which have to be completed in order to achieve a complete classification of positive multi-faced universal products.

A comparison between this article and corresponding results in Varšo's PhD thesis [30] can be found in Appendix A.

2. PRELIMINARIES AND NOTATION

We will have to deal a lot with tuples of all kinds, so we introduce some useful notation. Let X and Y be arbitrary sets. For any natural number n , denote by $[n]$ the set $\{1, \dots, n\}$. For an n -tuple $\mathbf{t} = (\mathbf{t}(1), \dots, \mathbf{t}(n)) \in X^n$ and a subset $I = \{i_1 < \dots < i_k\} \subset [n]$, we define the *restricted tuple* $\mathbf{t} \upharpoonright I := (\mathbf{t}(i_1), \dots, \mathbf{t}(i_k))$. Two tuples $\mathbf{t} \in X^n, \mathbf{s} \in Y^n$ of the same length may be combined to form the tuple $\mathbf{t} \times \mathbf{s} \in (X \times Y)^n$ with $(\mathbf{t} \times \mathbf{s})(i) = (\mathbf{t}(i), \mathbf{s}(i))$, and conversely, every tuple in $(X \times Y)^n$ is of that form. The set of n -tuples of arbitrary length n is denoted $X^* = \bigcup_{n \in \mathbb{N}_0} X^n$. When a set X does not carry any multiplicative structure, we might use the *word notation*, $\mathbf{t}(1) \cdots \mathbf{t}(n) := (\mathbf{t}(1), \dots, \mathbf{t}(n)) \in X^n$. The entries of a tuple \mathbf{t} might be written t_i instead of $\mathbf{t}(i)$ from time to time; or we might use \mathbf{t} as a shorthand for (t_1, \dots, t_n) without further comment when the t_i have been around before.

An *algebra* means a complex associative algebra, not necessarily unital. The *free product* of algebras A_1, A_2 is denoted $A_1 \sqcup A_2$. Recall that this is the coproduct in the category of algebras: for arbitrary algebra homomorphisms $h_i: A_i \rightarrow B$, there is a unique algebra homomorphism $h_1 \sqcup h_2: A_1 \sqcup A_2 \rightarrow B$ with $h_1 \sqcup h_2 \upharpoonright A_i = h_i$. We use the same symbol \sqcup to denote the canonical homomorphism $h_1 \sqcup h_2: A_1 \sqcup A_2 \rightarrow B_1 \sqcup B_2$ when $h_i: A_i \rightarrow B_i$, it should always be clear from the context which codomain is meant.

For a vector space V , we denote by $T_0(V) = \bigoplus_{n \in \mathbb{N}} V^{\otimes n}$ the (non-unital) free algebra over V . We will identify $T_0(V_1 \oplus V_2) = T_0(V_1) \sqcup T_0(V_2)$ without further commenting. Furthermore, $T_0(V_1) \oplus T_0(V_2)$ is identified with the corresponding subspace of $T_0(V_1 \oplus V_2)$ and linear functionals ψ on $T_0(V_1) \oplus T_0(V_2)$ are identified with linear functionals $T_0(V_1 \oplus V_2)$ by extending them as the 0-functional to the canonical complement, i.e.

$$\psi(v_1 \otimes \cdots \otimes v_k) := \begin{cases} \psi(v_1 \otimes \cdots \otimes v_k) & \text{if } \forall i: v_i \in V_1 \text{ or } \forall i: v_i \in V_2, \\ 0 & \text{if } \exists i, j: v_i \in V_1, v_j \in V_2. \end{cases}$$

In particular, this convention applies to the direct sum of two linear functionals $\psi_i: T_0(V_i) \rightarrow \mathbb{C}$, i.e. we identify $\psi_1 \oplus \psi_2$ with the linear functional on $T_0(V_1 \oplus V_2)$ given by

$$(1) \quad \psi_1 \oplus \psi_2(v_1 \otimes \cdots \otimes v_k) = \begin{cases} \psi_1(v_1 \otimes \cdots \otimes v_k) & \text{if } \forall i: v_i \in V_1, \\ \psi_2(v_1 \otimes \cdots \otimes v_k) & \text{if } \forall i: v_i \in V_2, \\ 0 & \text{if } \exists i, j: v_i \in V_1, v_j \in V_2. \end{cases}$$

The unital free algebra is denoted $T(V) = \bigoplus_{n \in \mathbb{N}_0} V^{\otimes n}$, and this unital algebra is the unitization of $T_0(V)$.

For the rest of this article, if not explicitly mentioned otherwise, \mathcal{F} denotes a fixed finite set, whose elements we call *faces* or *colors*. We could of course assume $\mathcal{F} = [m]$ for $m \in \mathbb{N}$, but since there will be a lot of integers around, we prefer to use more abstract symbols. We mostly use squared symbols such as \square, \blacksquare to denote arbitrary elements of \mathcal{F} . If there are exactly two faces, we assume $\mathcal{F} = \{\circ, \bullet\}$.

A multi-faced (or \mathcal{F} -faced⁽⁶⁾) algebra is an algebra A that is freely generated by given subalgebras $A^\square, \square \in \mathcal{F}$ (the *faces* of A), i.e. the canonical algebra homomorphism $\bigsqcup_{\square \in \mathcal{F}} A^\square \rightarrow A$ is an isomorphism; this is indicated by writing $A = \bigsqcup_{\square \in \mathcal{F}} A^\square$. A *multi-faced algebra homomorphism* is an algebra homomorphism $h: A \rightarrow B$ between multi-faced algebras A, B with $h(A^\square) \subset B^\square$. We consider the free product of multi-faced algebras again a multi-faced algebra with faces $(A \sqcup B)^\square := A^\square \sqcup B^\square$. Note that the free product of multi-faced algebras is the coproduct in the category $\text{Alg}_{\mathcal{F}}$ of multi-faced algebras with multi-faced algebra homomorphisms, i.e. for every pair of multi-faced algebra homomorphisms $h_i: A_i \rightarrow B$ there is a unique multi-faced algebra homomorphism $h_1 \sqcup h_2: A_1 \sqcup A_2 \rightarrow B$ restricting to h_i on A_i , respectively for $i = 1, 2$.

A multi-faced $*$ -algebra is a multi-faced algebra with an involution such that each face is a $*$ -subalgebra. Of course, the free product of multi-faced $*$ -algebras is again a multi-faced $*$ -algebra in the obvious way and the free product of multi-faced $*$ -homomorphisms is a $*$ -homomorphism.

We say that a linear functional $\varphi: A \rightarrow \mathbb{C}$ defined on a multi-faced $*$ -algebra is a *restricted state* if its unital extension to the unitization of A is a state (or, equivalently, positive).

3. UNIVERSAL PRODUCTS

DEFINITION 3.1 (cf. [8, Rem. 3.4]). A multi-faced universal product is a binary product operation for linear functionals on multi-faced algebras (with an a priori fixed set of faces \mathcal{F}) which associates with functionals φ_1, φ_2 on multi-faced algebras A_1, A_2 , respectively, a functional $\varphi_1 \odot \varphi_2$ on $A_1 \sqcup A_2$ such that

- $(\varphi_1 \circ h_1) \odot (\varphi_2 \circ h_2) = (\varphi_1 \odot \varphi_2) \circ (h_1 \sqcup h_2)$ for all multi-faced algebra homomorphisms $h_i: B_i \rightarrow A_i$ (universality)
- $(\varphi_1 \odot \varphi_2) \odot \varphi_3 = \varphi_1 \odot (\varphi_2 \odot \varphi_3)$ (associativity)
- $(\varphi_1 \odot \varphi_2) \upharpoonright A_i = \varphi_i$ (restriction property).

The product is called

- symmetric if $\varphi_1 \odot \varphi_2 = \varphi_2 \odot \varphi_1$,
- positive if the product of restricted states on multi-faced $*$ -algebras is a restricted state on the free product $*$ -algebra.

Note that we made several implicit identifications between isomorphic free products in the last definition. For a more detailed discussion see [8].

⁽⁶⁾We will usually write multi-faced instead of \mathcal{F} -faced. Nevertheless, use of the term always refers to the same fixed set of faces \mathcal{F} .

Universal products have been invented to encode independences. In the single-faced case, this has been worked out by Ben Ghorbal and Schürmann [1]. The multi-faced case is covered by [18] together with the categorical considerations from [7] and [12]. In a nutshell, given a universal product \odot and a linear functional Φ on an algebra \mathcal{A} , algebra homomorphisms $j_\kappa: B_\kappa \rightarrow \mathcal{A}$, $\kappa \in [k]$, defined on multi-faced algebras B_κ , are called \odot -independent w.r.t Φ if

$$\Phi \circ (j_1 \sqcup \cdots \sqcup j_k) = (\Phi \circ j_1) \odot \cdots \odot (\Phi \circ j_k),$$

or, in other words, if the joint distribution of the noncommutative random variables j_κ coincides with the universal product of their marginal distributions. This induces the usual definitions of independence for \mathcal{F} -tuples of elements or of subalgebras of \mathcal{A} . In the remainder of this article, we will not work with the independences themselves, but solely with the underlying universal products, so we refrain from giving more details here.

We will make extensive use of the “Central Structural Theorem” for universal products [18, Theorem 4.2]. Before we present a simplified version of it adapted to the special case of positive multi-faced universal products, we introduce some more notation and give an example.

Let A_1, \dots, A_k be multi-faced algebras and $A = A_1 \sqcup \cdots \sqcup A_k$ (i.e. we identify the A_i with subalgebras of their free product). For $\mathbf{s} = \mathbf{b} \times \mathbf{f} \in ([k] \times \mathcal{F})^n$, we denote

$$A^{\mathbf{s}} := \left\{ a_1 \cdots a_n \in A : a_i \in A_{\mathbf{b}(i)}^{\mathbf{f}(i)} \right\}.$$

Note that the $A^{\mathbf{s}}$ are not necessarily pairwise disjoint.⁽⁷⁾ Elements of $[k]^n$ are referred to as *block structures* and elements of \mathcal{F}^n are called *face structures*.

For $\mathbf{s} = \mathbf{b} \times \mathbf{f} \in ([k] \times \mathcal{F})^n$, put $\beta_\kappa(\mathbf{s}) := \{\ell \in [n] : \mathbf{b}(\ell) = \kappa\}$. We call a set partition π of $[n]$ *adapted to \mathbf{s}* , and write $\pi \prec \mathbf{s}$, if the following two conditions are met:

- each block $\beta \in \pi$ is contained in some $\beta_\kappa(\mathbf{s})$; in other words, π is a refinement of the set partition $\sigma = \{\beta_1(\mathbf{s}), \dots, \beta_k(\mathbf{s})\}$ (to adhere strictly to the usual definition of set partition, empty blocks should be removed from σ)
- if $\mathbf{s}(i) = \mathbf{s}(i+1)$, then $i, i+1$ belong to the same block of π .

Note that, obviously, σ is the maximal partition (w.r.t. refinement order) adapted to \mathbf{s} .

Given a multi-faced universal product \odot , we define its *linearized part* as

$$\varphi_1 \sqcup \cdots \sqcup \varphi_k(a) := \frac{\partial^k}{\partial t_1 \cdots \partial t_k} (t_1 \varphi_1) \odot \cdots \odot (t_k \varphi_k)(a) \Big|_{\mathbf{t}=0}$$

(that this expression is well-defined should be understood as part of the following theorem).

EXAMPLE 3.2. The deformed tensor product $\odot = \otimes \zeta = \zeta \otimes_1 \zeta$ according to [10, Proposition 5.10(1) and Example 5.7]⁽⁸⁾, $\zeta \in \mathbb{T}$, can be calculated for arbitrary 2-faced algebras A_κ , linear functionals $\varphi_\kappa: A_\kappa \rightarrow \mathbb{C}$ ($\kappa \in \{1, 2\}$), and elements $a = a_1^\circ a_2^\circ a_1^\bullet a_2^\bullet \in$

⁽⁷⁾Indeed, if $\mathbf{s}(i) = \mathbf{s}(i+1)$, then $A^{\mathbf{s}} \subseteq A^{\mathbf{s} \setminus \{1, \dots, i-1, i+1, \dots, n\}}$ because $A_{\mathbf{b}(i)}^{\mathbf{f}(i)}$ is a subalgebra of A . A typical way to deal with this is to only consider alternating sequences, i.e. demand $\mathbf{s}(i) \neq \mathbf{s}(i+1)$ for all $i \in [n-1]$. However, it does not cause problems to formulate the subsequent statements for all $A^{\mathbf{s}}$, so we decided to do so.

⁽⁸⁾In the notation of [10], \circ is face (1) and \bullet is face (2).

$A^{1212 \times \infty \bullet \bullet}$ as follows, abbreviating $\langle b \rangle := \varphi_\kappa(b)$ for $b \in A_\kappa$:

$$\begin{aligned} \varphi_1 \odot \varphi_2(a) &= \langle a_1^\circ \rangle \langle a_2^\circ \rangle \langle a_1^\bullet \rangle \langle a_2^\bullet \rangle + (\langle a_1^\circ a_1^\bullet \rangle - \langle a_1^\circ \rangle \langle a_1^\bullet \rangle) \langle a_2^\circ \rangle \langle a_2^\bullet \rangle + \langle a_1^\circ \rangle \langle a_1^\bullet \rangle (\langle a_2^\circ a_2^\bullet \rangle - \langle a_2^\circ \rangle \langle a_2^\bullet \rangle) \\ &\quad + \bar{\zeta} (\langle a_1^\circ a_1^\bullet \rangle - \langle a_1^\circ \rangle \langle a_1^\bullet \rangle) (\langle a_2^\circ a_2^\bullet \rangle - \langle a_2^\circ \rangle \langle a_2^\bullet \rangle) \\ &= \bar{\zeta} \cdot \langle a_1^\circ a_1^\bullet \rangle \langle a_2^\circ a_2^\bullet \rangle + (1 - \bar{\zeta}) \cdot \langle a_1^\circ a_1^\bullet \rangle \langle a_2^\circ \rangle \langle a_2^\bullet \rangle + (1 - \bar{\zeta}) \cdot \langle a_1^\circ \rangle \langle a_1^\bullet \rangle \langle a_2^\circ a_2^\bullet \rangle \\ &\quad - (1 - \bar{\zeta}) \cdot \langle a_1^\circ \rangle \langle a_1^\bullet \rangle \langle a_2^\circ \rangle \langle a_2^\bullet \rangle \end{aligned}$$

Consequently, the linearized part is given by

$$\varphi_1 \sqcap \varphi_2(a) = \bar{\zeta} \cdot \varphi_1(a_1^\circ a_1^\bullet) \varphi_2(a_2^\circ a_2^\bullet).$$

Note how the summands in the full expansion in Example 3.2 correspond to partitions adapted to \mathbf{s} ; the product element a is divided into some sort of “subproducts” which are then evaluated in the appropriate φ_κ . This general pattern is made precise in the following theorem and allows to describe a universal product in terms of the complex coefficients appearing in each summand, which are independent of the involved linear functionals, algebras and algebra elements.

THEOREM 3.3 (Adjusted and simplified from [18, Th. 4.2, Rem. 4.3, 4.4]).

Let \odot be a positive multi-faced universal product and $k \in \mathbb{N}$. Then there are unique coefficients $\alpha_{\mathbf{s}}^\pi$, $\mathbf{s} \in ([k] \times \mathcal{F})^*$, $\pi \prec \mathbf{s}$, such that, for all linear functionals $\varphi_\kappa: A_\kappa \rightarrow \mathbb{C}$ on multi-faced algebras A_κ ($\kappa \in [k]$) and all $a \in A^{\mathbf{s}}$,

$$(2) \quad \varphi_1 \odot \cdots \odot \varphi_k(a) = \sum_{\pi \prec \mathbf{s}} \alpha_{\mathbf{s}}^\pi \cdot \prod_{\kappa \in [k]} \prod_{\substack{\beta \in \pi \\ \beta \subset \beta_\kappa(\mathbf{s})}} \varphi_\kappa \left(\overrightarrow{\prod_{\ell \in \beta} a_\ell} \right).$$

(The symbol $\overrightarrow{\prod}$ indicates that the product is to be taken in the same order as the factors a_j appear in the product $a = a_1 \cdots a_n$.)

Putting $\alpha_{\mathbf{s}} := \alpha_{\mathbf{s}}^\sigma$ (σ the maximal partition adapted to \mathbf{s}), the linearized part is given by

$$(3) \quad \varphi_1 \sqcap \cdots \sqcap \varphi_k(a) = \alpha_{\mathbf{s}} \cdot \varphi_1 \left(\overrightarrow{\prod_{\mathbf{b}(\ell)=1} a_\ell} \right) \cdots \varphi_k \left(\overrightarrow{\prod_{\mathbf{b}(\ell)=k} a_\ell} \right).$$

The $\alpha_{\mathbf{s}}^\pi$ are called coefficients of \odot and the $\alpha_{\mathbf{s}}$ are called highest coefficients of \odot .

Proof. First assume that $\mathbf{s} \in ([k] \times \mathcal{F})^n$ is alternating, i.e. $\mathbf{s}(i) \neq \mathbf{s}(i+1)$ for $i = 1, \dots, n-1$. By [18, Rem. 4.3], the formula given in [18, Th. 4.2] can be applied. For a positive universal product, [18, Rem. 4.4] implies that there is only one summand for each $\pi \prec \mathbf{s}$, corresponding to the “right-ordered coefficient” (i.e. the a_j are multiplied in the same order in which they appear as factors in a) associated with π and \mathbf{s} , denoted $\alpha_{\mathbf{s}}^\pi$ in this article.

If \mathbf{s} is not alternating, then we define $\alpha_{\mathbf{s}}^\pi := \alpha_{\tilde{\mathbf{s}}}^{\tilde{\pi}}$, where $\tilde{\mathbf{s}}$ is the alternating tuple obtained from \mathbf{s} merging repeating entries into one, and $\tilde{\pi}$ the set partition adapted to $\tilde{\mathbf{s}}$ induced by π in the obvious way. By universality it follows that (2) extends to all $\mathbf{s} \in ([k] \times \mathcal{F})^*$; indeed, if $\tilde{\mathbf{s}} = \tilde{\mathbf{b}} \times \tilde{\mathbf{f}}$ has length m and

$$a_1, \dots, a_{r_1} \in A_{\tilde{\mathbf{b}}(1)}^{\tilde{\mathbf{f}}(1)}, \quad a_{r_1+1}, \dots, a_{r_2} \in A_{\tilde{\mathbf{b}}(2)}^{\tilde{\mathbf{f}}(2)}, \quad \dots, \quad a_{r_{m-1}+1}, \dots, a_{r_m} \in A_{\tilde{\mathbf{b}}(m)}^{\tilde{\mathbf{f}}(m)}$$

then for the multi-faced algebras B_κ ($\kappa \in [k]$) which are freely generated by $x_i \in B_{\tilde{\mathbf{b}}(i)}^{\tilde{\mathbf{f}}(i)}$ and multi-faced homomorphisms $h_\kappa: B_\kappa \rightarrow A_\kappa$ defined by

$$B_{\tilde{\mathbf{b}}(i)}^{\tilde{\mathbf{f}}(i)} \ni x_i \mapsto a_{r_{i-1}+1} \cdots a_{r_i} \in A_{\tilde{\mathbf{b}}(i)}^{\tilde{\mathbf{f}}(i)} \quad \text{for } i = 1, \dots, m \quad (r_0 := 0),$$

one finds that, using universality for the first equality,

$$\begin{aligned} \varphi_1 \odot \cdots \odot \varphi_k(a) &= (\varphi_1 \circ h_1) \odot \cdots \odot (\varphi_k \circ h_k)(x_1 \cdots x_m) \\ &= \sum_{\tilde{\pi} \prec \tilde{\mathbf{s}}} \alpha_{\tilde{\mathbf{s}}}^{\tilde{\pi}} \cdot \prod_{\kappa \in [k]} \prod_{\substack{\beta \in \tilde{\pi} \\ \tilde{\beta} \subset \beta_{\kappa}(\tilde{\mathbf{s}})}} (\varphi_{\kappa} \circ h_{\kappa}) \left(\prod_{\tilde{\ell} \in \tilde{\beta}} x_{\tilde{\ell}} \right) \\ &= \sum_{\pi \prec \mathbf{s}} \alpha_{\mathbf{s}}^{\pi} \cdot \prod_{\kappa \in [k]} \prod_{\substack{\beta \in \pi \\ \beta \subset \beta_{\kappa}(\mathbf{s})}} \varphi_{\kappa} \left(\prod_{\ell \in \beta} a_{\ell} \right). \end{aligned}$$

For each $\rho \prec \mathbf{s}$, one can easily construct multi-faced algebras and linear functionals $\varphi_{\kappa}: A_{\kappa} \rightarrow \mathbb{C}$ and an element $a \in A^{\mathbf{s}}$ in such a way that

$$\prod_{\kappa \in [k]} \prod_{\substack{\beta \in \pi \\ \beta \subset \beta_{\kappa}(\mathbf{s})}} \varphi_{\kappa} \left(\prod_{\ell \in \beta} a_{\ell} \right) = \delta_{\pi, \rho},$$

and, thus, $\alpha_{\mathbf{s}}^{\rho} = (\varphi_1 \odot \cdots \odot \varphi_k)(a)$. This shows uniqueness of the coefficients.

Equation (3) follows from Equation (2) because the summand corresponding to the maximal partition σ is the only one which is linear in each φ_{κ} . \square

Obviously, the family of coefficients determines the universal product. In fact, it follows from the cumulant theory developed in [18] that the highest coefficients alone are already enough to determine the universal product. We will come back to this in Section 7.

To end this section, we show that the highest coefficients can be recovered from the linearized part of a universal product using only linear functionals of a particularly well-behaved kind.

DEFINITION 3.4. *A restricted state $\varphi: A \rightarrow \mathbb{C}$ on a multi-faced algebra A is called trivially multi-faced if for all $\square, \blacksquare \in \mathcal{F}$ there exists a $*$ -isomorphism $a^{\square} \mapsto a^{\blacksquare}: A^{\square} \rightarrow A^{\blacksquare}$ with $\varphi(ab^{\square}c) = \varphi(ab^{\blacksquare}c)$ for all $b^{\square} \in A^{\square}$ and all a, c in the unitization of A .*

LEMMA 3.5. *For every $\mathbf{s} = \mathbf{b} \times \mathbf{f} \in ([k] \times \mathcal{F})^*$, there are trivially multi-faced restricted states φ_{κ} on multi-faced $*$ -algebras A_{κ} ($\kappa \in [k]$) and an element $a \in A^{\mathbf{s}}$ with $\varphi_{\kappa}(\prod_{\mathbf{b}(\ell)=\kappa} a_{\ell}) = 1$ for all $\kappa \in [k]$; in particular, for a positive multi-faced universal product \odot it follows that $\alpha_{\mathbf{s}} = \varphi_1 \square \cdots \square \varphi_k(a)$.*

Proof. Define $A_{\kappa}^{\blacksquare} := \mathbb{C}$ and $A_{\kappa} := \bigsqcup_{\blacksquare \in \mathcal{F}} A_{\kappa}^{\blacksquare}$. Then $\varphi_{\kappa} = \bigsqcup_{\blacksquare \in \mathcal{F}} \text{id}: A_{\kappa} \rightarrow \mathbb{C}$ is a state, in particular a restricted state, and trivially multi-faced. Put $a_{\kappa}^{\blacksquare} := 1$ for all $\kappa \in [k]$ and all $\blacksquare \in \mathcal{F}$. Now it is easy to see that $\varphi_{\kappa}(a_{\kappa}^{\blacksquare_1} \cdots a_{\kappa}^{\blacksquare_m}) = 1$ for all $m \in \mathbb{N}$ and $\blacksquare_{\mu} \in \mathcal{F}$ ($\mu \in [m]$). With $a := a_{\mathbf{b}(1)}^{\mathbf{f}(1)} \cdots a_{\mathbf{b}(n)}^{\mathbf{f}(n)}$ the first claim is obvious and the second claim follows from Theorem 3.3. \square

4. PARTITIONS

In general, a *multi-faced set* is a set S together with a map $\mathbf{f}: S \rightarrow \mathcal{F}$, the *face structure* of S . The subsets $S^{\blacksquare} := \mathbf{f}^{-1}(\{\blacksquare\})$ are called the faces of S . A multi-faced subset of S is just a subset of the underlying set viewed as a multi-faced set with respect to the restricted face structure.

In this article, we only deal with multi-faced sets whose underlying set S is finite and totally ordered; these properties are implicitly assumed whenever we write about multi-faced sets in the following.

Any word $\mathbf{f} = \mathbf{f}(1) \cdots \mathbf{f}(n) \in \mathcal{F}^*$ defines face structure on $[n]$, $k \mapsto \mathbf{f}(k)$, (which we identify with the word \mathbf{f}) thus turning $[n]$ into a multi-faced set, denoted by $[n]_{\mathbf{f}}$. Conversely, we associate with a multi-faced set $S = (\{s_1 < \dots < s_n\}, \mathbf{f})$ the word $|\mathbf{f}| := \mathbf{f}(s_1) \cdots \mathbf{f}(s_n) \in \mathcal{F}^*$. We choose this on first sight odd notation because the word \mathbf{f} plays the same role as the number of elements of a set plays in the single-faced case in the moment-cumulant formulas we are aiming at.

Let S be a multi-faced set and \sim an equivalence relation such that

- the equivalence classes are intervals,
- \mathbf{f} is constant on equivalence classes.

Then we understand the quotient S/\sim as a multi-faced set with the induced total order and face map.

EXAMPLE 4.1. We briefly discuss the two situations that will appear several times in this article.

- (1) Let $\mathbf{f} \in \mathcal{F}^n$ be a face and \sim the equivalence relation on $[n]$ that identifies two neighboring points $i, i+1$ in the same face, i.e. $\mathbf{f}(i) = \mathbf{f}(i+1)$. In this case we write $\mathbf{f}/(i \sim i+1)$ for the quotient $[n]_{\mathbf{f}}/\sim$ and denote its elements ℓ instead of $\{\ell\}$ for the trivial equivalence classes of $\ell \in [n] \setminus \{i, i+1\}$ and $\{i, i+1\}$ for the two-element equivalence class of i and $i+1$.
- (2) Let S be a multi-faced set and \sim the equivalence relation whose equivalence classes are the maximal intervals on which \mathbf{f} is constant. We then call the quotient $S_{\text{red}} := S/\sim$ the *reduction* of S . In the reduction, neighboring points will always have different faces, so that no further quotienting is possible.

A partition of a multi-faced set S is a collection of multi-faced subsets whose underlying sets form a set partition. The set of all partitions of a multi-faced set S is denoted $\mathcal{P}(S)$. An ordered partition of S is a partition of S together with a total order between the blocks. The set of all ordered partitions is denoted $\mathcal{P}_{<}(S)$.

For a word $\mathbf{f} \in \mathcal{F}^n$, we put $\mathcal{P}(\mathbf{f}) := \mathcal{P}([n]_{\mathbf{f}})$ and $\mathcal{P}_{<}(\mathbf{f}) := \mathcal{P}_{<}([n]_{\mathbf{f}})$. We also denote

$$\mathcal{P} := \bigcup_{\mathbf{f} \in \mathcal{F}^*} \mathcal{P}(\mathbf{f}), \quad \mathcal{P}_{<} := \bigcup_{\mathbf{f} \in \mathcal{F}^*} \mathcal{P}_{<}(\mathbf{f}).$$

EXAMPLE 4.2. Let $\mathcal{F} = \{\circ, \bullet\}$ and consider $\mathbf{f} = \circ\bullet\circ\bullet \in \mathcal{F}^*$. Then $\pi = \{\beta_1, \beta_2\}$ with $\beta_1 = \{1, 3, 4\}, \beta_2 = \{2, 5\}$ is an element of $\mathcal{P}(\mathbf{f})$ and we have $|\beta_1| = \circ\bullet\circ, |\beta_2| = \bullet\bullet$. This can be nicely drawn as an *arc diagram*, $\pi = \begin{array}{c} \circ & \bullet & \circ & \bullet & \circ \\ & \text{---} & & \text{---} & \\ & \text{---} & & \text{---} & \end{array}$.

In the following we will not distinguish between a partition and its arc diagram. In this article, we mostly use arc-diagrams to denote partitions in \mathcal{P} , i.e. without a block-order; the height of the blocks is then completely arbitrary. For a partition in $\mathcal{P}_{<}$, the height of the block corresponds to the order between blocks. If the underlying set S is not of the form $[n]_{\mathbf{f}}$ (typically because it was obtained as a quotient), we draw the diagram for the corresponding partition of $|\mathbf{f}|$.

$\mathcal{P}(\mathbf{f})$ is a partially ordered set by the order of reverse refinement. The maximum and minimum of $\mathcal{P}(\mathbf{f})$ are denoted $1_{\mathbf{f}}$ and $0_{\mathbf{f}}$, respectively, i.e. $1_{\mathbf{f}}$ is the one-block partition and in $0_{\mathbf{f}}$ all blocks are singletons.

There is a canonical bijection between $\mathcal{P}(S/\sim)$ and the set of $\pi \in \mathcal{P}(S)$ such that equivalent points of S lie in the same block of π .

For a multi-faced partition π , consider the equivalence relation \sim defined on the underlying multi-faced set S by

$$s \sim t : \Longleftrightarrow \text{all } r \in S \text{ with } s \leq r \leq t \text{ have the same color and belong to the same block of } \pi.$$

In other words, \sim is the equivalence relation whose equivalence classes are the maximal intervals I of S which fulfill the following two properties:

- \mathbf{f} is constant on I ;
- all elements of I belong to the same block of π .

We define the *reduction* of π as the induced multi-faced partition π_{red} on S/\sim . For example,

$$\left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right)_{\text{red}} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}.$$

Then π_{red} will not have neighboring legs that are in the same face and in the same block. For $\pi \in \mathcal{P}_{<}$, the block order remains unchanged.

For a multi-faced set S , we define its mirror image \overline{S} as the set with one element \overline{s} for each $s \in S$ (so that $s \mapsto \overline{s}$ is a bijection) with the face structure $\overline{\mathbf{f}}(\overline{s}) := \mathbf{f}(s)$ and reversed order, i.e. $\overline{s} \leq \overline{t} \iff s \geq t$. For $\pi \in \mathcal{P}(S)$, we put $\overline{\pi} \in \mathcal{P}(\overline{S})$ as the set partition with a block $\overline{\beta} = \{\overline{s}_1, \dots, \overline{s}_n\}$ for each block $\beta = \{s_1, \dots, s_n\} \in \pi$. For example,

$$\overline{\circ \bullet \bullet \circ} = \bullet \circ \bullet \circ, \quad \overline{\left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right)} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}.$$

If $S = [n]_{\mathbf{f}}$ for $\mathbf{f} \in \mathcal{F}^n$, so that the underlying set is $[n]$, we use the convention that $\overline{k} := n - k + 1$ (i.e. we identify \overline{k} with its image under the unique strictly increasing map $\overline{[n]} \rightarrow [n]$); this has the effect that $\overline{[n]}$ is identified with $[n]$ and $\overline{[n]_{\mathbf{f}}} = [n]_{\overline{\mathbf{f}}}$ for $\overline{\mathbf{f}} = \overline{\mathbf{f}}(1) \cdots \overline{\mathbf{f}}(n) = \mathbf{f}(n) \cdots \mathbf{f}(1)$ the mirror image of \mathbf{f} . This is clearly in accordance the diagrammatic representation. If $\pi = \{\beta_1 < \dots < \beta_k\} \in \mathcal{P}_{<}$, then $\overline{\pi}$ is defined as before together with the (non-reversed!) block order $\overline{\beta}_1 < \dots < \overline{\beta}_k$.

Finally, we introduce a notation for uniting blocks. Let $\pi = \{\beta_1 < \dots < \beta_k\} \in \mathcal{P}_{<}(S)$ with blocks β_i, β_{i+1} that are nearest neighbors for the order on π . Then we define $\pi_{\beta_i \cup \beta_{i+1}} := \{\beta_1 < \dots < \beta_{i-1} < \beta_i \cup \beta_{i+1} < \dots < \beta_k\}$. Similarly, for $\pi \in \mathcal{P}(\mathbf{f})$ and arbitrary blocks $\beta_1, \beta_2 \in \pi$, $\pi_{\beta_1 \cup \beta_2} := \pi \setminus \{\beta_1, \beta_2\} \cup \{\beta_1 \cup \beta_2\}$. For example,

$$\left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right)_{\{1,3\} \cup \{2,5\}} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}.$$

Let $\mathbf{f}_i \in \mathcal{F}^{m_i}$, $i \in [n]$, be face structures and \mathbf{f} their *concatenation*, i.e. $\mathbf{f}(m_1 + \dots + m_{i-1} + \ell) = \mathbf{f}_i(\ell)$ for all $i \in [n]$, $\ell \in [m_i]$. Given partitions $\pi_i \in \mathcal{P}(\mathbf{f}_i)$, we define their *concatenation* as the partition $\pi \in \mathcal{P}(\mathbf{f})$ which has for every block $\beta \in \pi_i$ with $i \in [n]$ a block $\tilde{\beta} := \{\ell : \ell + \sum_{j=1}^{i-1} m_j \in \beta\}$. Roughly speaking, π restricts to π_i on the legs corresponding to \mathbf{f}_i . For example, the concatenation of $\pi_1 = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$ and $\pi_2 = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$ is $\pi = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$. We do not define here the concatenation of ordered partitions.

5. HIGHEST COEFFICIENTS: NECESSARY CONDITIONS

DEFINITION 5.1. A family of complex numbers $\alpha = (\alpha_{\pi})_{\pi \in \mathcal{P}_{<}}$ is called (family of) weights on ordered partitions, a family $\alpha = (\alpha_{\pi})_{\pi \in \mathcal{P}}$ is called (family of) weights on partitions. Weights on (ordered) partitions are called monic if $\alpha_{\pi} = 1$ for every one-block partition.

For a family of numbers

$$\alpha_{\mathbf{s}} : \mathbf{s} \in ([k] \times \mathcal{F})^n, k, n \in \mathbb{N}$$

(as it is for example obtained from a universal product by Theorem 3.3) and $\pi = \{\beta_1 < \dots < \beta_k\} \in \mathcal{P}_{<}(\mathbf{f})$ an ordered multi-faced partition with k blocks, we define $\mathbf{s}_{\pi} \in ([k] \times \mathcal{F})^n$ via $\mathbf{s}_{\pi}(\ell) := (\kappa, \blacksquare)$ if $\ell \in \beta_{\kappa}$ and $\mathbf{f}(\ell) = \blacksquare$ and put

$$\alpha_{\pi} := \alpha_{\mathbf{s}_{\pi}}.$$

In this way, we associate with each universal product a family of weights on ordered partitions, and we say that the weights of a universal product are its highest coefficients. Note that such weights are always monic.

We say that weights on ordered partitions α are *invariant under permutation of blocks* if

$$\alpha_{\{\beta_1 < \dots < \beta_k\}} = \alpha_{\{\beta_{1'} < \dots < \beta_{k'}\}} \text{ for every permutation } \kappa \mapsto \kappa' \text{ of } [k].$$

In this case, define α_π for a non-ordered partition $\pi = \{\beta_1, \dots, \beta_k\} \in \mathcal{P}(\mathbf{f})$ simply as the value $\alpha_{\{\beta_1 < \dots < \beta_k\}}$ for an arbitrary ordered partition with the same blocks as π . In this way, we can identify weights on partitions and weights on ordered partitions which are invariant under block permutation.

REMARK 5.2. It is easy to check that the weights α coming from a universal product according to Theorem 3.3 are invariant under permutation of blocks if and only if the universal product is symmetric.

The question we wish to answer is the following: under which conditions on the weights α is there a (positive) universal product \odot with highest coefficients α ? The next theorem yields some necessary conditions.

THEOREM 5.3. *Let \odot be a positive multi-faced universal product. Then the highest coefficients fulfill:*

- (i) $\alpha_{1\mathbf{f}} = 1$ for all $\mathbf{f} \in \mathcal{F}^*$.
- (ii) $\alpha_{(\{1\} < \{2\}, \mathbf{f})} = \alpha_{(\{2\} < \{1\}, \mathbf{f})} = 1$ for every $\mathbf{f} \in \mathcal{F}^2$.
- (iii) $\alpha_\pi = \alpha_{\text{red}(\pi)}$.
- (iv) Suppose $\pi \in \mathcal{P}_<(\mathbf{f})$ has blocks $\beta_1 < \beta_2$ that are nearest neighbors for the order of π and have neighboring legs in the same face, i.e. there exist $i \in \beta_1, j \in \beta_2, |i - j| = 1, \mathbf{f}(i) = \mathbf{f}(j)$. Then

$$\alpha_\pi = \alpha_{\pi_{\beta_1 \sim \beta_2}} \cdot \alpha_{\{\beta_1 < \beta_2\}}$$

- (v) $\alpha_\pi = \alpha_\sigma$ whenever π and σ only differ in the faces of extremal legs.
- (vi) $\alpha_\pi = \overline{\alpha_\pi}$.

Proof. Recall the definition of $\mathbf{s}_\pi \in ([k] \times \mathcal{F})^*$ for $\pi = (\beta_1 < \dots < \beta_k) \in \mathcal{P}_<$ from the beginning of this section. By Lemma 3.5, we can express each coefficient α_π as

$$\alpha_\pi = \varphi_1 \square \dots \square \varphi_k(a)$$

with $a \in A^\pi := A^{\mathbf{s}_\pi}$ and $(\varphi_1 \otimes \dots \otimes \varphi_k)(a_\pi) = 1$, where $a_\pi := (\vec{\prod}_{\ell \in \beta_1} a_\ell) \otimes \dots \otimes (\vec{\prod}_{\ell \in \beta_k} a_\ell)$. We will freely use this notation in the rest of the proof.

(i) follows from the restriction property in Definition 3.1. (iii) holds by definition of the non-reduced coefficients in the proof of Theorem 3.3. For (iv) we have to carefully analyse the linearized universal product. If π has neighboring blocks $\beta_1 < \beta_2$ with neighboring legs in face $\blacksquare \in \mathcal{F}$, then $a \in A^\pi$ implies that $a = a_1 \dots a_r a_{r+1} \dots a_n$ with $a_r \in A_i^\blacksquare, a_{r+1} \in A_j^\blacksquare$ with $|i - j| = 1$. Without loss of generality, assume $j = i + 1$. Then

$$\begin{aligned} \alpha_\pi &= \varphi_1 \square \dots \square \varphi_k(a) \\ &= \frac{\partial^k}{\partial t_1 \dots \partial t_k} \left((t_1 \varphi_1) \odot \dots \odot ((t_i \varphi_i) \odot (t_{i+1} \varphi_{i+1})) \odot \dots \odot (t_k \varphi_k) \right) (a) \Big|_{\mathbf{t}=0}. \end{aligned}$$

Evaluating the full coefficient formula, Equation (2), for the universal product of the $k - 1$ functionals

$$\psi_\ell := \begin{cases} t_\ell \varphi_\ell & \text{if } \ell < i, \\ t_i \varphi_i \odot t_{i+1} \varphi_{i+1} & \text{if } \ell = i, \\ t_{\ell+1} \varphi_{\ell+1} & \text{if } \ell > i \end{cases}$$

every summand will contain a factor $F = \psi_i(\cdots a_r a_{r+1} \cdots)$ because the two factors a_r, a_{r+1} are from the same block and face and therefore have to be treated as one. Summands with more factors containing ψ_i vanish in the linearization procedure. Therefore, we obtain

$$\begin{aligned} & \frac{\partial^k}{\partial t_1 \cdots \partial t_k} \left((t_1 \varphi_1) \odot \cdots \odot ((t_i \varphi_i) \odot (t_{i+1} \varphi_{i+1})) \odot \cdots \odot (t_k \varphi_k) \right) (a) \Big|_{\mathbf{t}=0} \\ &= \alpha_{\pi_{\beta_1 \cup \beta_2}} \frac{\partial^2}{\partial t_i \partial t_{i+1}} ((t_i \varphi_i) \odot (t_{i+1} \varphi_{i+1})) (\cdots a_r a_{r+1} \cdots) \Big|_{\mathbf{t}=0} \\ &= \alpha_{\pi_{\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1 < \beta_2\}} \end{aligned}$$

as claimed

So far, we have not made significant use of positivity (except that we assumed that wrong ordered coefficients vanish), but positivity is important to prove the remaining two properties.

(vi) follows easily from the fact that positive functionals are hermitian and $a \in A^\pi$ if and only if $a^* \in A^{\bar{\pi}}$. All we have to do is choose some restricted states $\varphi_1, \dots, \varphi_k$ and $a \in A^\pi$ with $\varphi_1 \otimes \cdots \otimes \varphi_k(a_\pi) = 1$, then $\varphi_1 \otimes \cdots \otimes \varphi_k((a^*)_{\bar{\pi}}) = 1$ and we conclude $\overline{\alpha_\pi} = \varphi_1 \boxdot \cdots \boxdot \varphi_k(a^*) = \alpha_{\bar{\pi}}$.

To show (v), assume that $\alpha_\pi = \varphi_1 \boxdot \cdots \boxdot \varphi_k(a)$, $a = a_1^\square a_2 \cdots a_n$ with $a_1^\square \in A_i^\square$ and trivially multi-faced restricted states φ_κ , this is always possible by Lemma 3.5. Then $t_\kappa \varphi_\kappa$ is a restricted state for all $t_\kappa \leq 1$, and

$$\begin{aligned} & |t_1 \varphi_1 \odot \cdots \odot t_k \varphi_k((a_1^\square - a_1^\blacksquare) a_2 \cdots a_n)|^2 \\ & \leq t_i \varphi_i((a_1^\square - a_1^\blacksquare)^*(a_1^\square - a_1^\blacksquare)) (t_1 \varphi_1 \odot \cdots \odot t_k \varphi_k)((a_2 \cdots a_n)^*(a_2 \cdots a_n)) = 0 \end{aligned}$$

where a_1^\blacksquare is the image of a_1^\square under the isomorphism $A_i^\square \cong A_i^\blacksquare$ making φ_i trivially multi-faced. From this the statement for the first leg readily follows. For the corresponding statement for the last leg, we can either apply (vi) or perform an analogous computation.

Finally, let $\pi = (\{\{1\} < \{2\}\}, \mathbf{f})$ or $\pi = (\{\{2\} < \{1\}\}, \mathbf{f})$. By (v), we can assume without loss of generality that $\mathbf{f}(1) = \mathbf{f}(2)$. Therefore, (ii) follows from the single-faced case, which is settled in [2, Theorem 2.5]⁽⁹⁾. \square

REMARK 5.4. Note that the multi-faced universal products of bi-Boolean independence (defined by Gu and Skoufranis [14]) and bi-monotone independence of type I (defined by Gu, Hasebe and Skoufranis [13]) are not positive. Their associated highest coefficients do not fulfill (vi).

For the rest of this article, we restrict ourselves to the symmetric case. As noted before, symmetry of the universal product is equivalent to invariance under block-permutation of its highest coefficients, and in this case we denote its highest coefficients α_π with $\pi \in \mathcal{P}$.

DEFINITION 5.5. A family $\alpha = (\alpha_\pi)_{\pi \in \mathcal{P}}$ of complex numbers is called admissible weights if the corresponding block-permutation invariant family $\alpha_{(\pi, <)} := \alpha_\pi$ fulfills (i) – (vi) in Theorem 5.3; in particular, it fulfills

(iv') Suppose $\pi \in \mathcal{P}(\mathbf{f})$ has blocks $\beta_1 \neq \beta_2$ with neighboring legs $i \in \beta_1$, $i+1 \in \beta_2$ of the same face, $\mathbf{f}(i) = \mathbf{f}(i+1)$. Then

$$\alpha_\pi = \alpha_{\pi_{\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1, \beta_2\}}.$$

⁽⁹⁾In the statement, Ben Ghorbal and Schürmann assume “nondegenerateness”, but the proof does not use this assumption.

DEFINITION 5.6. A set of multi-faced partitions $\Pi \subset \mathcal{P}$ is called an admissible set of partitions if

$$\alpha_\pi := \begin{cases} 1 & \text{if } \pi \in \Pi, \\ 0 & \text{otherwise} \end{cases}$$

defines admissible weights.⁽¹⁰⁾

NOTATION 5.7. If π is described by a certain arc-diagram Diag , we will write $\alpha(\text{Diag})$ instead of α_π . Also, we will use a grey square (or circle in the 2-faced case) to indicate that the color of the extremal legs is arbitrary. For example $\alpha\left(\begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array}\right) = \alpha_\pi$ for $\pi = (\{1, 6\}, \{2, 4\}, \{3, 5\}, \square\square\square\square\square\square)$ or any other π with the same set partition and the same coloring of the non-extremal legs 2,3,4,5.

OBSERVATION 5.8. Let $(\alpha_\pi)_{\pi \in \mathcal{P}}$ be admissible weights. Then $\Pi_\alpha = \{\pi : \alpha_\pi \neq 0\}$ is an admissible set of partitions. There are, however, admissible families with $\alpha_\pi \notin \{0, 1\}$ for some $\pi \in \mathcal{P}$. Indeed, Example 3.2 in particular shows that, for α the highest coefficients of the deformed tensor product \otimes^ζ with $\zeta \neq 1$, one finds $\alpha(\begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \end{array}) = \bar{\zeta} \notin \{0, 1\}$.

OBSERVATION 5.9. A set $\Pi \subset \mathcal{P}$ of partitions is admissible if and only if Π contains the partitions

(P-i) $1_{\mathbf{f}}$ for all $\mathbf{f} \in \mathcal{F}^*$

(P-ii) $\begin{array}{c} \text{---} \\ \text{---} \end{array}$ for all $\square \in \mathcal{F}^2$

and is closed under the following operations used in [30]:

(P-iii) double a leg, including its color

(P-iii)' merge two neighboring legs of the same color in the same block into one

(P-iv) unite two blocks which have neighboring legs of the same color into one block,

$$\pi \mapsto \pi_{\beta_1 \cup \beta_2}$$

(P-iv)' remember a two-block partition formed by two blocks with neighboring legs of the same color, $\pi \mapsto \{\beta_1, \beta_2\}$

(P-iv)" replace a block of a partition from Π by a two-block partition from Π (of the same underlying multi-faced set as the original block) such that the blocks have neighboring legs of the same color, $(\pi_{\beta_1 \cup \beta_2}, \{\beta_1, \beta_2\}) \mapsto \pi$

(P-v) mirror a partition, $\pi \mapsto \bar{\pi}$

(P-vi) change color of an extremal leg of a partition from Π

Given any partitions $\pi_1, \dots, \pi_n \in \mathcal{P}$, we denote by $\langle \pi_1, \dots, \pi_n \rangle$ the minimal admissible set of partitions that contains all π_i . We say that $\langle \pi_1, \dots, \pi_n \rangle$ is generated by π_1, \dots, π_n ; note that $\langle \pi_1, \dots, \pi_n \rangle$ indeed consists of those partitions in \mathcal{P} which can be obtained in finitely many steps by applying the operations of Observation 5.9 to the partitions $1_{\mathbf{f}}$ ($\mathbf{f} \in \mathcal{F}^*$), $\begin{array}{c} \text{---} \\ \text{---} \end{array}$ ($\square \in \mathcal{F}^2$), and π_1, \dots, π_n .

6. PARTIAL CLASSIFICATION OF SYMMETRIC POSITIVE INDEPENDENCES

In this section we determine all admissible families $(\alpha_\pi)_{\pi \in \mathcal{P}}$.

DEFINITION 6.1. Let π be a partition.

- A leg ℓ is called inner if there exist legs $i < \ell < j$ and a block $\beta \in \pi$ with $i, j \in \beta$ and $\ell \notin \beta$. Otherwise it is called outer.

⁽¹⁰⁾Note that this is closely related to the definition of a universal class of partitions in [30], but not completely equivalent; the difference is that an admissible set must always contain the partitions $\begin{array}{c} \text{---} \\ \text{---} \end{array}$ for all $\square \in \mathcal{F}^2$.

- Two legs ℓ, ℓ' are called connected if they lie in the same block or if there is a sequence of blocks $\ell \in \beta_1, \dots, \beta_n \ni \ell'$ such that there is a crossing between β_k and β_{k+1} . Roughly speaking, ℓ and ℓ' are connected if and only if one can move from ℓ to ℓ' going only along the lines of the diagram associated with π .

We start by describing some simple consequences of the defining properties of admissible families of coefficients.

LEMMA 6.2. *Let $(\alpha_\pi)_{\pi \in \mathcal{P}}$ be admissible weights.*

- (1) $\alpha_\pi = 1$ for all interval partitions π .
- (2) Let π be the concatenation of π_1, \dots, π_n . Then $\alpha_\pi = \prod \alpha_{\pi_i}$.
- (3) $\alpha_\pi = \alpha_\sigma$ when σ is obtained by replacing one leg ℓ by two copies and splitting the block $\beta \ni \ell$ into β_1 and β_2 , where β_1 contains the first copy and all legs of β smaller than ℓ and β_2 contains the second copy and all legs of β larger than ℓ . We say that σ is obtained by splitting β at ℓ .
- (4) $\alpha_\pi = \alpha_\sigma$ when σ is obtained replacing an arbitrary number of connected outer legs by a single outer leg of arbitrary color. We call this process collapsing the outer legs.

Proof.

- (1) This is easily proved by induction. For a two-block interval partition π , we can consecutively change color of the extremal legs and merge them with their neighboring legs until we reach $\alpha_\pi = \alpha \left(\begin{smallmatrix} | & | \\ \hline | & | \end{smallmatrix} \right) = 1$. It is worth noting that for this step we needed to change the color of both extremal legs.

Assume that the statements holds for $(n-1)$ -block interval partitions and let $\pi = (\{\beta_1, \dots, \beta_n\}, \mathbf{f})$ be an interval partition with $n > 2$ blocks. Starting similar as before, we can without loss of generality assume that $\beta_1 = \{1\}$ and $\mathbf{f}(1) = \mathbf{f}(2)$. In that case, we find that $\alpha_\pi = \alpha_{\pi_{\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1, \beta_2\}} = 1$.

- (2) Clearly, it is enough to prove the claim for $n = 2$. We prove the claim by induction on the number of blocks $|\pi|$. If $|\pi| = 2$, then $|\pi_1| = |\pi_2| = 1$ and the three partitions are interval partitions, in particular $\alpha_\pi = 1 = \alpha_{\pi_1} \alpha_{\pi_2}$. If $|\pi| > 2$, then $|\pi_1| > 1$ or $|\pi_2| > 1$. In case $|\pi_1| > 2$, let $1 \in \beta_1 \in \pi_1$. We can assume without loss of generality that $2 \in \beta_2$ belongs to a different block $\beta_1 \neq \beta_2 \in \pi_1$ and $\mathbf{f}(1) = \mathbf{f}(2)$; if those conditions are not met, it does not change the coefficient to change the color of the first leg to match the color of the second leg and merge them into one until we are in the described situation. Now we find $\alpha_{\pi_1} = \alpha_{\pi_{1\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1, \beta_2\}}$ and $\alpha_\pi = \alpha_{\pi_{\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1, \beta_2\}}$. Of course, $|\pi_{\beta_1 \cup \beta_2}| = |\pi| - 1$, so we may assume that the statement holds for $\pi_{\beta_1 \cup \beta_2}$ which is the concatenation of $\pi_{1\beta_1 \cup \beta_2}$ and π_2 . Altogether,

$$\alpha_\pi = \alpha_{\pi_{\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1, \beta_2\}} = \alpha_{\pi_{1\beta_1 \cup \beta_2}} \cdot \alpha_{\pi_2} \cdot \alpha_{\{\beta_1, \beta_2\}} = \alpha_{\pi_1} \alpha_{\pi_2}.$$

If $|\pi_2| > 1$, we argue analogously, but we have to change the color of the last leg.

- (3) We have $\alpha_\pi = \alpha_\sigma \alpha_{\{\beta_1, \beta_2\}}$, and $\alpha_{\{\beta_1, \beta_2\}} = 1$ by (1).
- (4) Decompose π into a concatenation of *irreducible* π_1, \dots, π_n , i.e. no π_i can be deconcatenated any further. By Item 2, $\alpha_\pi = \prod \alpha_{\pi_i}$. Note that every outer leg of π is the outer leg of some π_i and that connected outer legs are necessarily in the same block. For each π_i , the outer legs can be collapsed by iteratively changing the face of the first or last leg to match the face of its successor or predecessor, respectively, and merging the legs using the fact that the weights don't change when we reduce the partition (Condition (iii) in Theorem 5.3). After collapsing the outer legs that way, the faces of the

outer legs can be changed once more in such a way that the concatenation of the obtained partitions σ_i is σ . It follows, using again Item 2, that $\alpha_\sigma = \prod \alpha_{\sigma_i} = \prod \alpha_{\pi_i} = \alpha_\pi$. \square

It is worth noting that, in the proof of Item 2, we need invariance of the coefficients under changing the faces of both extremal legs. For example, the weights associated with bi-Boolean independence defined in [14] do not share this property.

LEMMA 6.3. *Two admissible families coincide if and only if they coincide on 2-block partitions.*

Proof. Assume that $(\alpha_\pi), (\beta_\pi)$ are admissible families with $\alpha_\sigma = \beta_\sigma$ for all 2-block partitions σ . By definition, the value on 1-block partitions is 1. Given an n -block partition π with $n > 2$, we alternately

- change the color of the first leg to match the color of the second leg, cf. (v),
- combine the first two legs into one if they belong to the same block, cf. (iii),

to obtain a partition $\tilde{\pi}$ such that the first two legs of π have the same color but belong to different blocks β_1, β_2 . Then $\alpha_\pi = \alpha_{\tilde{\pi}}, \beta_\pi = \beta_{\tilde{\pi}}$ by definition of admissible weights. Using (iv), we then have $\alpha_\pi = \alpha_{\pi_{\beta_1 \cup \beta_2}} \cdot \alpha_{\{\beta_1, \beta_2\}}$ and $\beta_\pi = \beta_{\pi_{\beta_1 \cup \beta_2}} \cdot \beta_{\{\beta_1, \beta_2\}}$, where $\pi_{\beta_1 \cup \beta_2}$ is an $(n-1)$ -block partition and $\{\beta_1, \beta_2\}$ is a 2-block partition. We can iterate the procedure until we obtain α_π, β_π as products of coefficients of the same sequence of 2-block partitions, thus proving the claim. \square

COROLLARY 6.4. *Two admissible families coincide if and only if they coincide on 2-block partitions of at most four legs.*

Proof. Suppose that $\pi = \{\beta, \gamma\}$ has more than 4 legs and that the third leg lies in β . Without loss of generality, we assume that the first leg and the second leg belong different blocks but the same face; if they would belong to the same block, they could be collapsed and the face of the first leg can simply be adapted to that of the second leg without changing the coefficient. Without loss of generality assume that $1 \in \beta$. If all legs after the third leg belong to γ , they are necessarily outer and can be collapsed to reach a partition with four legs. If there is at least one leg from β after the third leg, then splitting β at the third leg yields a partition $\tilde{\pi} = \{\beta_1, \beta_2, \gamma\}$ where $\beta_1 = \{1, 3\}$ has two legs and $\beta_2 = \{3'\} \cup (\beta \setminus \{1, 3\})$ has exactly one leg less than β ; here $3'$ is the copy of 3 obtained from splitting such that $3 < 3'$. Now, $\alpha_\pi = \alpha_{\tilde{\pi}} = \alpha_{\tilde{\pi}_{\beta_1 \cup \gamma}} \alpha_{\{\beta_1, \gamma\}}$. Obviously, $\tau := \{\beta_1, \gamma\}$ has strictly less legs than π . Since the first three legs of $\tilde{\pi}_{\beta_1 \cup \gamma}$ belong to the same block, after collapsing those three legs, we get a partition σ with $\alpha_\sigma = \alpha_{\tilde{\pi}_{\beta_1 \cup \gamma}}$ which has one leg less than π (one leg more from the splitting are overcompensated by two legs less from collapsing). All in all, $\alpha_\pi = \alpha_\tau \alpha_\sigma$, where both, τ and σ are two-block partitions with a strictly smaller number of legs than π . This procedure can be iterated until α_π is expressed as a product of only 2-block partitions with at most 4 legs. \square

DEFINITION 6.5. *We introduce shorthand notations for the basic coefficients, where $\square, \blacksquare \in \mathcal{F}$:*

$$\nu_\square := \alpha \left(\begin{array}{|c|} \hline \square \\ \hline \end{array} \right), \quad \xi_\square := \alpha \left(\begin{array}{|c|} \hline \square \square \\ \hline \end{array} \right), \quad \nu_{\blacksquare} := \alpha \left(\begin{array}{|c|} \hline \blacksquare \\ \hline \end{array} \right), \quad \xi_{\blacksquare} := \alpha \left(\begin{array}{|c|} \hline \blacksquare \blacksquare \\ \hline \end{array} \right)$$

(Note that $\xi_{\square\square} = \xi_\square$, obviously, and $\nu_{\square\square} = \nu_\square$, because we can merge neighboring legs of the same face.)

COROLLARY 6.6. *Two admissible families coincide if and only if they have the same basic coefficients.*

Proof. A two-block partition with at most four legs is either an interval partition (in which case its coefficient is 1) or it can be reduced by changing color and combining legs to one of the partitions that define the basic coefficients. \square

LEMMA 6.7. *For all $\square, \blacksquare \in \mathcal{F}$, we have the following relations between the basic coefficients:*

- (1) $\nu_{\square}^2 = \nu_{\square}, \xi_{\square}^2 = \xi_{\square}$, i.e. $\nu_{\square}, \xi_{\square} \in \{0, 1\}$,
- (2) $t\nu_{\square} = t$ for $t \in \{\nu_{\square\blacksquare}, \xi_{\square}, \xi_{\square\blacksquare}\}$, i.e. $\nu_{\square} = 0 \implies \nu_{\square\blacksquare} = \xi_{\square} = \xi_{\square\blacksquare} = 0$,
- (3) $|t|^2 t = t$ for $t \in \{\nu_{\square\blacksquare}, \xi_{\square\blacksquare}\}$, i.e. $\nu_{\square\blacksquare}, \xi_{\square\blacksquare} \in \{0\} \cup \mathbb{T}$,
- (4) $\nu_{\square\blacksquare}\xi_{\square} = \xi_{\square\blacksquare}\xi_{\square}$, i.e. $\xi_{\square} = 1 \implies \nu_{\square\blacksquare} = \xi_{\square\blacksquare}$,
- (5) $\nu_{\square\blacksquare}\xi_{\square\blacksquare} = \nu_{\square\blacksquare}\xi_{\square\blacksquare}\xi_{\square}$, i.e. $\xi_{\square} = 0 \implies \nu_{\square\blacksquare} = 0$ or $\xi_{\square\blacksquare} = 0$.

Proof.

- (1) This follows as in the single-faced case, see [27]. Alternatively, this follows easily as a special case $\square = \blacksquare$ from the items below.
- (2) Consider $\begin{array}{|c|} \hline \square \\ \hline \blacksquare \\ \hline \end{array}$. Split the inner \square -leg and merge it's copy with the outer block to obtain $\alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right)\alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right)$. The other cases work analogously.
- (3) First note that $\alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = |\nu_{\square\blacksquare}|^2$. This leads to

$$|\nu_{\square\blacksquare}|^2 \nu_{\square\blacksquare} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) \nu_{\square\blacksquare} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \nu_{\square\blacksquare} \nu_{\square} = \nu_{\square\blacksquare}.$$

Similarly, $\alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = |\xi_{\square\blacksquare}|^2$ and hence

$$|\xi_{\square\blacksquare}|^2 \xi_{\square\blacksquare} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) \xi_{\square\blacksquare} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \xi_{\square\blacksquare} \nu_{\square} = \xi_{\square\blacksquare}.$$

- (4) This follows from

$$\nu_{\square\blacksquare}\xi_{\square} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) \nu_{\square\blacksquare} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) \nu_{\square} = \xi_{\square\blacksquare}\xi_{\square}\nu_{\square} = \xi_{\square\blacksquare}\xi_{\square}.$$

- (5) Reusing parts of the calculation above, we find

$$\nu_{\square\blacksquare}\xi_{\square\blacksquare} = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) = \alpha\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}\right) \nu_{\square\blacksquare} = \nu_{\square\blacksquare}\xi_{\square\blacksquare}\xi_{\square}. \quad \square$$

COROLLARY 6.8. *Two admissible sets of partitions coincide if and only if they have the same intersection with $\{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} : \square, \blacksquare \in \mathcal{F}\}$.*

Furthermore, for an admissible set Π we have the following implications:

- (1) *If Π contains at least one of the partitions $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}$, then it contains $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}$.*
- (2) *If Π contains at least one of the partitions $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}$, then it contains $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}$.*
- (3) *If Π contains two of the basic partitions $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}$, then Π contains all partitions with faces from $\{\square, \blacksquare\}$.*
- (4) *If Π contains two of the basic partitions $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}$, then Π contains all partitions with faces from $\{\square, \blacksquare\}$.*

DEFINITION 6.9. *A 2-faced partition π is called*

- *interval partition if all legs are outer or, equivalently, if all its blocks are intervals; I_{\bullet} denotes the set of all interval partitions,*
- *noncrossing if for all $i < j < k < \ell$ and blocks $\beta, \gamma \in \pi$,*

$$i, k \in \beta, j, \ell \in \gamma \implies \beta = \gamma;$$

NC_{\bullet} denotes the set of all noncrossing partitions,

- binoncrossing if for all $i < j < k < \ell$ and blocks $\beta \neq \gamma \in \pi$,

$$i, k \in \beta, j, \ell \in \gamma \implies \mathbf{f}(j) \neq \mathbf{f}(k),$$

$$i, \ell \in \beta, j, k \in \gamma \implies \mathbf{f}(j) = \mathbf{f}(k);$$

- interval-noncrossing if it is noncrossing and all \circ -legs are outer; $\mathbf{I}_\circ\mathbf{NC}_\bullet$ denotes the set of all interval-noncrossing partitions,
- noncrossing-interval if it is interval-noncrossing after swapping the colors \circ and \bullet ; $\mathbf{NC}_\circ\mathbf{I}_\bullet$ denotes the set of all noncrossing-interval partitions,
- interval-arbitrary if all \circ -legs are outer; $\mathbf{I}_\circ\mathbf{A}_\bullet$ denotes the set of all interval-arbitrary partitions,
- arbitrary-interval if it is interval-arbitrary after swapping the colors \circ and \bullet ; $\mathbf{A}_\circ\mathbf{I}_\bullet$ denotes the set of all arbitrary-interval partitions,
- noncrossing-arbitrary if every block that contains an inner \circ -leg is monochrome and does not cross any other block, i.e. for all legs i, j, k, ℓ and all blocks $\beta \neq \gamma \in \pi$,

$$j, k \in \beta, i, \ell \in \gamma, i < k < \ell, \mathbf{f}(k) = \circ \implies i < j < \ell, \mathbf{f}(j) = \circ;$$

$\mathbf{NC}_\circ\mathbf{A}_\bullet$ denotes the set of all noncrossing-arbitrary partitions,

- arbitrary-noncrossing if it is noncrossing-arbitrary after swapping the colors \circ and \bullet ; $\mathbf{A}_\circ\mathbf{NC}_\bullet$ denotes the set of all arbitrary-interval partitions,
- pure noncrossing if it is noncrossing and all inner blocks are monochrome; \mathbf{pNC}_\circ denotes the set of all pure noncrossing partitions,
- pure crossing if connected inner legs have the same color; \mathbf{pC}_\circ denotes the set of all pure noncrossing partitions,
- arbitrary without any conditions; the set of all bipartitions is also denoted \mathbf{A}_\circ .

THEOREM 6.10. *There are exactly 12 admissible sets of 2-faced partitions (9 if we identify a set with the one obtained by simply swapping the two colors), namely those given in Definition 6.9. Figure 1 displays their respective containment by means of a Hasse diagram and gives minimal generating sets of 2-block partitions.*

Proof. We know that a set obtained from a positive symmetric 2-faced universal product is automatically admissible. Of course, swapping the two colors turns an admissible set into an admissible set. This helps to settle admissibility of a large number of sets in the diagram:

- The sets $\mathbf{I}_\circ, \mathbf{NC}_\circ, \mathbf{A}_\circ$ are the sets of interval, noncrossing, and all partitions (ignoring the colors), and thus are known to come from the trivially two-faced Boolean, free and tensor universal product, respectively. Swapping the colors does not change these sets of partitions.
- The set $\mathbf{NC}_\circ\mathbf{I}_\bullet$ is the set of noncrossing-interval partitions, which originates from free-Boolean independence [17]. Swapping the colors leads to the set $\mathbf{I}_\circ\mathbf{NC}_\bullet$.
- The set $\mathbf{A}_\circ\mathbf{I}_\bullet$ comes from tensor-Boolean independence [10]. Swapping the colors leads to the set $\mathbf{I}_\circ\mathbf{A}_\bullet$.
- The set \mathbf{biNC} is the set of binoncrossing partitions, it comes from bifree independence [3, 32]. Swapping the colors does not change the set.

We are left with the sets of pure crossing and pure noncrossing partitions and with the sets of noncrossing-arbitrary and arbitrary-noncrossing partitions, where again by swapping the colors it is enough to deal with the noncrossing-arbitrary ones. All properties are easily verified.




FIGURE 1. Hasse diagram of all two-colored admissible sets of partitions.

The theorem now follows from the fact that each admissible set is uniquely determined by which basic two-block partitions have nonzero coefficients, and from the implications in Corollary 6.8. \square

COROLLARY 6.11. *Let \odot be a positive symmetric 2-faced universal product. Then the admissible set of partitions*

$$\Pi_{\odot} := \{\pi \in \mathcal{P} : \alpha_{\pi} \neq 0\}$$

is one of the 12 given in Definition 6.9. Furthermore:

- *If $\Pi_{\odot} \in \{NC_{\bullet}A_{\bullet}, A_oNC_{\bullet}, pNC_{\bullet\bullet}, pC_{\bullet\bullet}\}$, then the highest coefficients of \odot are given by the indicator function of Π_{\odot} , and \odot does not coincide with any of the positive symmetric two-faced universal product given in [10, Propositions 5.13 and 6.19].*
- *In all other cases, \odot does coincide with one of the positive symmetric two-faced universal product given in [10, Propositions 5.13 and 6.19]; more concretely,*
 - *if $\Pi_{\odot} = A_{\bullet\bullet}$, then $\odot = \bigotimes \zeta$ with $\zeta = \overline{\xi_{\bullet\bullet}} = \overline{\nu_{\bullet\bullet}}$ is a deformed tensor product,*
 - *if $\Pi_{\odot} = NC_{\bullet\bullet}$, then $\odot = \overrightarrow{\bigstar} \zeta$ with $\zeta = \overline{\nu_{\bullet\bullet}}$ is a deformed free product,*
 - *if $\Pi_{\odot} = biNC_{\bullet\bullet}$, then $\odot = \overrightarrow{\bigstar} \zeta$ with $\zeta = \overline{\nu_{\bullet\bullet}}$ is a deformed bifree product,*
 - *if $\Pi_{\odot} = I_oA_{\bullet}$, then $\odot = \bigotimes$ is the Boolean-tensor product,*
 - *if $\Pi_{\odot} = A_oI_{\bullet}$, then $\odot = \bigotimes$ is the tensor-Boolean product,*

- if $\Pi_{\odot} = \mathbf{I}_\bullet \mathbf{NC}_\bullet$, then $\odot = \diamond_{\ast}$ is the Boolean-free product,
- if $\Pi_{\odot} = \mathbf{NC}_\bullet \mathbf{I}_\bullet$, then $\odot = \overset{\ast}{\diamond}$ is the free-Boolean product,
- if $\Pi_{\odot} = \mathbf{I}_{\bullet\bullet}$, then $\odot = \diamond$ is the Boolean product.

Proof. If \odot is a positive symmetric universal product, then its highest coefficients form an admissible family of weights. If all the basic coefficients are 0 or 1, the family must be given by the indicator function of one of the admissible sets of partitions and all except the mentioned four are identified as positive products in [10]:

- $\mathbf{A}_{\bullet\bullet}$ corresponds to the tensor product
- $\mathbf{NC}_{\bullet\bullet}$ corresponds to the free product
- $\mathbf{biNC}_{\bullet\bullet}$ corresponds to the bifree product
- $\mathbf{I}_\bullet \mathbf{A}_\bullet$ and $\mathbf{A}_\bullet \mathbf{I}_\bullet$ corresponds to the Boolean-tensor and tensor-Boolean product, respectively
- $\mathbf{I}_\bullet \mathbf{NC}_\bullet$ and $\mathbf{NC}_\bullet \mathbf{I}_\bullet$ corresponds to the Boolean-free and free-Boolean product, respectively
- $\mathbf{I}_{\bullet\bullet}$ corresponds to the Boolean product

If one of the basic coefficients is not 0 or 1, Lemma 6.7 leaves only three possibilities, in each of which the universal product has been found to be positive in [10]:

- $\nu_{\bullet\bullet} = \xi_{\bullet\bullet} = q \in \mathbb{T} \setminus \{1\}$, in this case all other basic coefficients are forced to be equal to 1; by comparison of the basic coefficients, the corresponding universal product is the deformed tensor product with $\zeta = \bar{q}$,
- $\nu_{\bullet\bullet} = q \in \mathbb{T} \setminus \{1\}$, $\xi_{\bullet\bullet} = 0$; in this case, the product must coincide with the deformed free product with $\zeta = \bar{q}$,
- $\nu_{\bullet\bullet} = 0$, $\xi_{\bullet\bullet} = q \in \mathbb{T} \setminus \{1\}$; in this case, the product must coincide with the deformed bifree product with $\zeta = \bar{q}$. \square

REMARK 6.12. A remarkable property of freeness is that the free product of traces is again a trace. We cannot expect such a behaviour for any non-trivial multi-faced independence. Indeed, this would force the highest coefficients to be invariant under cyclic permutations, and since we may change the color of the first leg, we could change the color of every leg without changing the coefficient.

REMARK 6.13. Bifreeness allows to define a convolution for probability measures on \mathbb{R}^2 . This comes from the fact that for bifree pairs $(a_1^\circ, a_2^\bullet), (b_1^\circ, b_2^\bullet)$ one always has commutativity of a_1° with b_2^\bullet and of a_2^\bullet with b_1° . Consequently, $a_1^\circ + b_1^\circ$ commutes with $a_2^\bullet + b_2^\bullet$ whenever a_1°, a_2^\bullet commute and b_1°, b_2^\bullet commute. If independent variables in different faces commute, one must have $\xi_{\bullet\bullet} = 1$, which is only the case for tensor and bifree independence.

REMARK 6.14. There are other interesting symmetric two-faced universal products which are not positive, for example the bi-Boolean product. It seems very well possible to do a classification under slightly relaxed conditions, only assuming that one is allowed to change the color of the first leg and not assuming any mirror symmetry (recall that we used changing the color on both sides to show that highest coefficients for all interval partitions are 1). However, it is not clear how to motivate those properties when one does not aim for positivity. For the construction of a universal product in the algebraic sense (see Section 8), Conditions (v) and (vi) of Theorem 5.3 are not necessary at all.

7. MOMENT-CUMULANT RELATIONS

A key tool in the proofs of the subsequent sections are cumulants. In this section, we adapt the theory of cumulants developed in [18] to our special case of symmetric multi-faced independences.

OBSERVATION 7.1. *Let α be a family of weights such that α_π is invertible for every one-block partition. For every family of moments, $m = (m_{\mathbf{f}})_{\mathbf{f} \in \mathcal{F}^*} \in \mathbb{C}^{\mathcal{F}^*}$, there is a unique family of α -cumulants, $c = (c_{\mathbf{f}})_{\mathbf{f} \in \mathcal{F}^*} \in \mathbb{C}^{\mathcal{F}^*}$ such that*

$$(4) \quad m_{\mathbf{f}} = \sum_{\pi \in \mathcal{P}_{<}(\mathbf{f})} \frac{1}{|\pi|!} \alpha_\pi \prod_{\beta \in \pi} c_{|\beta|};$$

indeed, existence and uniqueness of the $c_{\mathbf{f}}$ follows by a standard induction argument. Obviously, the cumulants also determine the moments.

If α is invariant under permutation of blocks, then the formula simplifies to

$$(5) \quad m_{\mathbf{f}} = \sum_{\pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi \prod_{\beta \in \pi} c_{|\beta|}.$$

There is no problem extending formulas (4) and (5) to a multivariate situation. To this end, we think of the (multivariate) moments and cumulants as linear functionals $m, c: A \rightarrow \mathbb{C}$, where $A = \mathbb{C}\langle x_i^{\blacksquare} : \blacksquare \in \mathcal{F}, i \in I^{\blacksquare} \rangle$ is a multi-faced polynomial algebra with (possibly) several indeterminates x_i^{\blacksquare} , $i \in I^{\blacksquare}$, for each face $\blacksquare \in \mathcal{F}$. For a monomial $X = x_{i(1)}^{\mathbf{f}(1)} \cdots x_{i(n)}^{\mathbf{f}(n)}$ and a subset $\beta = \{\ell_1 < \dots < \ell_r\} \subset [n]$, let $X \upharpoonright \beta$ denote the monomial $x_{i(\ell_1)}^{\mathbf{f}(\ell_1)} \cdots x_{i(\ell_r)}^{\mathbf{f}(\ell_r)}$. Cumulants are then defined by the relations

$$(6) \quad m(X) = \sum_{\pi \in \mathcal{P}_{<}(\mathbf{f})} \frac{1}{|\pi|!} \alpha_\pi \prod_{\beta \in \pi} c(X \upharpoonright \beta),$$

$$(7) \quad m(X) = \sum_{\pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi \prod_{\beta \in \pi} c(X \upharpoonright \beta),$$

respectively. In case each I^{\blacksquare} is a one-element set, writing x^{\blacksquare} for the indeterminates, formulas (6) and (7) are recovered by setting $m_{\mathbf{f}} := m(x^{\mathbf{f}(1)} \cdots x^{\mathbf{f}(n)})$ and $c_{\mathbf{f}} := c(x^{\mathbf{f}(1)} \cdots x^{\mathbf{f}(n)})$ for $\mathbf{f} \in \mathcal{F}^n$.

DEFINITION 7.2. *An algebraic probability space is a pair (\mathcal{A}, Φ) , where \mathcal{A} is an algebra and $\Phi: \mathcal{A} \rightarrow \mathbb{C}$ is a linear functional.*

DEFINITION 7.3. *Let (\mathcal{A}, Φ) be an algebraic probability space and $\alpha = (\alpha_\pi)_{\pi \in \mathcal{P}_{(<)}}$ a family of weights on (ordered) multi-faced partitions. For a family $\mathbf{a} = (a_i^{\blacksquare} : \blacksquare \in \mathcal{F}, i \in I^{\blacksquare}) \subset \mathcal{A}$, put $j_{\mathbf{a}}: \mathbb{C}\langle x_i^{\blacksquare} : \blacksquare \in \mathcal{F}, i \in I^{\blacksquare} \rangle \rightarrow \mathcal{A}, x_i^{\blacksquare} \mapsto a_i^{\blacksquare}$. We define its moments by*

$$m_{\mathbf{a}}(X) := \Phi(j_{\mathbf{a}}(X))$$

and its α -cumulants $c_{\mathbf{a}}(X)$ according to the moment-cumulant relations (6) or (7), respectively.

DEFINITION 7.4. *Fix monic weights $(\alpha_\pi)_{\pi \in \mathcal{P}}$. Let $V = \bigoplus_{\blacksquare \in \mathcal{F}} V^{\blacksquare}$ be a vector space with a direct sum decomposition into subspaces according to the faces. Recall that $T_0(V) = \bigoplus_{n \in \mathbb{N}} V^{\otimes n} = \bigsqcup_{\blacksquare \in \mathcal{F}} T_0(V^{\blacksquare})$ denotes the (non-unital) free algebra over V and $T(V) := \bigoplus_{n \in \mathbb{N}_0} V^{\otimes n} = \mathbb{C}1 \oplus T_0(V)$ its unitization, the free unital algebra over V . On the dual space $T_0(V)' = \{\varphi: T_0(V) \rightarrow \mathbb{C} \text{ linear}\}$ we define for $x_i \in V^{\mathbf{f}(i)}$*

$$\exp_{\alpha}(\psi)(x_1 \cdots x_n) := \sum_{\pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi \psi^{\otimes |\pi|}(x_\pi)$$

where for $\pi = \{\beta_1, \dots, \beta_n\}$ we put $x_{\beta_k} := \prod_{i \in \beta_k} x_i$ and $x_\pi := x_{\beta_1} \otimes \dots \otimes x_{\beta_n}$. Then \exp_α is a bijection. We denote the inverse simply as \log_α , which can be calculated recursively,

$$\log_\alpha(\varphi)(x_1 \cdots x_n) = \varphi(x_1 \cdots x_n) - \sum_{\substack{\pi \in \mathcal{P}(\mathbf{f}) \\ |\pi| > 1}} \alpha_\pi \log_\alpha(\varphi)^{\otimes |\pi|}(x_\pi).$$

Note that often \exp_α and \log_α are interpreted as bijections between linear functionals on $T(V)$ vanishing on 1 and unital linear functionals on $T(V)$ by extending the linear functionals from $T_0(V)$ to $T(V)$ accordingly (i.e. ψ and $\log_\alpha(\varphi)$ are extended by annihilating the unit, while $\exp_\alpha(\psi)$ and φ are extended as unital maps).

We use the following conventions.

- If the weights α come from a universal product \odot , we write $\exp_\odot := \exp_\alpha$ and $\log_\odot := \log_\alpha$.
- If $(x_i^\blacksquare)_{i \in I^\blacksquare}$ form a basis of V^\blacksquare , we identify $T(V)$ and $T_0(V)$ with the noncommutative (unital or non-unital) polynomial algebras $\mathbb{C}\langle x_i^\blacksquare : i \in I^\blacksquare, \blacksquare \in \mathcal{F} \rangle$ and $\mathbb{C}\langle x_i^\blacksquare : i \in I^\blacksquare, \blacksquare \in \mathcal{F} \rangle_0$, respectively.

DEFINITION 7.5. Let A be a multi-faced algebra and $\varphi: A \rightarrow \mathbb{C}$ a linear functional. We define $\hat{A} := T_0\left(\bigoplus_{\blacksquare \in \mathcal{F}} A^\blacksquare\right)$ and $\hat{\varphi} := \varphi \circ \mu$, where $\mu: T_0\left(\bigoplus_{\blacksquare \in \mathcal{F}} A^\blacksquare\right) \rightarrow A$ is the canonical homomorphism.

OBSERVATION 7.6. Let α be monic weights. Let furthermore $\varphi: A \rightarrow \mathbb{C}$ be a linear functional on a multi-faced algebra A and $\mathbf{a} = (a_i^\blacksquare \in A^\blacksquare : i \in I^\blacksquare, \blacksquare \in \mathcal{F})$ a family of elements. With the notations from the previous definitions, for $X = x_{i_1}^{\blacksquare_1} \otimes \dots \otimes x_{i_n}^{\blacksquare_n}$, it holds that

$$\hat{\varphi}(a_{i_1}^{\blacksquare_1} \otimes \dots \otimes a_{i_n}^{\blacksquare_n}) = m_{\mathbf{a}}(x_{i_1}^{\blacksquare_1} \otimes \dots \otimes x_{i_n}^{\blacksquare_n}) \text{ and } \log_\alpha \hat{\varphi}(a_{i_1}^{\blacksquare_1} \otimes \dots \otimes a_{i_n}^{\blacksquare_n}) = c_{\mathbf{a}}(x_{i_1}^{\blacksquare_1} \otimes \dots \otimes x_{i_n}^{\blacksquare_n}).$$

OBSERVATION 7.7. Let $h: B \rightarrow A$ be an \mathcal{F} -faced homomorphism between \mathcal{F} -faced algebras B, A and define $\hat{h}: \hat{B} \rightarrow \hat{A}$ as the unique algebra homomorphism with $\hat{h}(b) = h(b)$ for all $b \in B^\blacksquare, \blacksquare \in \mathcal{F}$. Automatically, \hat{h} is an \mathcal{F} -faced homomorphism and fulfills $\mu_A \circ \hat{h} = h \circ \mu_B$. For $\varphi: A \rightarrow \mathbb{C}$ a linear functional, it follows that

$$\widehat{\varphi \circ h} = \varphi \circ h \circ \mu_B = \varphi \circ \mu_A \circ \hat{h} = \hat{\varphi} \circ \hat{h}.$$

Therefore, given monic weights $(\alpha_\pi)_{\pi \in \mathcal{P}}$, one finds that

$$\exp_\alpha(\widehat{\varphi \circ h}) = (\exp_\alpha \hat{\varphi}) \circ \hat{h}, \quad \log_\alpha(\widehat{\varphi \circ h}) = (\log_\alpha \hat{\varphi}) \circ \hat{h}.$$

THEOREM 7.8 (Adjusted and simplified from [18, Th. 7.2]).

A positive and symmetric universal product is uniquely determined by its highest coefficients. More precisely, for $a = a_1 \cdots a_n$ with $a_\ell \in A_{\mathbf{b}(\ell)}^{\mathbf{f}(\ell)}$ so that $a_1 \otimes \dots \otimes a_n \in$

$$T_0\left(\bigoplus_{\kappa \in [2], \blacksquare \in \mathcal{F}} A_\kappa^\blacksquare\right) = \hat{A}_1 \sqcup \hat{A}_2,$$

$$\varphi_1 \odot \varphi_2(a_1 \cdots a_n) = \exp_\odot(\log_\odot(\hat{\varphi}_1) \oplus \log_\odot(\hat{\varphi}_2))(a_1 \otimes \dots \otimes a_n);$$

here we use the direct sum as a shorthand notation for the corresponding linear functional on $T_0\left(\bigoplus_{\kappa \in [2], \blacksquare \in \mathcal{F}} A_\kappa^\blacksquare\right) = \hat{A}_1 \sqcup \hat{A}_2$ as described by Equation (1).

Proof. We only explain why this is a special case of [18, Th. 7.2] and refer the reader to [30, Theorems 2.4.12 and 2.5.13] for a detailed discussion. Since \odot is positive, their are no wrong-ordered highest coefficients. In the symmetric case, the exponential and logarithm map used in [18] coincide with the maps of Definition 7.4 and are therefore determined by the highest coefficients. Since \odot is symmetric, the second ingredient which is in general needed to determine the universal product, namely the n th order cumulant Lie algebra, is trivial for all n . \square

8. RECONSTRUCTION OF UNIVERSAL PRODUCTS FROM HIGHEST COEFFICIENTS

In this section we prove that every admissible family leads to a unique universal product. In particular, we can associate universal products with the admissible sets $\text{NC}_\bullet \mathbf{A}_\bullet$, $\mathbf{A}_\bullet \text{NC}_\bullet$, $\text{pNC}_{\bullet\bullet}$, $\text{pC}_{\bullet\bullet}$. However, it remains an open problem at the moment to decide whether or not those universal products are positive.

LEMMA 8.1. *Suppose that the weights $(\alpha_\pi)_{\pi \in \mathcal{P}}$ are admissible. Fix a family of elements $\mathbf{a} = (a_\ell^\square : \square \in \mathcal{F}, \ell \in I^\square) \subset \mathcal{A}$ in an algebraic probability space (\mathcal{A}, Φ) such that $[n]$ is the disjoint union of the I^\square . Put $\mathbf{f}(\ell) := \square$ if $\ell \in I^\square$ and assume that $\mathbf{f}(i) = \mathbf{f}(i+1) = \square \in \mathcal{F}$ for a certain index $i \in [n]$. We define a modified family $\tilde{\mathbf{a}} = (a_\ell^\square : \square \in \mathcal{F}, \ell \in \tilde{I}^\square)$ where $\tilde{I}^\square := I^\square \setminus \{i, i+1\} \cup \{\{i, i+1\}\}$ and $a_{\{i, i+1\}}^\square := a_i^\square a_{i+1}^\square$. For $X := x_1^{\mathbf{f}(1)} \cdots x_n^{\mathbf{f}(n)}$, $\tilde{X} := x_1^{\mathbf{f}(1)} \cdots x_{i-1}^{\mathbf{f}(i-1)} x_{\{i, i+1\}}^\square x_{i+2}^{\mathbf{f}(i+2)} \cdots x_n^{\mathbf{f}(n)}$ the moments and cumulants according to Definition 7.3 fulfill $m_{\tilde{\mathbf{a}}}(\tilde{X}) = m_{\mathbf{a}}(X)$ and*

$$c_{\tilde{\mathbf{a}}}(\tilde{X}) = c_{\mathbf{a}}(X) + \sum_{\substack{\sigma = \{\beta_1, \beta_2\} \in \mathcal{P}(\mathbf{f}) \\ i \in \beta_1 \neq \beta_2 \ni i+1}} \alpha_\sigma c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2).$$

Proof. The claimed equality for the moments is obvious. The claim for the cumulants is proved by induction on n . For $n = 2$, i.e. $X = x_1^\square x_2^\square$, we have

$$c_{\tilde{\mathbf{a}}}(x_{\{1,2\}}^\square) = m_{\tilde{\mathbf{a}}}(x_{\{1,2\}}^\square) = m_{\mathbf{a}}(x_1^\square x_2^\square) = c_{\mathbf{a}}(x_1^\square x_2^\square) + \alpha \left(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix} \right) c_{\mathbf{a}}(x_1^\square) c_{\mathbf{a}}(x_2^\square).$$

For general n , we can use the moment-cumulant relations for $m_{\mathbf{a}}(X) = m_{\tilde{\mathbf{a}}}(\tilde{X})$ and obtain

$$\begin{aligned} m_{\mathbf{a}}(X) &= \sum_{\pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi \prod_{\beta \in \pi} c_{\mathbf{a}}(X \upharpoonright \beta) \\ &= \sum_{\substack{\pi \in \mathcal{P}(\mathbf{f}) \\ i, i+1 \in \beta \in \pi}} \alpha_\pi c_{\mathbf{a}}(X \upharpoonright \hat{\beta}) \prod_{\beta \in \pi \setminus \{\hat{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta) \\ &\quad + \sum_{\substack{\rho \in \mathcal{P}(\mathbf{f}) \\ \beta_1, \beta_2 \in \rho \\ i \in \beta_1 \neq \beta_2 \ni i+1}} \alpha_\rho c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2) \prod_{\beta \in \rho \setminus \{\beta_1, \beta_2\}} c_{\mathbf{a}}(X \upharpoonright \beta) \\ &= \sum_{\substack{\pi \in \mathcal{P}(\mathbf{f}) \\ i, i+1 \in \beta \in \pi}} \alpha_\pi \left(c_{\mathbf{a}}(X \upharpoonright \hat{\beta}) + \sum_{\substack{\{\beta_1, \beta_2\} \in \mathcal{P}(\hat{\beta}) \\ i \in \beta_1 \neq \beta_2 \ni i+1}} \alpha_{\{\beta_1, \beta_2\}} c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2) \right) \\ &\quad \cdot \prod_{\beta \in \pi \setminus \{\hat{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta) \\ &= c_{\mathbf{a}}(X) + \sum_{\substack{\{\beta_1, \beta_2\} \in \mathcal{P}(\mathbf{f}) \\ i \in \beta_1 \neq \beta_2 \ni i+1}} \alpha_{\{\beta_1, \beta_2\}} c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2) \\ &\quad + \sum_{\substack{\mathbf{f} \neq \pi \in \mathcal{P}(\mathbf{f}) \\ i, i+1 \in \beta \in \pi}} \alpha_\pi \left(c_{\mathbf{a}}(X \upharpoonright \hat{\beta}) + \sum_{\substack{\{\beta_1, \beta_2\} \in \mathcal{P}(\hat{\beta}) \\ i \in \beta_1, i+1 \in \beta_2}} \alpha_{\{\beta_1, \beta_2\}} c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2) \right) \\ &\quad \cdot \prod_{\beta \in \pi \setminus \{\hat{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta) \end{aligned}$$

where we used $\alpha_\rho = \alpha_\pi \alpha_{\{\beta_1, \beta_2\}}$ for $\pi = \rho_{\beta_1 \sim \beta_2}$. On the other hand, with $\tilde{\mathbf{f}} \in \mathcal{F}^{[n]/(i \sim i+1)}$, $\tilde{\mathbf{f}}(\{i, i+1\}) := \mathbf{f}(i) = \mathbf{f}(i+1)$ and $\tilde{\mathbf{f}}(\ell) := \mathbf{f}(\ell)$ for $\ell \neq \{i, i+1\}$,

$$\begin{aligned} m_{\tilde{\mathbf{a}}}(\tilde{X}) &= \sum_{\sigma \in \mathcal{P}(\tilde{\mathbf{f}})} \alpha_\sigma \prod_{\beta \in \sigma} c_{\tilde{\mathbf{a}}}(\tilde{X} \upharpoonright \beta) \\ &= \sum_{\substack{\sigma \in \mathcal{P}(\tilde{\mathbf{f}}) \\ \{i, i+1\} \in \tilde{\beta}}} \alpha_\sigma c_{\tilde{\mathbf{a}}}(\tilde{X} \upharpoonright \tilde{\beta}) \prod_{\beta \in \sigma \setminus \{\tilde{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta) \\ &= c_{\tilde{\mathbf{a}}}(\tilde{X}) + \sum_{\substack{1_{\tilde{\mathbf{f}}} \neq \sigma \in \mathcal{P}(\tilde{\mathbf{f}}) \\ \{i, i+1\} \in \tilde{\beta}}} \alpha_\sigma c_{\tilde{\mathbf{a}}}(\tilde{X} \upharpoonright \tilde{\beta}) \prod_{\beta \in \sigma \setminus \{\tilde{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta). \end{aligned}$$

Recall that there is a canonical bijection between partitions $\sigma \in \mathcal{P}(\tilde{\mathbf{f}})$ and partitions $\pi \in \mathcal{P}(\mathbf{f})$ with $i, i+1$ in the same block $\tilde{\beta} \in \pi$. Also, the highest coefficients α_σ and α_π agree under this bijection by Theorem 5.3 (iii). Using the induction hypothesis on $c_{\mathbf{a}}(X \upharpoonright \tilde{\beta})$ finishes the proof. \square

THEOREM 8.2. *Suppose that the weights $(\alpha_\pi)_{\pi \in \mathcal{P}}$ are admissible. Then there exists a unique symmetric universal product with highest coefficients $(\alpha_\pi)_{\pi \in \mathcal{P}}$.*

Proof. The uniqueness statement is proved in [18, Th. 7.2], see Theorem 7.8.

Let $\varphi_\kappa: A_\kappa \rightarrow \mathbb{C}$ be linear functionals on 2-faced algebras ($\kappa \in \{1, 2\}$). Recall Definition 7.4 of \exp_α and \log_α and Definition 7.5, which sets the notation for lifting φ_k to linear functionals $\hat{\varphi}_\kappa = \varphi_\kappa \circ \mu_\kappa$ on the tensor algebras $T_0(\bigoplus_{\bullet \in \mathcal{F}} A_\kappa^\bullet)$. We simply write $\exp := \exp_\alpha$ and $\log := \log_\alpha$ in the following. We define

$$(8) \quad \varphi_1 \tilde{\odot} \varphi_2 := \exp(\log(\hat{\varphi}_1) \oplus \log(\hat{\varphi}_2)) \in T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1^\bullet \oplus A_2^\bullet) \right)'.$$

The main task is now to prove that $\varphi_1 \tilde{\odot} \varphi_2$ vanishes on the ideal $\mathcal{I} := \ker(\mu_1 \sqcup \mu_2)$ in

$$T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1^\bullet \oplus A_2^\bullet) \right) = \bigsqcup_{\bullet \in \mathcal{F}} \left(T_0(A_1^\bullet) \sqcup T_0(A_2^\bullet) \right)$$

(i.e. the ideal generated by the relations $a \otimes b = ab$ for $a, b \in A_\kappa^\bullet$), so that $\varphi_1 \tilde{\odot} \varphi_2$ descends to a functional $\varphi_1 \odot \varphi_2$ with $(\varphi_1 \odot \varphi_2) \circ (\mu_1 \sqcup \mu_2) = (\varphi_1 \tilde{\odot} \varphi_2)$ on the quotient

$$A_1 \sqcup A_2 = T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1^\bullet \oplus A_2^\bullet) \right) / \mathcal{I}.$$

Let $\mathbf{s} = \mathbf{f} \times \mathbf{b} \in ([2] \times \mathcal{F})^n$ with $\mathbf{s}(i) = \mathbf{s}(i+1) = (j, \blacksquare)$ for some $i \in [n-1], j \in [2], \blacksquare \in \mathcal{F}$. Let $a_1 \cdots a_n \in A^\mathbf{s} \subset A_1 \sqcup A_2$ with $a_\ell \in A_{\mathbf{b}(\ell)}^{\mathbf{f}(\ell)}$, in particular, a_i and a_{i+1} lie in the same direct summand A_j^\bullet of the free product $A_1 \sqcup A_2$. Define $\tilde{\mathbf{f}} \in \mathcal{F}^{[n]/(i \sim i+1)}$, $\tilde{\mathbf{f}}(\{i, i+1\}) = \mathbf{f}(i) = \mathbf{f}(i+1)$ and $\tilde{\mathbf{f}}(\ell) = \mathbf{f}(\ell)$ for $\ell \neq \{i, i+1\}$. Analogously, we define $\tilde{\mathbf{b}}$ and $\tilde{\mathbf{s}}$. With $\mathbf{a} := (a_1, \dots, a_n)$, $X := x_1 \cdots x_n \in \mathbb{C}\langle x_1, \dots, x_n \rangle$, $\tilde{\mathbf{a}} := (a_1, \dots, a_i a_{i+1}, \dots, a_n)$, and $\tilde{X} := x_1 \cdots x_{\{i, i+1\}} x_n \in \mathbb{C}\langle x_1, \dots, x_{\{i, i+1\}}, x_n \rangle$, we have

- for $\beta \subset \{1, \dots, i, i+1, \dots, n\}$ with $a_\ell \in A_\kappa$ for all $\ell \in \beta$

$$\log \hat{\varphi}_\kappa(\mathbf{a} \upharpoonright \beta) = c_{\mathbf{a}}(X \upharpoonright \beta),$$

- for $\beta \subset \{1, \dots, \{i, i+1\}, \dots, n\}$ with $a_\ell \in A_\kappa$ for all $\ell \in \beta$

$$\log \hat{\varphi}_\kappa(\tilde{\mathbf{a}} \upharpoonright \beta) = c_{\tilde{\mathbf{a}}}(\tilde{X} \upharpoonright \beta).$$

Let us say that a partition π is *adapted to \mathbf{b}* , and write $\pi \prec \mathbf{b}$, if \mathbf{b} is constant on blocks of π (this is the first condition of π being adapted to \mathbf{s}). Note that $(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi|}(a_\pi) = 0$ when π is not adapted to \mathbf{b} ; indeed, this follows directly from the

way we identify the direct sum of linear functionals with a linear functional on the tensor algebra in Equation (1). With this in hand, we calculate

$$(9) \quad \exp(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)(a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n) \\ = \sum_{\pi \in \mathcal{P}(\hat{\mathbf{f}})} \alpha_\pi (\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi|} (a_\pi) = \sum_{\substack{\pi \prec \hat{\mathbf{b}} \\ \{i, i+1\} \in \tilde{\beta} \in \pi}} \alpha_\pi c_{\tilde{\mathbf{a}}}(\tilde{X} \upharpoonright \tilde{\beta}) \prod_{\beta \in \pi \setminus \{\tilde{\beta}\}} c_{\tilde{\mathbf{a}}}(\tilde{X} \upharpoonright \beta).$$

On the other hand, if the two legs i and $i+1$ are not identified, then there are partitions adapted to \mathbf{b} for which $i, i+1$ lie in the same block as well as ones for which $i, i+1$ lie in different blocks. This leads to

$$(10) \quad \exp(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)(a_1 \otimes \cdots \otimes a_i \otimes a_{i+1} \otimes \cdots \otimes a_n) \\ = \sum_{\pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi (\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi|} (a_\pi) \\ = \sum_{\substack{\pi \prec \mathbf{b} \\ i, i+1 \in \hat{\beta} \in \pi}} \alpha_\pi c_{\mathbf{a}}(X \upharpoonright \hat{\beta}) \prod_{\beta \in \pi \setminus \{\hat{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta) \\ + \sum_{\substack{\sigma \prec \mathbf{b} \\ \beta_1, \beta_2 \in \sigma \\ i \in \beta_1 \neq \beta_2 \ni i+1}} \alpha_\sigma c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2) \prod_{\beta \in \sigma \setminus \{\beta_1, \beta_2\}} c_{\mathbf{a}}(X \upharpoonright \beta) \\ = \sum_{\substack{\pi \prec \mathbf{b} \\ i, i+1 \in \hat{\beta} \in \pi}} \alpha_\pi \left(c_{\mathbf{a}}(X \upharpoonright \hat{\beta}) + \sum_{\beta_1 \dot{\cup} \beta_2 = \hat{\beta}} \alpha_{\{\beta_1, \beta_2\}} c_{\mathbf{a}}(X \upharpoonright \beta_1) c_{\mathbf{a}}(X \upharpoonright \beta_2) \right) \\ \cdot \prod_{\beta \in \pi \setminus \{\hat{\beta}\}} c_{\mathbf{a}}(X \upharpoonright \beta),$$

using $\alpha_\pi \alpha_{\{\beta_1, \beta_2\}} = \alpha_\sigma$ when $\sigma = \pi \setminus \{\hat{\beta}\} \cup \{\beta_1, \beta_2\}$, i.e. $\pi = \sigma_{\beta_1 \cup \beta_2}$. The two expressions derived in (9) and (10) agree by Lemma 8.1 and, therefore, we have a well-defined map $\varphi_1 \odot \varphi_2(a_1 \cdots a_n) = \varphi_1 \odot \varphi_2(a_1 \otimes \cdots \otimes a_n + \mathcal{I}) := \varphi_1 \tilde{\odot} \varphi_2(a_1 \otimes \cdots \otimes a_n)$.

Let us verify that \odot is indeed a symmetric universal product. To prove universality, recall Observation 7.7. Let $h_\kappa: B_\kappa \rightarrow A_\kappa$ be \mathcal{F} -faced algebra homomorphisms and $\varphi_\kappa: A_\kappa \rightarrow \mathbb{C}$ linear functionals. Then, for $b = b_1 \cdots b_n \in B_1 \sqcup B_2$ and $\hat{b} = b_1 \otimes \cdots \otimes b_n \in \hat{B}_1 \sqcup \hat{B}_2$,

$$\begin{aligned} (\varphi_1 \circ h_1) \odot (\varphi_2 \circ h_2)(b) &= \exp(\log(\widehat{\varphi_1 \circ h_1}) \oplus \log(\widehat{\varphi_2 \circ h_2}))(\hat{b}) \\ &= \exp((\log \hat{\varphi}_1 \circ \hat{h}_1) \oplus (\log \hat{\varphi}_2 \circ \hat{h}_2))(\hat{b}) \\ &= \exp((\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2) \circ (\hat{h}_1 \sqcup \hat{h}_2))(\hat{b}) \\ &= \exp(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2) \circ (\hat{h}_1 \sqcup \hat{h}_2)(\hat{b}) \\ &= ((\varphi_1 \odot \varphi_2) \circ (h_1 \sqcup h_2))(b). \end{aligned}$$

Symmetry and unitality are immediate for $\tilde{\odot}$ and therefore descend to \odot . To prove associativity is slightly more involved. We write $b \in A_1 \sqcup A_2 \sqcup A_3$ as $(\mu_1 \sqcup \mu_2 \sqcup \mu_3)(\hat{b})$ with

$$\hat{b} \in T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1^\bullet \oplus A_2^\bullet \oplus A_3^\bullet) \right) = \bigsqcup_{\bullet \in \mathcal{F}} T_0(A_1^\bullet) \sqcup T_0(A_2^\bullet) \sqcup T_0(A_3^\bullet)$$

and claim that

$$((\varphi_1 \odot \varphi_2) \odot \varphi_3)(b) = (\varphi_1 \odot (\varphi_2 \odot \varphi_3))(b) = \exp(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2 \oplus \log \hat{\varphi}_3)(\hat{b}).$$

The crucial observation is that $\log \widehat{\varphi_1 \odot \varphi_2} = (\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2) \circ \lambda_{12}$ with

$$\begin{aligned} \lambda_{12}: T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1 \sqcup A_2)^\bullet \right) &= \bigsqcup_{\bullet \in \mathcal{F}} T_0(A_1^\bullet \sqcup A_2^\bullet) \\ &\rightarrow T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1^\bullet \oplus A_2^\bullet) \right) = \bigsqcup_{\bullet \in \mathcal{F}} (T_0(A_1^\bullet) \sqcup T_0(A_2^\bullet)) \end{aligned}$$

the unique algebra homomorphism extending the canonical embeddings $A_1^\bullet \sqcup A_2^\bullet \hookrightarrow T_0(A_1^\bullet) \sqcup T_0(A_2^\bullet)$. Indeed, $\widehat{\varphi_1 \odot \varphi_2} = (\varphi_1 \odot \varphi_2) \circ \mu_{12}$ for the canonical map

$$\mu_{12}: T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1 \sqcup A_2)^\bullet \right) \rightarrow A_1 \sqcup A_2,$$

which factorizes as $\mu_{12} = (\mu_1 \sqcup \mu_2) \circ \lambda_{12}$. From $(\varphi_1 \odot \varphi_2) \circ (\mu_1 \sqcup \mu_2) = \varphi_1 \widetilde{\odot} \varphi_2$, we conclude that

$$\widehat{\varphi_1 \odot \varphi_2} = (\varphi_1 \odot \varphi_2) \circ \mu_{12} = (\varphi_1 \odot \varphi_2) \circ (\mu_1 \sqcup \mu_2) \circ \lambda_{12} = (\varphi_1 \widetilde{\odot} \varphi_2) \circ \lambda_{12}.$$

From Definition 7.4 it is obvious that $\exp(\psi \circ \lambda_{12}) = \exp(\psi) \circ \lambda_{12}$ for all $\psi \in T_0 \left(\bigoplus_{\bullet \in \mathcal{F}} (A_1^\bullet \oplus A_2^\bullet) \right)'$, therefore $\log \widehat{\varphi_1 \odot \varphi_2} = (\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2) \circ \lambda_{12}$ as claimed. The rest is easy:

$$\begin{aligned} ((\varphi_1 \odot \varphi_2) \odot \varphi_3)(b) &= \exp(\log(\widehat{\varphi_1 \odot \varphi_2}) \oplus \log \hat{\varphi}_3)(\hat{b}) \\ &= \exp(\log(\hat{\varphi}_1) \oplus \log(\hat{\varphi}_2) \oplus \log(\hat{\varphi}_3))((\lambda_{12} \sqcup \text{id})(\hat{b})) \\ &= \exp(\log(\hat{\varphi}_1) \oplus \log(\hat{\varphi}_2) \oplus \log(\hat{\varphi}_3))(\hat{b}); \end{aligned}$$

note that λ_{12} is actually a projection onto a subalgebra, so we can safely identify \hat{b} with the corresponding element in the domain of $\lambda_{12} \sqcup \text{id}$ instead of introducing yet another symbol for its preimage. The other direction, i.e. $(\varphi_1 \odot (\varphi_2 \odot \varphi_3))(b) = \exp(\log(\hat{\varphi}_1) \oplus \log(\hat{\varphi}_2) \oplus \log(\hat{\varphi}_3))(\hat{b})$, follows by symmetry.

To check that the highest coefficients of \odot are indeed given by α , it is enough to consider products of two functionals φ_1, φ_2 . For $a = a_1 \cdots a_n \in A^{\mathbf{s}}$, $\mathbf{s} = \mathbf{b} \times \mathbf{f} \in ([2] \times \mathcal{F})^*$, and $\sigma \in \mathcal{P}(\mathbf{f})$ the partition with blocks $\beta_\kappa = \{\ell : \mathbf{b}(\ell) = \kappa\}$, we find

$$\begin{aligned} &\left. \frac{\partial^2}{\partial t_1 \partial t_2} (t_1 \varphi_1) \odot (t_2 \varphi_2)(a_1 \cdots a_n) \right|_{\mathbf{t}=0} \\ &= \sum_{\pi \in \mathcal{P}(\mathbf{f})} \frac{\partial^2}{\partial t_1 \partial t_2} \alpha_\pi \cdot (\log t_1 \hat{\varphi}_1 \oplus \log t_2 \hat{\varphi}_2)^{\otimes |\pi|}(a_\pi) \Big|_{\mathbf{t}=0} \\ &= \frac{\partial^2}{\partial t_1 \partial t_2} \alpha_\sigma \cdot \log t_1 \hat{\varphi}_1(a_{\beta_1}) \log t_2 \hat{\varphi}_2(a_{\beta_2}) \Big|_{\mathbf{t}=0} \\ &= \alpha_\sigma \cdot \varphi_1(a_{\beta_1}) \varphi_2(a_{\beta_2}) \end{aligned}$$

as needed.⁽¹¹⁾ □

The formula to compute mixed moments can be considerably simplified in the special case where the the highest coefficients are only 0 or 1.

DEFINITION 8.3. *We say a symmetric universal product is combinatorial with partition set Π if its highest coefficients are all either 0 or 1 and $\Pi = \{\pi \in \mathcal{P} : \alpha_\pi = 1\}$.*

⁽¹¹⁾The notation $a_\beta = \prod_{i \in \beta} a_i$, $a_\pi = \bigotimes_{\beta \in \pi} a_\beta$ refers to $a_1 \otimes \cdots \otimes a_n$, but note that by well-definedness the choice of decomposition of $a_1 \cdots a_n \in A_1 \sqcup A_2$ as a tensor in $T_0(\bigoplus_{\bullet \in \mathcal{F}} A_1^\bullet \oplus A_2^\bullet)$ does not influence the result!

THEOREM 8.4. Let \odot be a combinatorial universal product with admissible partition set Π one of the 12 sets of Theorem 6.10 (in particular, $\mathcal{F} = \{\circ, \bullet\}$ a two element set). Furthermore, let φ_κ be a linear functional on a multi-faced algebra A_κ ($\kappa \in [k]$), and $a = a_1 \cdots a_n \in A^{\mathbf{s}}$ for $\mathbf{s} = \mathbf{b} \times \mathbf{f} \in ([k] \times \mathcal{F})^n$. Denote

- $\pi \in \mathcal{P}(\mathbf{f})$ the multi-faced partition with blocks $\beta_\kappa := \{i : \mathbf{b}(i) = \kappa\}$ (whenever non-empty),
- $\Pi_{\leq \pi} := \{\sigma \in \Pi : \sigma \leq \pi\}$ the set of refinements of π inside $\Pi \cap \mathcal{P}(\mathbf{f})$,
- S the set of maximal elements of $\Pi_{\leq \pi}$ (i.e. coarsest refinements of π inside Π),
- $\wedge R$ is the maximal common refinement of partitions in $R \subset \Pi \cap \mathcal{P}(\mathbf{f})$, $\wedge \emptyset := 1_{\mathbf{f}}$,
- $\Phi := \varphi_1 \odot \cdots \odot \varphi_k$,
- $\hat{\Phi}$ the lift of Φ to $T_0\left(\bigoplus_{\ell \in [k], \bullet \in \mathcal{F}} A_\ell^\bullet\right)$.

Then

$$\Phi(a) = \sum_{\emptyset \neq R \subseteq S} (-1)^{\#R-1} \hat{\Phi}^{\otimes |\wedge R|}(a_{\wedge R}).$$

Proof. Put $\Psi := \log \hat{\Phi} = \log \hat{\varphi}_1 \oplus \cdots \oplus \log \hat{\varphi}_k$. The key observation is that a refinement σ of a partition $\rho \in \Pi$ belongs to Π if and only if $\sigma \upharpoonright \beta \in \Pi$ for all blocks $\beta \in \rho$; this can be easily seen for each of the 12 admissible sets of partitions individually. Using the moment cumulant formula on each block of $\wedge R$ and the observation on refinements just made, we find

$$(11) \quad \sum_{R \subseteq S} (-1)^{\#R} \hat{\Phi}^{\otimes |\wedge R|}(a_{\wedge R}) = \sum_{R \subseteq S} \sum_{\substack{\sigma \leq \wedge R \\ \sigma \in \Pi}} (-1)^{\#R} \Psi^{\otimes |\sigma|}(a_\sigma)$$

(equality of the summands for $R = \emptyset$ will be discussed below.) Now, the same partition $\sigma \in \Pi$ can of course be a refinement of $\wedge R$ for different $R \subseteq S$. Denote $T(\sigma) := \{\rho \in S : \sigma \leq \rho\}$ and $n(\sigma) := \#T(\sigma)$. Then $\sigma \leq \wedge R$ if and only if $R \subseteq T(\sigma)$, and for every $k \in \{0, 1, \dots, n(\sigma)\}$ there are $\binom{n(\sigma)}{k}$ many such R with $\#R = k$. If $n(\sigma) = 0$, i.e. if σ is not a refinement of π , then $\Psi^{\otimes |\sigma|}(a_\sigma) = 0$ because mixed cumulants vanish. This leads to

$$\text{RHS of (11)} = \sum_{\sigma \in \Pi_{\leq \pi}} \underbrace{\sum_{k=0}^{n(\sigma)} (-1)^k \binom{n(\sigma)}{k}}_{=0} \Psi^{\otimes |\sigma|}(a_\sigma) = 0.$$

Recall that we defined $\wedge \emptyset := 1_{\mathbf{f}}$, so that

$$\Phi(a) = \hat{\Phi}(a_{1_{\mathbf{f}}}) = \sum_{\sigma \in \mathcal{P}(\mathbf{f}) \cap \Pi} \Psi^{\otimes |\sigma|}(a_\sigma) = \sum_{\substack{\sigma \leq 1_{\mathbf{f}} \\ \sigma \in \Pi}} \Psi^{\otimes |\sigma|}(a_\sigma);$$

this confirms that the choice is consistent with Equation (11), and it also shows that the statement of the theorem is equivalent to LHS of (11) = 0. \square

EXAMPLE 8.5. Let \odot be the universal product associated with $\text{NC}_\bullet \text{A}_\bullet$. Then $\overline{\downarrow \downarrow \downarrow \downarrow}$ has set of coarsest refinements $S = \{\overline{\downarrow \downarrow \downarrow \downarrow}, \overline{\downarrow \downarrow \downarrow \downarrow}, \downarrow \downarrow \downarrow \downarrow\}$ in $\text{NC}_\bullet \text{A}_\bullet$ with $\wedge S = \downarrow \downarrow \downarrow \downarrow$, leading to

$$\begin{aligned} & \varphi \odot \psi(a_1^\circ b_1^\bullet a_2^\bullet a_3^\circ b_2^\bullet) \\ &= \varphi(a_1^\circ a_2^\bullet) \varphi(a_3^\circ) \psi(b_1^\bullet b_2^\bullet) + \varphi(a_1^\circ a_2^\bullet a_3^\circ) \psi(b_1^\bullet) \psi(b_2^\bullet) - \varphi(a_1^\circ a_2^\bullet) \varphi(a_3^\circ) \psi(b_1^\bullet) \psi(b_2^\bullet) \end{aligned}$$

for all $\varphi \in A'$, $\psi \in B'$, $a_i^\bullet \in A^*$ ($i = 1, 2, 3$), and $b_j^\bullet \in B^*$ ($j = 1, 2$).

9. UNIT PRESERVING UNIVERSAL PRODUCTS

In [4], Diaz-Aguilera, Gaxiola, Santos, and Vargas characterize when the moment cumulant relation associated with weights on partitions leads to independent constants, finding this to be the case if and only if the weights do not change when removing or inserting a singleton from or to the partition. Manzel and Schürmann discuss in [18, Rem. 3.1] the relation between universal products in the category of multi-faced algebras and in the category of multi-faced unital algebras and observe that while a product for the unital category always gives rise to a product for the non-unital category, the other way round requires a condition, namely that the universal product *respects the units* or is *unit preserving* as we prefer to write in this article. In this section we briefly review universal products in the category of multi-faced unital algebras, define what exactly it means to be unit preserving, generalize the definition of *singleton inductive weights* to the multi-faced setting, and finally characterize unit preserving symmetric universal product as those whose highest coefficients are singleton inductive.

In the category of unital algebras with unital algebra homomorphisms, the coproduct is given by the *unital free product*, which can be constructed from the non-unital free product as

$$\mathcal{A}_1 \sqcup \mathcal{A}_2 := (\mathcal{A}_1 \sqcup \mathcal{A}_2) / \langle 1_{\mathcal{A}_1} - 1_{\mathcal{A}_2} \rangle;$$

here $\langle \cdot \rangle$ denotes the generated two-sided ideal.

DEFINITION 9.1.

- A multi-faced unital algebra is a unital algebra A with unital subalgebras \mathcal{A}^\blacksquare , $\blacksquare \in \mathcal{F}$, such that the canonical unital algebra homomorphism $\bigsqcup_{\blacksquare \in \mathcal{F}} \mathcal{A}^\blacksquare \rightarrow A$ is an isomorphism, in which case we write $A = \bigsqcup_{\blacksquare \in \mathcal{F}} \mathcal{A}^\blacksquare$.
- A multi-faced unital algebra homomorphism is a unital algebra homomorphism which maps face into face.
- The unital free product of multi-faced unital algebras $\mathcal{A}_1, \mathcal{A}_2$ is a multi-faced unital algebra with $(\mathcal{A}_1 \sqcup \mathcal{A}_2)^\blacksquare := \mathcal{A}_1^\blacksquare \sqcup \mathcal{A}_2^\blacksquare$.
- A linear functional $\phi: A \rightarrow \mathbb{C}$ on a multi-faced unital algebra is unital if $\phi(1_A) = 1$.

Multi-faced unital algebras with multi-faced unital algebra homomorphisms form a category, in which \sqcup is a coproduct. One can adapt Definition 3.1 to the unital situation and obtains the following.

DEFINITION 9.2. A universal product in the category of multi-faced unital algebras is a binary product operation for unital linear functionals on multi-faced unital algebras which associates with unital functionals ϕ_1, ϕ_2 on multi-faced unital algebras $\mathcal{A}_1, \mathcal{A}_2$, respectively, a unital functional $\phi_1 \odot \phi_2$ on $\mathcal{A}_1 \sqcup \mathcal{A}_2$ such that

- $(\phi_1 \circ h_1) \odot (\phi_2 \circ h_2) = (\phi_1 \odot \phi_2) \circ (h_1 \sqcup h_2)$ for all multi-faced unital algebra homomorphisms $h_i: \mathcal{B}_i \rightarrow \mathcal{A}_i$ (universality)
- $(\phi_1 \odot \phi_2) \odot \phi_3 = \phi_1 \odot (\phi_2 \odot \phi_3)$ (associativity)
- $(\phi_1 \odot \phi_2) \upharpoonright \mathcal{A}_i = \phi_i$ (restriction property).

As Manzel and Schürmann noticed in [18, Rem. 3.1], every universal product $\widetilde{\odot}$ in the category of multi-faced unital algebras gives rise to a universal product in the sense of Definition 3.1, simply putting

$$\varphi_1 \odot \varphi_2 := \tilde{\varphi}_1 \widetilde{\odot} \tilde{\varphi}_2 \upharpoonright A_1 \sqcup A_2 \subset \tilde{A}_1 \sqcup \tilde{A}_2$$

where \tilde{A}_i denotes the unitization of a multi-faced algebra and $\tilde{\varphi}_i$ the unital extension of a linear functional.

Conversely, if a universal product \odot in the non-unital case is given, one would like to define

$$(12) \quad \phi_1 \widetilde{\odot} \phi_2(p(a)) := \varphi_1 \odot \varphi_2(a)$$

with the following conventions:

- $A_\kappa := \bigsqcup_{\blacksquare \in \mathcal{F}} \mathcal{A}_\kappa^\blacksquare$, so that $\mathcal{A}_\kappa \cong A_\kappa / \mathcal{I}_{\mathcal{A}_\kappa}$ with $\mathcal{I}_{\mathcal{A}_\kappa} := \langle 1^\blacksquare - 1^\blacksquare : \blacksquare, \blacksquare \in \mathcal{F} \rangle \subset A_\kappa$,
- $p_{\mathcal{A}_\kappa} : A_\kappa \rightarrow \mathcal{A}_\kappa$ denotes the canonical homomorphism,
- $\varphi_\kappa := \phi_\kappa \circ p_{\mathcal{A}_\kappa} : A_\kappa \rightarrow \mathbb{C}$, i.e. $\varphi_\kappa(a) := \phi_\kappa(a + \mathcal{I}_{\mathcal{A}_\kappa})$,
- $p : A_1 \sqcup A_2 \rightarrow \mathcal{A}_1 \sqcup \mathcal{A}_2$ denotes the canonical homomorphism.

DEFINITION 9.3. A universal product is unit preserving (or respects units) if, whenever A_1, A_2 are multi-faced algebras with each A_i^\blacksquare unital and φ_i a linear functional on A_i which vanishes on the ideal $\langle 1_i^\blacksquare - 1_i^\blacksquare : \blacksquare, \blacksquare \in \mathcal{F} \rangle \subset A_i$ and such that $\varphi_i \upharpoonright A_i^\blacksquare$ is unital for every $\blacksquare \in \mathcal{F}$, then $\varphi_1 \odot \varphi_2$ vanishes on the ideal $\langle 1_i^\blacksquare - 1_j^\blacksquare : i, j \in [2], \blacksquare, \blacksquare \in \mathcal{F} \rangle \subset A_1 \sqcup A_2$ and $\varphi_1 \odot \varphi_2 \upharpoonright A_i^\blacksquare$ is unital for every $i \in [2], \blacksquare \in \mathcal{F}$.

REMARK 9.4. A multi-faced universal product is unit preserving if and only if (12) is well-defined, in which case it yields a universal product in the category of multi-faced unital algebras [18, Rem. 3.1]. Since Manzel and Schürmann do not give a definition of “respecting units”, let us briefly check that Definition 9.3 captures what they mean.

Assume that \odot is unit preserving. The $\varphi_i = \phi_i \circ p_{\mathcal{A}_i}$ in (12) are linear functionals on A_i , vanish on $\ker p_{\mathcal{A}_i} = \mathcal{I}_{\mathcal{A}_i} = \langle 1_i^\blacksquare - 1_i^\blacksquare : \blacksquare, \blacksquare \in \mathcal{F} \rangle \subset A_i$ and fulfill $\varphi_i(1_i^\blacksquare) = \phi_i(1_i) = 1$. Therefore, we may conclude that $\varphi_1 \odot \varphi_2$ vanishes on the ideal $\langle 1_i^\blacksquare - 1_j^\blacksquare : i, j \in [2], \blacksquare, \blacksquare \in \mathcal{F} \rangle \subset A_1 \sqcup A_2$, which coincides with the kernel of the canonical homomorphism $p : A_1 \sqcup A_2 \rightarrow \mathcal{A}_1 \sqcup \mathcal{A}_2$. This means that there is a well-defined linear functional $\phi_1 \widetilde{\odot} \phi_2$ with $\varphi_1 \odot \varphi_2 = (\phi_1 \widetilde{\odot} \phi_2) \circ p$. This functional is also unital because $\phi_1 \widetilde{\odot} \phi_2(1) = \phi_1 \widetilde{\odot} \phi_2(p(1_i^\blacksquare)) = \varphi_i(1_i^\blacksquare) = 1$.

We leave the rest of the simple, but notationally cumbersome proof of the claim (in particular universality and associativity of $\widetilde{\odot}$) to the interested reader.

In the following we will need often remove a singleton block $\beta = \{s\}$ from a partition $\pi \ni \beta$. While consistent use of notation would dictate to write $\pi \setminus \{\beta\} = \pi \setminus \{\{s\}\}$, we will prefer to write $\pi \setminus \{s\}$ for better legibility.

DEFINITION 9.5 (multi-faced version of [4, Def. 3.2]). A family of weights $(\alpha_\pi)_{\pi \in \mathcal{P}}$ is singleton inductive if $\alpha_\pi = \alpha_{\pi \setminus \{s\}}$ for every singleton block $\{s\} \in \pi$.

LEMMA 9.6 (multi-faced version of [4, Th. 3.2]). Let α be monic, singleton inductive weights. Suppose that $A = \bigsqcup_{\blacksquare \in \mathcal{F}} A^\blacksquare$ is a multi-faced algebra such that each face A^\blacksquare is unital (with unit 1^\blacksquare) and that φ fulfills $\varphi(1^\blacksquare) = 1$ and φ vanishes on the ideal $\langle 1^\blacksquare - 1^\blacksquare : \blacksquare, \blacksquare \in \mathcal{F} \rangle \subset A$. Then

$$\log_\alpha \hat{\varphi}(a_1 \otimes \cdots \otimes a_n) = 0 \text{ whenever } n > 1 \text{ and } a_s = 1^\blacksquare \text{ for some } s \in [n], \blacksquare \in \mathcal{F};$$

here $\hat{\varphi}$ is the lift of φ to a $T_0(\bigoplus_{\blacksquare \in \mathcal{F}} A^\blacksquare)$.

Proof. We prove the claim by induction. For $n = 2$ and arbitrary $\blacksquare, \blacksquare \in \mathcal{F}$,

$$\log_\alpha \hat{\varphi}(1^\blacksquare \otimes a^\blacksquare) = \varphi(1^\blacksquare a^\blacksquare) - \varphi(1^\blacksquare) \varphi(a^\blacksquare) = \varphi(1^\blacksquare a^\blacksquare) - \varphi(a^\blacksquare) = 0$$

and analogously $\log_\alpha \hat{\varphi}(a^\blacksquare \otimes 1^\blacksquare) = 0$. Now assume the statement holds for all $1 < m < n$ and consider $a = a_1 \otimes \cdots \otimes a_n$ with $a_i \in A^{\mathbf{f}(i)}$, $a_s = 1^{\mathbf{f}(s)}$. Note that $\varphi(a_1 \cdots a_n) = \varphi(a_1 \cdots \check{a}_s \cdots a_n)$ (here \check{a}_s means omission of the factor) and $\log_\alpha \hat{\varphi}(a_s) = \log_\alpha \hat{\varphi}(1^\blacksquare) =$

$\varphi(1^\bullet) = 1$. We find

$$\begin{aligned}
 \log_\alpha \hat{\varphi}(a_1 \otimes \cdots \otimes a_n) &= \varphi(a_1 \cdots a_n) - \sum_{\pi \in \mathcal{P}(\mathbf{f}) \setminus \{1_\mathbf{f}\}} \alpha_\pi(\log_\alpha \hat{\varphi})^{\otimes |\pi|}(a_\pi) \\
 &= \varphi(a_1 \cdots a_n) - \sum_{\{s\} \in \pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi(\log_\alpha \hat{\varphi})^{\otimes |\pi|}(a_\pi) \\
 &\quad - \underbrace{\sum_{\{s\} \notin \pi \in \mathcal{P}(\mathbf{f}) \setminus \{1_\mathbf{f}\}} \alpha_\pi(\log_\alpha \hat{\varphi})^{\otimes |\pi|}(a_\pi)}_{=0 \text{ by induction hypothesis}} \\
 &= \varphi(a_1 \cdots \check{a}_s \cdots a_n) - \sum_{\{s\} \in \pi \in \mathcal{P}(\mathbf{f})} \alpha_{\pi \setminus \{s\}}(\log_\alpha \hat{\varphi})^{\otimes |\pi|-1}(a_{\pi \setminus \{s\}}) \log_\alpha \hat{\varphi}(a_s) \\
 &= \varphi(a_1 \cdots \check{a}_s \cdots a_n) - \sum_{\sigma \in \mathcal{P}(\mathbf{f}[n] \setminus \{s\})} \alpha_\sigma(\log_\alpha \hat{\varphi})^{\otimes |\sigma|}(a_\sigma) = 0,
 \end{aligned}$$

where we used that the weights are singleton inductive as well as the moment cumulant relation for $a_1 \cdots \check{a}_s \cdots a_n$. \square

THEOREM 9.7. *For a multi-faced positive symmetric universal product \odot , the following are equivalent.*

- (1) \odot is unit preserving,
- (2) $\nu_\bullet = 1$ for all $\bullet \in \mathcal{F}$,
- (3) the highest coefficients of \odot are singleton inductive.

Proof. Let \odot be unit preserving. To calculate $\nu_\bullet = \alpha(\begin{smallmatrix} \square \\ \bullet \end{smallmatrix})$, we can assume that also the extremal legs are \bullet -legs. We can therefore ignore the faces and calculate, as in the single-faced case,

$$\varphi_1 \odot \varphi_2(aba') = \nu_\bullet \cdot \varphi_1(aa')\varphi_2(b) + \gamma \cdot \varphi_1(a)\varphi_1(a')\varphi_2(b)$$

for all $a, a' \in A_1^\bullet, b \in A_2^\bullet$, with some universal constant $\gamma \in \mathbb{C}$. Suppose that φ_1, φ_2 are as in Definition 9.3 and furthermore $b = 1_2^\bullet, \varphi_1(a) = \varphi_1(a') = 0, \varphi_1(aa') = 1$, then

$$\varphi_1 \odot \varphi_2(a1_2^\bullet a') = \varphi_1 \odot \varphi_2(a1_1^\bullet a') = \varphi_1 \odot \varphi_2(aa') = \varphi_1(aa') = 1$$

because \odot preserves units. We also have $\varphi_2(b) = \varphi_2(1_2^\bullet) = 1$. Putting everything together, $\nu_\bullet = 1$.

A simple induction on the number of blocks shows that $\alpha_\pi = \nu_\bullet \cdot \alpha_{\pi \setminus \{s\}}$ whenever $\pi \in \mathcal{P}$ has a singleton block $\{s\} \in \pi$ of color \bullet . Therefore, $\nu_\bullet = 1$ for all $\bullet \in \mathcal{F}$ implies that the highest coefficients are singleton inductive.

Now assume that the highest coefficients of a positive symmetric universal product are singleton inductive. Let A_1, A_2 be multi-faced algebras with unital faces, $\mathbf{s} = \mathbf{b} \times \mathbf{f} \in ([2] \times \mathcal{F})^n$, $a_\ell \in A_{\mathbf{b}(\ell)}^{\mathbf{f}(\ell)}$ for $\ell \in [n]$, $\hat{a} = a_1 \otimes \cdots \otimes a_n$, $a = a_1 \cdots a_n$ and $a_s = 1_i^\bullet$. Then, with $\log := \log_\odot$,

$$\begin{aligned}
 \varphi_1 \odot \varphi_2(a) &= \sum_{\pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi|}(\hat{a}_\pi) \\
 &= \sum_{\{s\} \in \pi \in \mathcal{P}(\mathbf{f})} \alpha_\pi(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi|}(\hat{a}_\pi)
 \end{aligned}$$

by Lemma 9.6. Because α is singleton inductive, $\alpha_\pi = \alpha_{\pi \setminus \{s\}}$. Also, for any π with $\{s\} \in \pi$, we have

$$(\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi|}(\hat{a}_\pi) = (\log \hat{\varphi}_1 \oplus \log \hat{\varphi}_2)^{\otimes |\pi \setminus \{s\}|}(\hat{a}_{\pi \setminus \{s\}}) \underbrace{\log \hat{\varphi}_{\mathbf{b}(s)}(a_s)}_{=1}.$$

Therefore, $\varphi_1 \odot \varphi_2(a_1 \cdots a_n) = \varphi_1 \odot \varphi_2(a_1 \cdots \check{a}_s \cdots a_n)$. This calculation works for any $i \in [2]$ and any $\blacksquare \in \mathcal{F}$, so the statement follows. \square

COROLLARY 9.8. *A 2-faced positive symmetric universal product is unit preserving if and only if its associated set of partitions contains $\text{pNC}_{\bullet\bullet}$.*

10. SUMMARY AND OUTLOOK

We found conditions on weights that are necessarily satisfied by the highest coefficients of a positive two-faced universal product. In the symmetric case, we showed that weights which fulfill these conditions are always the highest coefficients of a uniquely determined universal product. We could also determine all families of weights which fulfill these conditions, thereby providing a list of candidates for positive symmetric universal products.

We hope that the methods developed in this work will eventually lead to a complete classification of positive multi-faced universal products. To that end, the following problems will have to be overcome:

- Prove or disprove positivity of the “exceptional cases” which do not admit a representation on free or tensor product.
- Extend the classification of admissible weights to more than two faces.
- Extend the classification of admissible weights to the non-symmetric case.
- Extend the reconstruction theorem to the non-symmetric case. This might be significantly more difficult because the cumulants have to be combined using the Campbell-Baker-Hausdorff formula instead of just the direct sum.

APPENDIX A. COMPARISON WITH [30]

Most ideas behind the proofs in Sections 5, 6 and 8 go more or less back to [30]. A crucial difference between this article and the exposition in [30] is that our main results are consistently formulated and proved for families of weights on partitions, while Varšo often works with sets of partitions instead, which means that in [30] several results are only proved in the *combinatorial* case in the sense of Definition 8.3. The weight-based approach often helped us to streamline proofs. Another difference is that we decided to focus on *positive* universal products here.

In the following we give a more detailed comparison of the results.

- Theorem 5.3 (iv) is basically [30, Corollary 5.2.6]. The remaining claims of Theorem 5.3 generalize [30, Theorem 5.2.17] to possibly non-symmetric universal products. Because we put more emphasis on positive products, for ease of reading, we only formulated Theorem 5.3 for positive products while Varšo formulates his results more generally for products with the “right ordered monomials property”, i.e. those products for which the conclusion of Theorem 3.3 holds; however, we mention in the proof where exactly the positivity condition is used and where the right-ordered monomials property is enough.
- Lemma 6.3 is closely related to [30, Theorem 5.2.20] (since admissible families of weights are not defined in [30], the statement is formulated for families of highest coefficients of certain universal products). Lemma 6.7 has overlap with [30, Lemma 5.2.23]; however, from Lemma 6.7 (3) it follows that all coefficients have absolute value in $\{0, 1\}$, which goes beyond what was found in [30]. Regarding the main classification results, Theorem 6.10 corresponds to [30, Theorem 4.2.44] and Corollary 6.11 strengthens [30, Remark 5.2.28].
- Observation 5.9 draws the connection between the admissible sets of partitions as defined from admissible weights in Definitions 5.5 and 5.6 and Varšo’s

(m -colored) universal classes of partitions [30, Definition 3.4.9]. The only difference is that admissible sets are assumed to contain the interval partitions, while a universal class of partitions is also allowed to consist of the 1-block partitions alone.

- Our reconstruction theorem, Theorem 8.2, also covers universal products with non-0-1 highest coefficients, in contrast to [30, Theorem 3.4.32]. The crucial Lemma 8.1 corresponds to [30, Lemma 3.4.24] (formulated and proved for admissible weights instead of universal classes of partitions).

Theorem 8.4 and the results of Section 9 have no counterpart in [30].

Acknowledgements. We are grateful to Michael Schürmann for numerous fruitful discussions in the course of this research. MG thanks Moritz Weber and Roland Speicher for stimulating discussions and atmosphere during his stay in Saarbrücken for the focus semester on quantum information Autumn 2022. We thank both anonymous referees for their comments, which helped us improve the quality of this article. We truly appreciate one of the referees extraordinary detailed reading and in particular his thoughtful suggestions regarding notation.

REFERENCES

- [1] Anis Ben Ghorbal and Michael Schürmann, *Non-commutative notions of stochastic independence*, Math. Proc. Cambridge Philos. Soc. **133** (2002), no. 3, 531–561, <https://doi.org/10.1017/S0305004102006072>.
- [2] ———, *Quantum Lévy processes on dual groups*, Math. Z. **251** (2005), no. 1, 147–165, <https://doi.org/10.1007/s00209-005-0793-x>.
- [3] Ian Charlesworth, Brent Nelson, and Paul Skoufranis, *On two-faced families of non-commutative random variables*, Canad. J. Math. **67** (2015), no. 6, 1290–1325, <https://doi.org/10.4153/CJM-2015-002-6>.
- [4] Carlos Diaz-Aguilera, Tulio Gaxiola, Jorge Santos, and Carlos Vargas, *Combinatorics of NC-probability spaces with independent constants*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **25** (2022), no. 2, article no. 2250009 (20 pages), <https://doi.org/10.1142/S0219025722500096>.
- [5] Kurusch Ebrahimi-Fard and Frédéric Patras, *Cumulants, free cumulants and half-shuffles*, Proc. Royal Soc. A **471** (2015), no. 2176, article no. 20140843 (18 pages), <https://doi.org/10.1098/rspa.2014.0843>.
- [6] Kurusch Ebrahimi-Fard, Frédéric Patras, and Roland Speicher, *ϵ -noncrossing partitions and cumulants in free probability*, Int. Math. Res. Not. IMRN **2018** (2018), no. 23, 7156–7170, <https://doi.org/10.1093/imrn/rnx098>.
- [7] Uwe Franz, *Lévy processes on quantum groups and dual groups*, in Quantum independent increment processes. II, Lecture Notes in Math., vol. 1866, Springer, Berlin, 2006, pp. 161–257, https://doi.org/10.1007/11376637_3.
- [8] Malte Gerhold, *Schoenberg correspondence for multifaced independence*, 2021, <https://arxiv.org/abs/2104.02985>.
- [9] ———, *Bi-monotone Brownian motion*, in Infinite Dimension Analysis, Quantum Probability and Related Topics (Noboru Watanabe, Luigi Accardi, and Si Si, eds.), QP-PQ: Quantum Probab. White Noise Anal., vol. 32, World Sci. Publ., Hackensack, NJ, 2023, Proceedings of the International Conference on Infinite Dimensional Analysis, Quantum Probability and Related Topics, QP38 (2017), https://doi.org/10.1142/9789811275999_0005.
- [10] Malte Gerhold, Takahiro Hasebe, and Michaël Ulrich, *Towards a classification of multi-faced independence: A representation-theoretic approach*, J. Funct. Anal. **285** (2023), no. 3, article no. 109907, <https://doi.org/10.1016/j.jfa.2023.109907>.
- [11] Malte Gerhold and Stephanie Lachs, *Classification and GNS-construction for general universal products*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **18** (2015), no. 1, article no. 1550004 (29 pages), <https://doi.org/10.1142/S0219025715500046>.
- [12] Malte Gerhold, Stephanie Lachs, and Michael Schürmann, *Categorical independence and Lévy processes*, SIGMA Symmetry Integrability Geom. Methods Appl. **18** (2022), article no. 075 (27 pages), <https://doi.org/10.3842/SIGMA.2022.075>.

- [13] Yinzhen Gu, Takahiro Hasebe, and Paul Skoufranis, *Bi-monotonic independence for pairs of algebras*, J. Theoret. Probab. **33** (2020), no. 1, 533–566, <https://doi.org/10.1007/s10959-019-00884-2>.
- [14] Yinzhen Gu and Paul Skoufranis, *Bi-Boolean independence for pairs of algebras*, Complex Anal. Oper. Theory **13** (2019), no. 7, 3023–3089, <https://doi.org/10.1007/s11785-017-0750-9>.
- [15] Takahiro Hasebe and Hayato Saigo, *The monotone cumulants*, Ann. Inst. Henri Poincaré Probab. Stat. **47** (2011), no. 4, 1160–1170, <https://doi.org/10.1214/10-AIHP379>.
- [16] Weihua Liu, *Free-free-boolean independence for triples of algebras*, 2018, <https://arxiv.org/abs/1801.03401>.
- [17] ———, *Free-Boolean independence for pairs of algebras*, J. Funct. Anal. **277** (2019), no. 4, 994–1028, <https://doi.org/10.1016/j.jfa.2019.05.005>.
- [18] Sarah Manzel and Michael Schürmann, *Non-commutative stochastic independence and cumulants*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **20** (2017), no. 2, article no. 1750010 (38 pages), <https://doi.org/10.1142/S0219025717500102>.
- [19] Wojciech Młotkowski, *Λ -free probability*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **7** (2004), no. 1, 27–41, <https://doi.org/10.1142/S0219025704001517>.
- [20] Naofumi Muraki, *Monotonic independence, monotonic central limit theorem and monotonic law of small numbers*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **4** (2001), no. 1, 39–58, <https://doi.org/10.1142/S0219025701000339>.
- [21] ———, *The five independences as quasi-universal products*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **5** (2002), no. 1, 113–134, <https://doi.org/10.1142/S0219025702000742>.
- [22] ———, *The five independences as natural products*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **6** (2003), no. 3, 337–371, <https://doi.org/10.1142/S0219025703001365>.
- [23] ———, *A simple proof of the classification theorem for positive natural products*, Probab. Math. Statist. **33** (2013), no. 2, 315–326, <https://www.math.uni.wroc.pl/~pms/files/33.2/Article/33.2.12.pdf>.
- [24] Alexandru Nica and Roland Speicher, *Lectures on the combinatorics of free probability*, London Mathematical Society Lecture Note Series, vol. 335, Cambridge University Press, Cambridge, 2006, <https://doi.org/10.1017/CB09780511735127>.
- [25] Michael Schürmann and Stefan Voß, *Schoenberg correspondence on dual groups*, Comm. Math. Phys. **328** (2014), no. 2, 849–865, <https://doi.org/10.1007/s00220-013-1872-1>.
- [26] Roland Speicher, *Multiplicative functions on the lattice of noncrossing partitions and free convolution*, Math. Ann. **298** (1994), no. 4, 611–628, <https://doi.org/10.1007/BF01459754>.
- [27] ———, *On universal products*, in Free probability theory (Waterloo, ON, 1995), Fields Inst. Commun., vol. 12, Amer. Math. Soc., Providence, RI, 1997, pp. 257–266, <https://doi.org/10.1090/fic/012/12>.
- [28] Roland Speicher and Reza Woroudi, *Boolean convolution*, in Free probability theory (Waterloo, ON, 1995), Fields Inst. Commun., vol. 12, Amer. Math. Soc., Providence, RI, 1997, pp. 267–279, <https://doi.org/10.1090/fic/012/13>.
- [29] Roland Speicher and Janusz Wysoczański, *Mixtures of classical and free independence*, Arch. Math. (Basel) **107** (2016), no. 4, 445–453, <https://doi.org/10.1007/s00013-016-0955-6>.
- [30] Philipp Varšo, *Studies on positive and symmetric two-faced universal products*, Ph.D. thesis, University of Greifswald, 2021, <https://nbn-resolving.org/urn:nbn:de:gbv:9-opus-75524>, pp. 271 + xii.
- [31] Dan Voiculescu, *Symmetries of some reduced free product C^* -algebras*, in Operator algebras and their connections with topology and ergodic theory (Buşteni, 1983), Lecture Notes in Math., vol. 1132, Springer, Berlin, 1985, pp. 556–588, <https://doi.org/10.1007/BFb0074909>.
- [32] Dan-Virgil Voiculescu, *Free probability for pairs of faces I*, Comm. Math. Phys. **332** (2014), no. 3, 955–980, <https://doi.org/10.1007/s00220-014-2060-7>.
- [33] Wilhelm von Waldenfels, *An approach to the theory of pressure broadening of spectral lines*, in Probability and information theory, II (M. Behara, K. Krickeberg, and J. Wolfowitz, eds.), Springer, Berlin, 1973, pp. 19–69. Lecture Notes in Math., Vol. 296, <https://doi.org/10.1007/BFb0059819>.
- [34] ———, *Interval partitions and pair interactions*, in Séminaire de Probabilités, IX (Seconde Partie, Univ. Strasbourg, Strasbourg, années universitaires 1973/1974 et 1974/1975), Springer, Berlin, 1975, pp. 565–588. Lecture Notes in Math., Vol. 465, https://www.numdam.org/item/SPS_1975__9__565_0/.

MALTE GERHOLD, Saarland University, Faculty of Mathematics, Postbox 151150, D-66041 Saarbruecken, Germany

E-mail : gerhold@math.uni-sb.de

Url : <https://sites.google.com/view/malte-gerhold/>

PHILIPP VARŠO

Url : <https://orcid.org/0000-0001-9199-2516>