



ALGEBRAIC COMBINATORICS


Joseph Johnson & Ricky Ini Liu

Birational rowmotion and the octahedron recurrence

Volume 7, issue 5 (2024), p. 1453-1477.

<https://doi.org/10.5802/alco.385>

© The author(s), 2024.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Algebraic Combinatorics is published by *The Combinatorics Consortium*
and is a member of the *Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Birational rowmotion and the octahedron recurrence

Joseph Johnson & Ricky Ini Liu

ABSTRACT We use the octahedron recurrence to give a simplified statement and proof of a formula for iterated birational rowmotion on a product of two chains, first described by Musiker and Roby. Using this, we show that weights of certain chains in rectangles shift in a predictable way under the action of rowmotion. We then define generalized Stanley–Thomas words whose cyclic rotation uniquely determines birational rowmotion on the product of two chains. We also discuss the relationship between rowmotion and birational RSK and give a birational analogue of Greene’s theorem in this setting.

1. INTRODUCTION

For any poset P , (*combinatorial*) *rowmotion* is the action on the set of order ideals of P that sends I to the ideal generated by the minimal elements of $P \setminus I$. This action is well studied in the dynamical algebraic combinatorics literature; for background on rowmotion, see [27]. On certain classes of posets (triangles, skeletal posets, rectangles, root posets, and others), rowmotion has a surprisingly small period, and it also sometimes exhibits other interesting phenomena such as homomesy and cyclic sieving: see [5, 12, 15, 19, 23, 28].

Rowmotion also has a description in terms of local, involutive transformations called *toggles* [2]. Reinterpreting these toggles as acting on lattice points in \mathbb{R}^n , one can lift toggles and hence rowmotion to piecewise-linear maps. One can then lift these further to the birational realm by replacing max with addition, addition with multiplication, and subtraction with division [4, 5]. Surprisingly, many results from the combinatorial level remain true on the birational level. For instance, for some posets, the period of birational rowmotion remains small, even though a priori it need not even be finite [9, 10].

The main poset of interest in this paper is the product of two chains, called the *rectangle poset*. Grinberg and Roby [9] show that birational rowmotion on the $r \times s$ rectangle has the same order as combinatorial rowmotion, $r + s$. Musiker and Roby [19] then give an explicit combinatorial formula for all powers of birational rowmotion on rectangles in terms of nonintersecting lattice paths. However, their impressive formula is notationally dense, and their proof requires a rather intricate bijection on lattice paths. The main result of this paper is a simplified statement and proof of Musiker and Roby’s iterated birational rowmotion formula for rectangles. (This essentially

Manuscript received 7th September 2022, revised 25th March 2024, accepted 20th June 2024.

KEYWORDS. rowmotion, RSK correspondence, octahedron recurrence, Stanley–Thomas words.

ACKNOWLEDGEMENTS. R. I. Liu was partially supported by National Science Foundation grants DMS-1700302 and CCF-1900460.

gives a new proof of periodicity of birational rowmotion on rectangles.) Our proof relies mainly on the connections between rowmotion, the octahedron recurrence, the solid minors of a matrix, and nonintersecting lattice paths via the Lindström–Gessel–Viennot Lemma.

We also touch upon a number of topics related to this work. For instance, associated to any antichain of the $r \times s$ rectangle poset is a certain 0/1-sequence of length $r + s$ called the *Stanley–Thomas word*. Rowmotion on order ideals equivariantly induces a rowmotion action on antichains, which cyclically shifts the Stanley–Thomas word [23, 26]. Previously, the Stanley–Thomas word has been defined in the birational realm to prove homomesy results [15], but this word alone is not enough to uniquely specify a general labeling of a rectangle. In this article, we define *generalized Stanley–Thomas words* in terms of certain sums of weights of chains and show that birational rowmotion is the unique function that cyclically shifts all of them.

Finally, we discuss the relationship between the birational version of the *Robinson–Schensted–Knuth (RSK) correspondence* [3, 20] and rowmotion by defining birational RSK in terms of toggles. We use the iterated birational rowmotion formula to show that this definition satisfies a birational version of Greene’s theorem and also compare it to existing constructions in the literature.

ROAD MAP OF THE PAPER. In Section 2 we review background on birational rowmotion, the octahedron recurrence, and the relationship between the two. In Section 3 we state and prove our main result, the iterated birational rowmotion formula on rectangles, using the Lindström–Gessel–Viennot Lemma and the octahedron recurrence. Using this framework we prove a chain shifting lemma in Section 4 and define generalized Stanley–Thomas words. Finally in Section 5 we define birational RSK and prove a birational analogue of Greene’s theorem, which we then use to show that the cyclic shifting of the generalized Stanley–Thomas words uniquely determines birational rowmotion.

2. BACKGROUND

2.1. POSETS AND RECTANGLES. We first review some basic terminology about posets. Typically we represent a finite poset $P = (P, \leq)$ by its *Hasse diagram*, a directed graph with vertex set P and edges $x \rightarrow y$ when y covers x , which we denote $x \lessdot y$. Since all edges are directed upward in the Hasse diagram, we omit the direction in figures.

DEFINITION 2.1. *Let P be a poset. A chain in P is a sequence of elements $p_1 < p_2 < \dots < p_k$. An antichain in P is a set $A \subseteq P$ such that for any distinct $p, q \in A$, neither $p \leq q$ nor $q \leq p$.*

The antichains are related to the order ideals (and order filters) of a poset.

DEFINITION 2.2. *Let P be a finite poset.*

- (1) *An order ideal of P is a set $I \subseteq P$ such that if $p \leq q$ in P and $q \in I$, then $p \in I$.*
- (2) *An order filter of P is a set $F \subseteq P$ such that if $p \leq q$ in P and $p \in F$, then $q \in F$.*
- (3) *An interval of P is a subset of the form $[p, q] = \{x \mid p \leq x \leq q\}$.*

Let $[r]$ be the chain with r elements $1 < 2 < \dots < r$. Of particular interest is the *rectangle poset* given by the Cartesian product of two chains $R = [r] \times [s]$. We will distinguish between a rectangle poset and general posets by using R exclusively for the rectangle.

DEFINITION 2.3. *Let $R = [r] \times [s]$ be a rectangle poset.*




FIGURE 1. The rectangle $[2] \times [3]$.

- For fixed i , the i th row of R is the set of all elements in R of the form (i, j) .
- For fixed j , the j th column of R is the set of all elements in R of the form (i, j) .
- The k th rank of R is the set of all elements (i, j) such that $i + j = k$.
- The k th file of R is the set of all elements (i, j) such that $j - i = k$.

We will typically draw rectangles oriented as in Figure 1, so that rows run southwest to northeast, columns run southeast to northwest, ranks are aligned horizontally, and files are aligned vertically. (As a word of caution, note that the minimum element has rank 2.)

2.2. ROWMOTION. An important object of study in dynamical algebraic combinatorics is a certain dynamical process on order ideals called combinatorial rowmotion.

DEFINITION 2.4. Let I be an order ideal of P . Combinatorial rowmotion is the map ρ that sends I to the order ideal generated by the minimal elements of $P \setminus I$.

Cameron and Fon-der-Flaass [2] give another description of rowmotion in terms of combinatorial toggles. Let $J(P)$ denote the set of order ideals of P . For each $p \in P$, we associate a toggle map $t_p: J(P) \rightarrow J(P)$ by

$$t_p(I) = \begin{cases} I \cup \{p\} & \text{if } p \notin I \text{ and } I \cup \{p\} \in J(P), \\ I \setminus \{p\} & \text{if } p \in I \text{ and } I \setminus \{p\} \in J(P), \\ I & \text{otherwise.} \end{cases}$$

Combinatorial rowmotion can then be defined as the composition of toggles on P in the order of a linear extension $\mathcal{L}: P \rightarrow [n]$ (where $n = |P|$) from top to bottom, that is,

$$\rho = t_{\mathcal{L}^{-1}(1)} \circ \cdots \circ t_{\mathcal{L}^{-1}(n)}.$$

We note that $t_p \circ t_q = t_q \circ t_p$ if and only if p and q do not form a cover relation. Consequently this description is independent of the choice of linear extension.

In [26], Stanley gives a bijection (also discovered independently by Thomas) between the order ideals of $R = [r] \times [s]$ and 0/1-sequences with r zeroes and s ones.

DEFINITION 2.5. Let A be an antichain of $R = [r] \times [s]$. The Stanley–Thomas word of A is $w(A) = (w_1, \dots, w_{r+s})$, where

$$w_i = \begin{cases} 1 & \text{if } 1 \leq i \leq r \text{ and } A \text{ has an element in row } i, \\ 1 & \text{if } r + 1 \leq i \leq r + s \text{ and } A \text{ has no element in column } i - r, \\ 0 & \text{otherwise.} \end{cases}$$

Rowmotion performs a cyclic shift of the Stanley–Thomas word [23]. Since the map w is a bijection, this gives a simple proof of the following result, originally proved by Brouwer and Schrijver [1].

THEOREM 2.6 ([1]). *The order of ρ on $R = [r] \times [s]$ is $r + s$.*

Note that $r + s$ is one more than the number of distinct ranks of R . This is the smallest possible order that rowmotion can have on a graded poset: iteratively applying rowmotion to the empty order ideal simply adds one rank of elements at a time to the order ideal until arriving at the entire poset, which is then mapped back to the empty order ideal.

2.3. PIECEWISE-LINEAR ROWMOTION. In [25], Stanley defines two polytopes related to a finite poset.

DEFINITION 2.7. *Let P be a finite poset.*

- (1) *The order polytope $\mathcal{O}(P) \subseteq \mathbb{R}^P$ is the set of all $x \in \mathbb{R}^P$ satisfying the inequalities $0 \leq x_p \leq 1$ for all $p \in P$, and $x_p \leq x_q$ for all $p, q \in P$ satisfying $p \leq q$.*
- (2) *The chain polytope $\mathcal{C}(P) \subseteq \mathbb{R}^P$ is the set of all $x \in \mathbb{R}^P$ satisfying the inequalities $x_p \geq 0$ for all $p \in P$, and $\sum_{p \in C} x_p \leq 1$ for all (maximal) chains $C \subseteq P$.*

The vertices of $\mathcal{O}(P)$ and $\mathcal{C}(P)$ are the indicator vectors of the order filters and antichains of P , respectively. We may also identify the vertices of $\mathcal{O}(P)$ with the order ideals of P by complementation.

In [25], Stanley defines a piecewise-linear, continuous, and volume-preserving bijection $\phi: \mathcal{O}(P) \rightarrow \mathcal{C}(P)$ called the *transfer map*, defined by

$$\phi(x)_p = x_p - \max_{q < p} x_q,$$

where we interpret an empty max as 0. The inverse of this map is given by

$$\phi^{-1}(x)_p = \max_{c_1 < c_2 < \dots < c_k = p} \left(\sum_{i=1}^k x_{c_i} \right).$$

The transfer map can be thought of as a piecewise-linear extension of the map that sends an order filter to its minimal elements.

Combinatorial rowmotion permutes order ideals and so can also be thought of as a permutation of the vertices of $\mathcal{O}(P)$. As is the case with the transfer map, there is a natural piecewise-linearization of this bijection described by Einstein and Propp [5].

DEFINITION 2.8. *The piecewise-linear toggle on the order polytope $\mathcal{O}(P)$ corresponding to an element $p \in P$ is the map $t_p: \mathcal{O}(P) \rightarrow \mathcal{O}(P)$ that changes the p th coordinate by*

$$x_p \mapsto \min_{q > p} x_q + \max_{q < p} x_q - x_p$$

and fixes all other coordinates, where an empty min is interpreted as 1 and an empty max is interpreted as 0.

(We will abuse notation and use the same symbol for combinatorial toggles and piecewise-linear toggles.) Note that t_p only depends on the coordinates in the neighborhood of $p \in P$ in the Hasse diagram. Consequently for $p, q \in P$, $t_p \circ t_q = t_q \circ t_p$ if and only if neither $p < q$ nor $p > q$ as in the combinatorial realm.

We then define *piecewise-linear rowmotion* $\rho: \mathcal{O}(P) \rightarrow \mathcal{O}(P)$ using piecewise-linear toggles by

$$\rho = t_{\mathcal{L}^{-1}(1)} \circ \dots \circ t_{\mathcal{L}^{-1}(|P|)},$$

where \mathcal{L} is any linear extension of P .

2.4. BIRATIONAL ROWMOTION. Piecewise-linear rowmotion can be lifted even further to a birational analogue that uses addition, multiplication, and division in place of max, addition, and subtraction, respectively. This lifting process is called *detropicalization*. (See [4, 5, 20] for more detailed discussion.) The functions resulting from detropicalization are generally *subtraction-free* and therefore well-defined on positive labelings of P .

As an example, the *birational transfer map* ϕ acts on positive labelings $x \in \mathbb{R}_{>0}^P$ via coordinate functions

$$\phi(x)_p = \frac{x_p}{\sum_{q < p} x_q}$$

where an empty sum is interpreted as 1.

Since $\min(a, b) = -\max(-a, -b)$, detropicalizing \min yields the *parallel sum* $\#$ defined by

$$a \# b = \frac{1}{\frac{1}{a} + \frac{1}{b}} = \frac{ab}{a + b}.$$

Parallel sum is associative and commutative. For a finite multiset $S \subseteq \mathbb{R}_{>0}$, we denote the parallel sum of all elements in S by $\sum_{s \in S}^{\#} s$.

DEFINITION 2.9. *The birational toggle on $\mathbb{R}_{>0}^P$ corresponding to an element $p \in P$ is the birational map $t_p: \mathbb{R}_{>0}^P \rightarrow \mathbb{R}_{>0}^P$ that changes the p th coordinate by*

$$x_p \mapsto \left(\sum_{q > p}^{\#} x_q \right) \left(\sum_{q < p} x_q \right) \cdot \frac{1}{x_p}$$

and fixes all other coordinates, where an empty sum or parallel sum is interpreted as 1.

(We again abuse notation by using the same notation for birational toggles as piecewise-linear toggles.) Similarly to piecewise-linear rowmotion, we define *birational rowmotion* as

$$\rho = t_{\mathcal{L}^{-1}(1)} \circ \cdots \circ t_{\mathcal{L}^{-1}(|P|)}$$

for any linear extension \mathcal{L} of P . From the birational setting, we can obtain the piecewise-linear analogue via a *valuation* or *tropicalization*: see, for instance, [4, 5].

For any poset, one can compute birational rowmotion in terms of the dual transfer map.

DEFINITION 2.10. *Let P be a poset and $x \in \mathbb{R}_{>0}^P$ be a labeling. The dual transfer map is the birational function $\phi^*: \mathbb{R}_{>0}^P \rightarrow \mathbb{R}_{>0}^P$ with coordinate functions*

$$\phi^*(x)_p = \frac{x_p}{\sum_{q > p} x_q}$$

for all $p \in P$, where an empty sum is interpreted as 1.

In other words, ϕ^* acts on labelings of P in the same way that ϕ acts on the associated labeling of the dual of P .

The following lemma is due to Einstein and Propp (see for instance [4, 14]). We include a short proof for completeness.

LEMMA 2.11. *Let P be a finite poset. Then for any $x \in \mathbb{R}_{>0}^P$ and $p \in P$,*

$$\rho \circ \phi^{-1}(x)_p = \frac{1}{(\phi^*)^{-1}(x)_p}.$$

Proof. Let $y = \phi^{-1}(x)$ and $z = (\phi^*)^{-1}(x)$. For any $p \in P$, suppose that $\rho(y)_q = \frac{1}{z_q}$ for all $q > p$. When applying the toggle at p in the computation of $\rho(y)$, the elements at or below p are labeled as in y while those above p are labeled as in $\rho(y)$. Hence by the definition of t_p ,

$$\rho(y)_p = \left(\sum_{q>p} \# \frac{1}{z_q} \right) \left(\sum_{q<p} y_q \right) \cdot \frac{1}{y_p} = \frac{1}{\sum_{q>p} z_q} \cdot \frac{1}{x_p} = \frac{1}{z_p}.$$

The result follows easily by induction starting at the top of P . □

Computing iterated applications of rowmotion is more difficult. However, one can sometimes prove results about rowmotion using an associated combinatorial construction instead. As mentioned previously, one such example is the Stanley–Thomas word, which lifts to the birational level.

DEFINITION 2.12. *Let $R = [r] \times [s]$. The birational Stanley–Thomas word w for x is the word of length $r + s$ defined by*

$$w_i = \begin{cases} \prod_{j=1}^s x_{ij} & \text{if } 1 \leq i \leq r, \\ \prod_{j=1}^r x_{j,i-r}^{-1} & \text{if } r + 1 \leq i \leq r + s. \end{cases}$$

In [15] it is shown that $\phi \circ \rho \circ \phi^{-1}$ cyclically shifts the birational Stanley–Thomas word. Though this is not sufficient to show that the order of birational rowmotion on the product of two chains has finite order, it can be used to extend other results (such as instances of homomesy) to the birational level. In Section 4 we will define a collection of *generalized Stanley–Thomas words*, and in Section 5 we show that the cyclic rotation of these words uniquely determines rowmotion.

2.5. DODGSON CONDENSATION AND THE OCTAHEDRON RECURRENCE. Given a matrix $A = (a_{ij})_{i,j=1}^n$, let $A_{ij}^{(k)}$ denote the $k \times k$ submatrix of A formed by the intersection of rows i through $i+k-1$ and columns j through $j+k-1$ whenever $1 \leq i, j \leq n-k+1$. These minors satisfy the following algebraic relation known as the *Desnanot–Jacobi identity*, which forms the basis of a recursive algorithm for computing the determinant of a matrix called *Dodgson condensation*.

PROPOSITION 2.13. *For $k \geq 0$,*

$$\det(A_{ij}^{(k+1)}) \det(A_{i+1,j+1}^{(k-1)}) = \det(A_{ij}^{(k)}) \det(A_{i+1,j+1}^{(k)}) - \det(A_{i,j+1}^{(k)}) \det(A_{i+1,j}^{(k)}).$$

(By convention, we set $\det(A_{ij}^{(0)}) = 1$ and $\det(A_{ij}^{(-1)}) = 0$ for all integers i and j .)

We visualize this relation by placing the values $\det(A_{ij}^{(k)})$ into a three-dimensional array. In all figures containing such an array, we place the entries $\det(A_{ij}^{(k)})$ at height k so that, for $k > 1$, $\det(A_{ij}^{(k)})$ lies directly above the center of the submatrix at height 1 for which it is the determinant. See Figure 2.

Proposition 2.13 then implies that the entries $M_{ij}^{(k)} = \det(A_{ij}^{(k)})$ satisfy the following *octahedron recurrence*:

$$M_{ij}^{(k)} M_{i+1,j+1}^{(k)} = M_{i+1,j+1}^{(k-1)} M_{ij}^{(k+1)} + M_{i+1,j}^{(k)} M_{i,j+1}^{(k)}.$$

The elements in this relation lie at the vertices of an octahedron that is a translation of the one shown in Figure 2. By convention, we extend the array $M = (M_{ij}^{(k)})$ to all integers i, j , and k by setting any undefined values equal to 0, which does not violate the octahedron recurrence.




FIGURE 2. The coordinate system we will use for figures depicting three-dimensional arrays $(M_{ij}^{(k)})$. (Note that the k direction does not point vertically but rather along the edge of the pyramid.) The octahedron recurrence involves the vertices of translations of the octahedron as shown on the right.




FIGURE 3. The relationship between the octahedron recurrence and birational toggles as described in Lemma 2.14. Two labelings of the poset by quotients of entries in the octahedron recurrence are related by a birational toggle at the central element.

We now demonstrate the known relationship between toggles and the octahedron recurrence appearing in [9]. A visualization of this lemma is shown in Figure 3. If part of the rectangle poset is labeled via the quotients $z_{ij}^{(k)}$ as shown, then Lemma 2.14 shows that a toggle at (i, j) transforms $z_{ij}^{(k-1)}$ to $z_{ij}^{(k)}$.

LEMMA 2.14. *Suppose $M = (M_{ij}^{(k)})$ is an array of indeterminates satisfying the octahedron recurrence, and let $z_{ij}^{(k)} = \frac{M_{k+2, i+k+1}^{(j-1)}}{M_{k+1, i+k+1}^{(j)}}$. Then as rational functions in the entries of M ,*

$$z_{ij}^{(k)} = \frac{(z_{i, j-1}^{(k)} + z_{i-1, j}^{(k)})(z_{i+1, j}^{(k-1)} \# z_{i, j+1}^{(k-1)})}{z_{ij}^{(k-1)}}.$$

Proof. For ease of notation, we relabel the relevant part of the array as in Figure 3. (For instance, $x = M_{k+1,i+k}^{(j-1)}$ and $y = M_{k,i+k}^{(j)}$, so $z_{ij}^{(k-1)} = \frac{x}{y}$.) Applying the octahedron recurrence to the two octahedra shown and dividing gives

$$\frac{xx'}{yy'} = \frac{ad + bc}{cf + de} = \frac{\frac{a}{c} + \frac{b}{d}}{\frac{f}{d} + \frac{e}{c}} = \left(\frac{a}{c} + \frac{b}{d}\right) \left(\frac{d}{f} \# \frac{c}{e}\right).$$

Solving for $\frac{x'}{y'}$ yields the desired identity

$$\frac{x'}{y'} = \frac{\left(\frac{a}{c} + \frac{b}{d}\right) \left(\frac{c}{e} \# \frac{d}{f}\right)}{\frac{x}{y}}. \quad \square$$

REMARK 2.15. In most cases when we apply Lemma 2.14, all quantities will specialize to well-defined values. There are two exceptions: if we set $e = M_{k,i+k+1}^{(j)} = 0$ (which would make $\frac{c}{e} = z_{i+1,j}^{(k-1)}$ undefined), then in the proof we instead have

$$\frac{xx'}{yy'} = \frac{ad + bc}{cf} = \frac{\frac{a}{c} + \frac{b}{d}}{\frac{f}{d}} = \left(\frac{a}{c} + \frac{b}{d}\right) \left(\frac{d}{f}\right),$$

which yields the same result as Lemma 2.14 but with the undefined term $z_{i+1,j}^{(k-1)}$ removed from the parallel sum. A similar result holds if we instead set $f = M_{k,i+k}^{(j+1)} = 0$ (which would make $\frac{d}{f} = z_{i,j+1}^{(k-1)}$ undefined).

Lemma 2.14 suggests that toggling/rowmotion should be thought of as translation with respect to the octahedron recurrence. In the next section, we will see how this relationship can be used to prove the iterated birational rowmotion formula for rectangles.

3. BIRATIONAL ROWMOTION IN THE RECTANGLE POSET

In this section, we will use the relationship between the octahedron recurrence and toggles to prove a formula for any power of birational rowmotion on a rectangle similar to one given by Musiker–Roby [19]. This formula will be described in terms of nonintersecting paths inside a certain graph \mathcal{G}_R . We will begin by proving a lemma that relates nonintersecting paths in \mathcal{G}_R to nonintersecting paths in R .

3.1. NONINTERSECTING PATHS. Let $R = [r] \times [s]$, and let the element (i, j) have weight x_{ij} . For $1 \leq k \leq s$, define $\mathcal{P}_R^{(k)}$ to be the set of all collections \mathcal{L} of k nonintersecting (i.e. vertex disjoint) paths traveling up along the edges of the Hasse diagram of R , starting at $\{(1, 1), (1, 2), \dots, (1, k)\}$ and ending at $\{(r, s - k + 1), \dots, (r, s)\}$. The weight $w(\mathcal{L})$ is the product of the weights of the elements in the paths of \mathcal{L} . We will write $w_R^{(k)} = w_R^{(k)}(x)$ for the sum of the weights of all collections of paths in $\mathcal{P}_R^{(k)}$. We also write $w_R = w_R^{(s)}$ for the product of all weights in R . Similarly, if I is any interval in R , then we define $\mathcal{P}_I^{(k)}$, $w_I^{(k)}(x)$, and w_I in an analogous manner.

Note that while the definition of $\mathcal{P}_R^{(k)}$ is asymmetric in r and s , if $k \leq \min\{r, s\}$, then the paths in any $\mathcal{L} \in \mathcal{P}_R^{(k)}$ must pass through all k elements at rank $k + 1$ as well as those at rank $r + s - k + 1$. Thus in this case $w_R^{(k)}$ will remain unchanged if we transpose R . We will generally assume that $k \leq \min\{r, s\}$ when considering $\mathcal{P}_R^{(k)}$.

Define \mathcal{G}_R to be the graph with vertex set $[r + 1] \times [s]$, directed edges from (i, j) to $(i + 1, j)$ of weight x_{ij}^{-1} , and directed edges from (i, j) to $(i + 1, j - 1)$ of weight 1. (The vertices of R are in bijection with the edges of \mathcal{G}_R whose weights are not 1.) As an example, Figure 4 shows $R = [2] \times [3]$ and the corresponding graph \mathcal{G}_R .




FIGURE 4. The poset R and graph \mathcal{G}_R for $r = 2$ and $s = 3$. Edges are directed up and to the left in \mathcal{G}_R , and unlabeled edges have weight 1. The highlighted paths correspond via the bijection in Lemma 3.3.




FIGURE 5. Bijection between $\mathcal{P}_R^{(k)}$ and $\mathcal{S}_{k+1, r+1}^{(s-k)}$ as in Lemma 3.1. Tiles on the left are replaced with the corresponding tiles on the right.

We label the vertices on the boundary of \mathcal{G}_R as follows: for $1 \leq i \leq r$ and $1 \leq j \leq s$, define

$$P_j = (1, j), \quad P_{s+i} = (i + 1, s), \quad Q_i = (i, 1), \quad Q_{r+j} = (r + 1, j).$$

We then define $\mathcal{S}_{ij}^{(k)}$ to be the set of all collections \mathcal{L}' of k nonintersecting paths starting from $\{P_i, \dots, P_{i+k-1}\}$ and ending at $\{Q_j, \dots, Q_{j+k-1}\}$. The weight $w(\mathcal{L}')$ of \mathcal{L}' is the product of the weights of the edges in the paths of \mathcal{L}' . Let $W_{ij}^{(k)} = W_{ij}^{(k)}(x)$ be the total weight of all $\mathcal{L}' \in \mathcal{S}_{ij}^{(k)}$. (By convention, $W_{ij}^{(0)} = 1$ and $W_{ij}^{(k)} = 0$ when $k < 0$.)

The following result relates paths in R to paths in \mathcal{G}_R .

LEMMA 3.1. *There exists a bijection from $\mathcal{P}_R^{(k)}$ to $\mathcal{S}_{k+1, r+1}^{(s-k)}$ that divides weight by w_R .*

Proof. In the drawing of $\mathcal{L} \in \mathcal{P}_R^{(k)}$, extend the start and end of each path by a half-edge from northwest to southeast. Such a drawing can be built by piecing together square tiles centered at each vertex showing the five ways in which the paths can pass through that vertex in such a way that the tiles match along sides—see Figure 5.

We can replace each tile with a new tile containing edges/half-edges parallel to the edges of \mathcal{G}_R in a way that preserves the exits along the southwest and northeast sides but inverts the exits along the southeast and northwest sides as shown in Figure 5. Placing vertices on the southeast and northwest edges of these new tiles, the resulting drawing will then depict a collection of paths \mathcal{L}' in \mathcal{G}_R that forms an element of $\mathcal{S}_{k+1,r+1}^{(s-k)}$. (This bijection also has the following alternate description: when the two drawings are overlaid, each of the k horizontal steps in the i th path in \mathcal{L}' crosses the i th northeast step in one of the k paths in \mathcal{L} .)

The weighted edges in \mathcal{L}' correspond to elements of R not lying in any path of \mathcal{L} . Thus $w(\mathcal{L}')$ is the inverse of the weight of the complement of \mathcal{L} in R , which is exactly $\frac{w(\mathcal{L})}{w_R}$. \square

Let $A = (a_{ij})_{i,j=1}^{r+s}$ be the square matrix such that $a_{ij} = W_{ij}^{(1)}$, the total weight of all paths from P_i to Q_j in \mathcal{G}_R . By the Lindström–Gessel–Viennot Lemma [7, 18], each minor $\det(A_{ij}^{(k)})$ is equal to $W_{ij}^{(k)}$, the total weight of $\mathcal{S}_{ij}^{(k)}$, the set of all collections of nonintersecting paths from $\{P_i, P_{i+1}, \dots, P_{i+k-1}\}$ to $\{Q_j, Q_{j+1}, \dots, Q_{j+k-1}\}$. By the discussion in Section 2.5, the three-dimensional array $W = (W_{ij}^{(k)})$ formed by these minors satisfies the octahedron recurrence. See Figure 7 for a depiction of W when $R = [2] \times [3]$.

We will need a few simple properties of the array W .

PROPOSITION 3.2. *Let $R = [r] \times [s]$, and let $W = (W_{ij}^{(k)})$ be the corresponding three-dimensional array. Assume $k > 0$ and $0 \leq i, j \leq r + s + 1 - k$.*

- (a) *If $W_{ij}^{(k)} \neq 0$, then $i \leq j \leq i + r$.*
- (b) *If $i = j$, then $W_{ij}^{(k)} = 1$.*
- (c) *If $i < j$ and $k > s$, then $W_{ij}^{(k)} = 0$.*

Proof. All of these follow from the description of $W_{ij}^{(k)}$ as the total weight of $\mathcal{S}_{ij}^{(k)}$. For (a), there are no paths from P_i to Q_j if $i > j$ or $j - i > r$. For (b), if $i = j$, then $\mathcal{S}_{ij}^{(k)}$ has only one element consisting only of horizontal steps of weight 1.

For (c), if $i < j$ and $k > s$, then the first $s + 1$ starting points P_i, \dots, P_{i+s} all lie at or below row $i + 1$, while the ending points for these paths lie at or above row $j \geq i + 1$. But row $i + 1$ only has s vertices, so it is impossible for there to be $s + 1$ nonintersecting paths passing through it. \square

We can translate Lemma 3.1 into a statement expressing weights of certain collections of nonintersecting paths in R in terms of the array of minors $W = (W_{ij}^{(k)})$. In fact, we can formulate a similar result for paths inside certain intervals $I \subseteq R$, specifically those for which the leftmost point of I lies on the left boundary of R and the rightmost point of I lies on the right boundary of R .

COROLLARY 3.3. *Let $R = [r] \times [s]$, and let $I = [i_1, i_2] \times [j_1, j_2]$ be an interval in R such that $i_1 = 1$ or $j_2 = s$, and $j_1 = 1$ or $i_2 = r$.*

- (a) *There exists a bijection from $\mathcal{P}_I^{(k)}$ to $\mathcal{S}_{i_1+j_1+k-1, i_2+j_1}^{(j_2-j_1-k+1)}$ that divides weight by w_I , so that*

$$W_{i_1+j_1+k-1, i_2+j_1}^{(j_2-j_1-k+1)} = \frac{w_I^{(k)}}{w_I}.$$

- (b) *The following equality holds:*

$$w_I^{(k)} = \frac{W_{i_1+j_1+k-1, i_2+j_1}^{(j_2-j_1-k+1)}}{W_{i_1+j_1-1, i_2+j_1}^{(j_2-j_1+1)}}.$$




FIGURE 6. Illustration of \mathcal{G}_R and \mathcal{G}_I (dotted) for $R = [5] \times [7]$ and $I = [3, 5] \times [2, 7]$ with $k = 2$ as in Corollary 3.3 (note $j_2 = s$ and $i_2 = r$). Nonintersecting paths from $\{P_6, \dots, P_9\}$ to $\{Q_7, \dots, Q_{10}\}$ in \mathcal{G}_R must contain the horizontal steps shown.

Proof. Any nonintersecting paths in \mathcal{G}_R starting at $P_{i_1+j_1+k-1}, \dots, P_{i_1+j_2-1}$ must begin with horizontal steps until reaching row i_1 at points $(i_1, j_1+k), \dots, (i_1, j_2)$. See Figure 6. (This statement is trivial if $i_1 = 1$.) Similarly, any nonintersecting paths ending at $Q_{i_2+j_1}, \dots, Q_{i_2+j_2-k}$ must pass through row $i_2 + 1$ at $(i_2 + 1, j_1), \dots, (i_2 + 1, j_2 - k)$ and end with horizontal steps. (This is trivial if $i_2 = r$.) The remaining parts of the paths lie inside the subgraph of \mathcal{G}_R from rows i_1 through $i_2 + 1$ and columns j_1 through j_2 , which is isomorphic as a weighted graph to \mathcal{G}_I . Part (a) follows by applying Lemma 3.1 to I and summing over all paths.

For part (b), setting $k = 0$ in part (a) gives $W_{i_1+j_1-1, i_2+j_1}^{(j_2-j_1+1)} = \frac{1}{w_I}$. Combining with part (a) gives the result. \square

In particular, note that the conditions of Corollary 3.3 hold whenever I is an order ideal (when $i_1 = j_1 = 1$) or order filter (when $i_2 = r$ and $j_2 = s$) of R .

3.2. BIRATIONAL ROWMOTION FORMULA. Using the relation between the octahedron recurrence and toggles, we will prove the following birational rowmotion formula.

THEOREM 3.4. Let $R = [r] \times [s]$, $x \in \mathbb{R}_{>0}^R$, and $y = \phi^{-1}(x)$. Fix $(i, j) \in R$.

(a) If $0 \leq k \leq r + s - i - j$, then

$$\rho^{-k}(y)_{ij} = \frac{W_{k+2, i+k+1}^{(j-1)}}{W_{k+1, i+k+1}^{(j)}}.$$

(b) For all $k \in \mathbb{Z}$,

$$\rho^k(y)_{ij} = \frac{1}{\rho^{k-i-j+1}(y)_{r+1-i, s+1-j}}.$$

In particular, if $0 < k < i + j$, then the right hand side can be computed using part (a).

(c) For all $k \in \mathbb{Z}$, $\rho^{k+r+s}(y) = \rho^k(y)$. In other words, the action of rowmotion on R has order $r + s$.




FIGURE 7. $R = [2] \times [3]$, \mathcal{G}_R , and the corresponding array $W = W_{ij}^{(k)}$ for $1 \leq k \leq 5$, where \bar{a} denotes a^{-1} for readability. (All entries at height 0 equal 1, and all other entries not shown are 0.) Red and blue lines indicate quotients used to compute $\rho^0(y) = y = \phi^{-1}(x)$ and $\rho^{-1}(y)$ (apart from the topmost label), respectively.

Theorem 3.4 can be used to find an explicit formula for the entries of any power of ρ applied to y . (Using parts (a) and (b) one can compute $\rho^k(y)_{ij}$ whenever $i + j - r - s \leq k < i + j$, and part (c) can be used to bring k into this range.) While the birational rowmotion formulas given in [19] are stated in terms of complements of paths in R , the formulation given here can be seen to be equivalent using Corollary 3.3.

Geometrically, Theorem 3.4(a) states that one can find the values of $\rho^{-k}(y)_{ij}$ as quotients of nearby entries inside the array W , and increasing the value of k corresponds to a translation in the direction $(1, 1, 0)$ as long as these entries remain inside the defined pyramidal region of W . Upon passing outside this region, one needs to use part (b) to relocate inside the pyramid to the entries corresponding to the antipodal point of R . See Figure 7.

EXAMPLE 3.5. Let $R = [2] \times [3]$. Let x be the labeling of R as in Figure 7 with the array $(W_{ij}^{(k)})$ as previously described. When $(i, j) = (2, 2)$, Theorem 3.4(a) gives

$$\begin{aligned} \rho^0(y)_{22} &= \frac{W_{23}^{(1)}}{W_{13}^{(2)}} = \frac{b^{-1} + c^{-1}}{(abcd)^{-1}} = acd + abd, \\ \rho^{-1}(y)_{22} &= \frac{W_{34}^{(1)}}{W_{24}^{(2)}} = \frac{d^{-1} + e^{-1}}{(cdef)^{-1}} = cef + cdf. \end{aligned}$$

To compute $\rho^{-2}(y)_{22}$, $r + s - i - j = 1 < 2$, so part (a) does not apply. Instead, we apply parts (c) and (b) first to obtain

$$\begin{aligned} \rho^{-2}(y)_{22} &= \rho^3(y)_{22} = \frac{1}{\rho^0(y)_{12}} = \frac{W_{12}^{(2)}}{W_{22}^{(1)}} \\ &= (ac)^{-1}, \\ \rho^{-3}(y)_{22} &= \rho^2(y)_{22} = \frac{1}{\rho^{-1}(y)_{12}} = \frac{W_{23}^{(2)}}{W_{33}^{(1)}} \\ &= (bd)^{-1} + (be)^{-1} + (ce)^{-1}, \\ \rho^{-4}(y)_{22} &= \rho(y)_{22} = \frac{1}{\rho^{-2}(y)_{12}} = \frac{W_{34}^{(2)}}{W_{44}^{(1)}} \\ &= (df)^{-1}. \end{aligned}$$

We will prove each part of Theorem 3.4 separately. Part (a) follows primarily from Lemma 2.14, though some care is needed along the boundary of R .

Proof of Theorem 3.4(a). Let $z_{ij}^{(k)} = \frac{W_{k+2, i+k+1}^{(j-1)}}{W_{k+1, i+k+1}^{(j)}}$, which we wish to equal $\rho^{-k}(y)_{ij}$ for $0 \leq k \leq r + s - i - j$. We proceed by induction on k . When $k = 0$, $\rho^0(y)_{ij} = y_{ij} = \phi^{-1}(x)_{ij}$ is the total weight of all maximal chains in $[i] \times [j]$ (with respect to the labeling x). By Corollary 3.3(b), this is $\frac{W_{2, i+1}^{(j-1)}}{W_{1, i+1}^{(j)}} = z_{ij}^{(0)}$, as desired.

For the inductive step, assume $k > 0$ and suppose that the claim is true for $\rho^{-k+1}(y)$ as well as for $\rho^{-k}(y)_{i'j'}$ when $(i', j') < (i, j)$. The value of $\rho^{-k}(y)$ is obtained by applying toggles from bottom to top on $\rho^{-k+1}(y)$, so the toggle at (i, j) gives

$$(*) \quad \rho^{-k}(y)_{ij} = \left(\sum_{(i', j') \leq (i, j)} \rho^{-k}(y)_{i'j'} \right) \left(\sum_{(i', j') \geq (i, j)} \rho^{-k+1}(y)_{i'j'} \right) \cdot \frac{1}{\rho^{-k+1}(y)_{ij}},$$

where an empty sum is replaced with 1.

We claim that we can replace the first sum in $(*)$ with $z_{i, j-1}^{(k)} + z_{i-1, j}^{(k)}$. If $i > 1$ and $j > 1$, then this is immediate by induction. Otherwise note that

$$z_{i0}^{(k)} = \frac{W_{k+2, i+k+1}^{(-1)}}{W_{k+1, i+k+1}^{(0)}} = \frac{0}{1} = 0, \quad z_{0j}^{(k)} = \frac{W_{k+2, k+1}^{(j-1)}}{W_{k+1, k+1}^{(j)}} = \frac{\delta_{j1}}{1} = \delta_{j1}$$

by Proposition 3.2(a) and (b). Thus if exactly one of i and j is 1, then one of $z_{i, j-1}^{(k)}$ and $z_{i-1, j}^{(k)}$ equals 0 while the other is the only term in the first sum in $(*)$ by induction. If instead $i = j = 1$, then the empty sum in $(*)$ is replaced with $1 = 0 + \delta_{11} = z_{10}^{(k)} + z_{01}^{(k)}$ as needed.

Similarly, we claim that we can replace the second (parallel) sum in (*) with either $z_{i,j+1}^{(k-1)} \# z_{i+1,j}^{(k-1)}$, or just one of these terms if the other is undefined. If $i < r$ and $j < s$, then this is again immediate by induction (since the inequality $k - 1 \leq r + s - (i + j + 1)$ holds). If $i = r$ and $j < s$, then $z_{r+1,j}^{(k-1)}$ is undefined since $W_{k,r+k+1}^{(j)} = 0$ by Proposition 3.2(a), so we are left with $z_{r,j+1}^{(k-1)} = \rho^{-k+1}(y)_{r,j+1}$ by induction. If instead $i < r$ and $j = s$, then $z_{i,s+1}^{(k-1)}$ is undefined since $W_{k,i+k}^{(s+1)} = 0$ by Proposition 3.2(c), which leaves just $z_{i+1,s}^{(k-1)} = \rho^{-k+1}(y)_{i+1,s}$ by induction. Finally, $i = r$ and $j = s$ cannot both occur since $0 < k \leq r + s - i - j$.

The final factor in (*) is equal to $\frac{1}{z_{ij}^{(k-1)}}$ by induction. The result then follows by comparing (*) to Lemma 2.14, using Remark 2.15 when needed. □

Next we prove part (b). In some sense, this reflects two symmetries: toggling is respected by dualizing a poset and inverting each label, and the octahedron recurrence is respected by reflecting over the plane perpendicular to the main diagonal.

Proof of Theorem 3.4(b). Note that if $0 < k < i + j$, then $0 \leq i + j - k - 1 \leq r + s - (r + 1 - i) - (s + 1 - j)$, so the right hand side of (b) will satisfy the condition of part (a), as claimed.

Via the change of coordinates $z = \rho^{k-1}(y)$, it suffices to prove the case when $k = 1$. By part (a),

$$\rho^{2-i-j}(y)_{r+1-i,s+1-j} = \frac{W_{i+j,r+j}^{(s-j)}}{W_{i+j-1,r+j}^{(s-j+1)}}.$$

By Corollary 3.3, this is the total weight of all maximal chains in $[i, r] \times [j, s]$ in R (with respect to the labeling x). Thus

$$\rho^{2-i-j}(y)_{r+1-i,s+1-j} = (\phi^*)^{-1}(x)_{ij} = \frac{1}{\rho(y)_{ij}}$$

by Lemma 2.11. □

Finally, we deduce part (c) from part (b).

Proof of Theorem 3.4(c). Apply Theorem 3.4(b) twice:

$$\begin{aligned} \rho^k(y)_{i,j} &= (\rho^{k-i-j+1}(y)_{r+1-i,s+1-j})^{-1} \\ &= \left[\left(\rho^{k-i-j+1-(r+1-i)-(s+1-j)+1}(y)_{i,j} \right)^{-1} \right]^{-1} \\ &= \rho^{k-r-s}(y)_{i,j}. \end{aligned} \quad \square$$

In the next two sections, we will show how to use the perspective relating rowmotion and the octahedron recurrence to prove various identities about rowmotion.

4. A CHAIN SHIFTING LEMMA

In this section we will use the birational rowmotion formula to derive a chain shifting lemma, which is a birational generalization of the rowmotion action on the Stanley–Thomas word appearing in [15].




FIGURE 8. Two labelings x and $z = \phi \circ \rho^{-1} \circ \phi^{-1}(x)$.

4.1. CHAIN SHIFTING. Recall from Section 3.1 that if I is an interval of R and x is a labeling of R , then $w_I^{(k)}(x)$ is the total weight of the collections of k nonintersecting paths in $\mathcal{P}_I^{(k)}$. We will prove the following lemma relating these sums in two labelings x and $\phi \circ \rho^{-1} \circ \phi^{-1}(x)$.

LEMMA 4.1 (Chain shifting). *Let $R = [r] \times [s]$ and let $1 < u \leq v \leq s$. Define intervals $I = [r] \times [u, v]$ and $I' = [r] \times [u - 1, v - 1]$ of R . Then for any labeling $x \in \mathbb{R}_{>0}^R$,*

$$w_I^{(k)}(x) = w_{I'}^{(k)}(\phi \circ \rho^{-1} \circ \phi^{-1}(x)).$$

(The symmetric statement obtained by reflecting R also holds.)

Note that the intervals I and I' differ only by shifting by one unit in R .

EXAMPLE 4.2. Consider the labeling x of $R = [2] \times [3]$ and its image $z = \phi \circ \rho^{-1} \circ \phi^{-1}(x)$ in Figure 8. Then

$$z_{11}z_{21} = \frac{bc}{b+c} \cdot \frac{d(b+c)}{b} = cd = x_{12}x_{22}.$$

Similarly, we can compute all of the products along the rows and columns of R :

$$\begin{aligned} z_{12}z_{22} &= x_{13}x_{23} & z_{13}z_{23} &= (x_{11}x_{12}x_{13})^{-1} \\ (z_{11}z_{12}z_{13})^{-1} &= (x_{21}x_{22}x_{23})^{-1} & (z_{21}z_{22}z_{23})^{-1} &= x_{11}x_{21} \end{aligned}$$

From the above computation, we see that $\phi \circ \rho^{-1} \circ \phi^{-1}$ rotates the birational Stanley–Thomas word

$$(x_{11}x_{21}, \quad x_{12}x_{22}, \quad x_{13}x_{23}, \quad (x_{11}x_{12}x_{13})^{-1}, \quad (x_{21}x_{22}x_{23})^{-1}).$$

However, the shifting of chain sums happens more generally for sums of chains within subrectangles of R . For instance, if $I = [2] \times [2, 3]$, $I' = [2] \times [1, 2]$, and $k = 1$, then we have

$$\begin{aligned} z_{11}z_{21}z_{22} + z_{11}z_{12}z_{22} &= \frac{bc}{b+c} \cdot \frac{f(bd+cd+ce)}{d(b+c)} \left(\frac{d(b+c)}{b} + \frac{de(b+c)}{bd+cd+ce} \right) \\ &= \frac{cf(bd+cd+ce)}{b+c} + \frac{bcdf}{b+c} \\ &= cd f + ce f \\ &= x_{12}x_{22}x_{23} + x_{12}x_{13}x_{22}. \end{aligned}$$

We now prove Lemma 4.1. As we will see, it can be thought of as a manifestation of the translation invariance of the octahedron recurrence.

Proof of Lemma 4.1. Let x be a labeling of R , let $y = \phi^{-1}(x)$, and let $(W_{ij}^{(k)})$ be the corresponding three-dimensional array. Also define $\tilde{x} = \phi \circ \rho^{-1} \circ \phi^{-1}(x)$, $\tilde{y} = \phi^{-1}(\tilde{x}) = \rho^{-1}(y)$, and $(\tilde{W}_{ij}^{(k)})$ the corresponding array.

We claim that $\widetilde{W}_{ij}^{(k)} = W_{i+1,j+1}^{(k)}$ for $1 \leq i, j \leq r+s-k$. Note that by the octahedron recurrence/Dodgson condensation formula, it suffices to prove the case $k = 1$ since the entries for $k > 1$ are determinants of submatrices of these entries at height 1.

Assume $k = 1$. If $i \geq j$, then the claim follows by Proposition 3.2. If $i < j$, then applying Theorem 3.4 twice gives

$$\widetilde{W}_{ij}^{(1)} = \frac{1}{\rho^{-i+1}(\widetilde{y})_{j-i,1}} = \frac{1}{\rho^{-i}(y)_{j-i,1}} = W_{i+1,j+1}^{(1)},$$

which proves the claim.

To complete the proof, we need only observe that applying Corollary 3.3(b) to the two sides of the desired equality expresses them as quotients of two entries of the form $W_{i+1,j+1}^{(k)}$ or $\widetilde{W}_{ij}^{(k)}$, respectively, which are then equal by the claim. \square

4.2. GENERALIZED STANLEY–THOMAS WORDS. Given Lemma 4.1, it is natural to want to generalize the definition of the Stanley–Thomas word to include more words that are cyclically shifted by the action of rowmotion.

The usual birational Stanley–Thomas word is obtained as the orbit of $\prod_{j=1}^s x_{1j} = \phi^{-1}(x)_{1s}$ under the action of $\phi \circ \rho \circ \phi^{-1}$ on x , or, put another way, the orbit of y_{1s} under the action of ρ on $y = \phi^{-1}(x)$. We can generalize this by replacing y_{1s} with any other value y_{is} or y_{rj} lying on the upper boundary of R .

DEFINITION 4.3. Let $x \in \mathbb{R}_{>0}^R$, and let $y = \phi^{-1}(x)$. A generalized Stanley–Thomas word for x is a sequence of one of the following two forms:

$$\begin{aligned} ST_i(x) &= (y_{is}, \quad \rho^{-1}(y)_{is}, \quad \rho^{-2}(y)_{is}, \quad \dots, \quad \rho^{-r-s+1}(y)_{is}), \\ \overline{ST}_j(x) &= (y_{rj}, \quad \rho^{-1}(y)_{rj}, \quad \rho^{-2}(y)_{rj}, \quad \dots, \quad \rho^{-r-s+1}(y)_{rj}). \end{aligned}$$

The usual birational Stanley–Thomas word is therefore ST_1 . The reason for choosing these words in particular is that they have a clean description in terms of x . We consider the case of ST_i below; the other case is similar.

Recall that $y_{is} = \phi^{-1}(x)_{is} = w_{[i] \times [s]}^{(1)}(x)$. By Lemma 4.1,

$$\rho^{-1}(y)_{is} = w_{[i] \times [s]}^{(1)}(\phi \circ \rho^{-1}(y)) = w_{[2,i+1] \times [s]}^{(1)}(x).$$

Iterating, we find that for $0 \leq k \leq r - i$,

$$\rho^{-k}(y)_{is} = w_{[k+1,k+i] \times [s]}^{(1)}(x).$$

To find the remaining coordinates, we apply Theorem 3.4 (c), (b), and (a) in order to find that, for $r - i < k < r + s$,

$$\rho^{-k}(y)_{is} = \rho^{r+s-k}(y)_{is} = \frac{1}{\rho^{r-i-k+1}(y)_{r+1-i,1}} = W_{k+i-r,k+1}^{(1)}.$$

While $W_{k+i-r,k+1}^{(1)}$ is defined in terms of paths in \mathcal{G}_R , it is also easy to describe it in terms of the labeling x directly. Choosing the northwest edges in a path in \mathcal{G}_R from P_{k+i-r} to Q_{k+1} corresponds to choosing elements of R at ranks $k+i-r+1, \dots, k+1$ traveling to the northwest (i.e. with increasing first coordinate), not necessarily using the edges of R . In other words, define

$$\omega_{ab} = \omega_{ab}(x) = \sum (x_{i_a j_a} x_{i_{a+1} j_{a+1}} \cdots x_{i_b j_b})^{-1},$$

where the sum ranges over all sequences $(i_a, j_a), (i_{a+1}, j_{a+1}), \dots, (i_b, j_b) \in R$ such that $i_t + j_t = t$ for all t , and $i_a < i_{a+1} < \dots < i_b$ (or equivalently $j_a \geq j_{a+1} \geq \dots \geq j_b$). Then we have proved the following proposition.




FIGURE 9. Computation of ST_2 in $R = [3] \times [3]$ as in Example 4.5. In the first two diagrams, the sums of the weights of the maximum length chains in the blue posets are $w_{[1,2] \times [3]}^{(1)}$ and $w_{[2,3] \times [3]}^{(1)}$, respectively. In the remaining four diagrams, the sums of the reciprocals of the weights of the maximum length chains in the red posets are ω_{23} , ω_{34} , ω_{45} , and ω_{56} , respectively.

PROPOSITION 4.4. For any $x \in \mathbb{R}_{>0}^R$, the generalized Stanley–Thomas word $ST_i = ST_i(x)$ is given by

$$ST_i = (w_{[1,i] \times [s]}^{(1)}, w_{[2,i+1] \times [s]}^{(1)}, \dots, w_{[k+1,k+i] \times [s]}^{(1)}, \omega_{2,r-i+2}, \omega_{3,r-i+3}, \dots, \omega_{i+s,r+s}).$$

An analogous formula for \overline{ST}_j can be obtained by reflecting R .

EXAMPLE 4.5. Let $R = [3] \times [3]$ and $x \in \mathbb{R}_{>0}^R$. The generalized Stanley–Thomas word ST_2 is

$$ST_2 = (w_{[1,2] \times [3]}^{(1)}, w_{[2,3] \times [3]}^{(1)}, \omega_{23}, \omega_{34}, \omega_{45}, \omega_{56}).$$

In the x -coordinates we have the following:

$$\begin{aligned} w_{[1,2] \times [3]}^{(1)} &= x_{11}x_{21}x_{22}x_{23} + x_{11}x_{12}x_{22}x_{23} + x_{11}x_{12}x_{13}x_{23}, \\ w_{[2,3] \times [3]}^{(1)} &= x_{21}x_{31}x_{32}x_{33} + x_{21}x_{22}x_{32}x_{33} + x_{21}x_{22}x_{23}x_{33}, \\ \omega_{23} &= \frac{1}{x_{11}x_{21}}, \\ \omega_{34} &= \frac{1}{x_{12}x_{22}} + \frac{1}{x_{12}x_{31}} + \frac{1}{x_{21}x_{31}}, \\ \omega_{45} &= \frac{1}{x_{13}x_{23}} + \frac{1}{x_{13}x_{32}} + \frac{1}{x_{22}x_{32}}, \\ \omega_{56} &= \frac{1}{x_{23}x_{33}}. \end{aligned}$$

See Figure 9 for a visualization of this example.

As noted in [15], the cyclic rotation of ST_1 is not sufficient to uniquely determine birational rowmotion. However, the cyclic rotation of all ST_i and \overline{ST}_j does uniquely determine birational rowmotion, and in fact the chain shifting lemma alone nearly suffices. We make this statement precise in Section 5.




FIGURE 10. An example of the map $A \mapsto RSK(A)$ in the classical setting. See Example 5.1.

5. BIRATIONAL RSK AND GREENE’S THEOREM

In this section, we will define birational RSK in terms of toggles and show how our perspective gives a simple proof of the birational version of Greene’s theorem.

5.1. CLASSICAL RSK. We first review some background on the classical RSK correspondence, which gives a bijection between nonnegative integer matrices A and pairs of semistandard tableaux (P, Q) of the same shape λ . (See, for instance, [24] for more details.)

In the case when A is the matrix of a permutation π , Greene’s theorem [8] states that $\lambda_1 + \dots + \lambda_k$ is the maximum size of a union of k increasing subsequences of π . In fact, one can use Greene’s theorem to compute not just the shape of P and Q but the entire tableaux: the shape $P^{\leq m}$ formed by the entries at most m in P corresponds to the permutation formed by the letters $1, \dots, m$ in π , while the shape $Q^{\leq m}$ corresponds to the permutation formed by the first m letters of π .

When A is a general $n \times n$ nonnegative integer matrix, a routine standardization argument gives the following generalization of Greene’s theorem: $\lambda_1 + \dots + \lambda_k$ is the maximum weight of k nonintersecting paths (traveling weakly southeast) from $(1, 1), \dots, (1, k)$ to $(n, n - k + 1), \dots, (n, n)$ in A , where the weight of a path is the sum of the entries it contains. (As noted in [6], this result appears to be somewhat folklore, but see [20, Thm. 2.5] as well as [17, Thm. 12], [24, Thm. 4.8.10].) As in the permutation case, this can be used to characterize the entire P and Q tableaux since $P^{\leq m}$ (resp. $Q^{\leq m}$) corresponds to the submatrix formed by the first m columns (resp. rows) of A .

One way to see the impact of Greene’s theorem more directly is to transform P and Q into Gelfand-Tsetlin patterns and glue them along their top rows to form an $n \times n$ matrix with weakly increasing rows and columns that we denote by $RSK(A)$. By construction, this matrix has the property that, for $1 \leq k \leq i, j \leq n$ with either $i = n$ or $j = n$,

$$\sum_{t=0}^{k-1} RSK(A)_{i-t, j-t}$$

is the maximum weight of k nonintersecting paths from $(1, 1), \dots, (1, k)$ to $(i, j - k + 1), \dots, (i, j)$ in A .

EXAMPLE 5.1. Figure 10 depicts a 4×4 matrix $A = (a_{ij})$ and its image under the classical RSK correspondence as a pair of semistandard tableaux P and Q of the same




FIGURE 11. Computing $RSK(x)$ by applying toggles to $y = \phi^{-1}(x)$. At each step, the highlighted elements are toggled from bottom to top.

shape. These tableaux are then turned into triangular Gelfand-Tsetlin patterns (the rows of which give the shapes of the tableaux $P^{\leq m}$ and $Q^{\leq m}$ for $m = 4, 3, 2, 1$). These two Gelfand-Tsetlin patterns are then glued together to form the matrix $RSK(A)$.

By Greene’s theorem, the entries of $RSK(A)$ tell us about weights of nonintersecting paths in A . For example:

- $RSK(A)_{34} = 5$ is the maximum weight of a path from a_{11} to a_{34} ;
- $RSK(A)_{34} + RSK(A)_{23} = 5 + 2 = 7$ is the maximum weight of two nonintersecting paths from a_{11} and a_{12} to a_{33} and a_{34} ; and
- $RSK(A)_{34} + RSK(A)_{23} + RSK(A)_{12} = 5 + 2 + 1 = 8$ is the maximum weight of three nonintersecting paths from a_{11}, a_{12}, a_{13} to a_{32}, a_{33}, a_{34} , which is just the sum of all the entries in the first three rows of A .

In the next section, we will see how the map $A \mapsto RSK(A)$ can be generalized to the birational setting.

5.2. BIRATIONAL RSK. For any interval $I \subseteq R$, denote by ρ_I the map on labelings of R given by the composition of toggles at the elements of I (applied in the order of a linear extension from top to bottom).

DEFINITION 5.2. Let $R = [r] \times [s]$, and let $m = \min\{r, s\} - 1$. The birational RSK map on labelings of R is given by

$$RSK = \rho_{[r-m] \times [s-m]}^{-1} \circ \cdots \circ \rho_{[r-2] \times [s-2]}^{-1} \circ \rho_{[r-1] \times [s-1]}^{-1} \circ \phi^{-1}.$$

We will compare this to other formulations of birational RSK below, but for now, this can be treated as a definition. We will abuse notation and denote the piecewise-linear version of this map by RSK as well.

EXAMPLE 5.3. Let $R = [3] \times [4]$, and let $x \in \mathbb{R}_{\geq 0}^R$ and $y = \phi^{-1}(x)$. Then

$$RSK(x) = \rho_{[1] \times [2]}^{-1} \circ \rho_{[2] \times [3]}^{-1}(y) = (t_{12} \circ t_{11}) \circ (t_{23} \circ t_{22} \circ t_{21} \circ t_{13} \circ t_{12} \circ t_{11})(y).$$

See Figure 11. Note that each toggle involves essentially the same calculation that is done when applying some power of ρ^{-1} to y . As a result, each label of $RSK(x)$ is equal to a label in some $\rho^{-k}(y)$. These can then be computed using Theorem 3.4.

In general, one can always describe the entries of $RSK(x)$ using the action of rowmotion on all of R as in the following proposition.

PROPOSITION 5.4. $RSK(x)_{r-i, s-j} = \rho^{-\min\{i, j\}} \circ \phi^{-1}(x)_{r-i, s-j}$.

Proof. Let $y = \phi^{-1}(x)$. We claim that $\rho_{[r-k] \times [s-k]}^{-1} \circ \cdots \circ \rho_{[r-1] \times [s-1]}^{-1}(y)$ and $\rho^{-k}(y)$ agree on $[r - k] \times [s - k]$. Indeed, suppose inductively that the claim holds for $k - 1$.

Since any neighbor of an element in $[r - k] \times [s - k]$ lies in $[r - k + 1] \times [s - k + 1]$, applying $\rho_{[r-k, s-k]}^{-1}$ to both $\rho_{[r-k+1] \times [s-k+1]}^{-1} \circ \dots \circ \rho_{[r-1] \times [s-1]}^{-1}(y)$ and $\rho^{-k+1}(y)$ yields labelings that agree on $[r - k] \times [s - k]$. Then applying the remaining toggles in ρ^{-1} (at elements of $R \setminus ([r - k] \times [s - k])$) to the latter also does not affect any of these values, proving the claim.

If $\min\{i, j\} = k$, then $\rho_{[r-k'] \times [s-k']}^{-1}$ does not affect the label at $(r - i, s - j)$ for any $k' > k$, so the result follows from the claim. \square

We are now ready to prove the following theorem, which will serve as a birational analogue to Greene's theorem as described in Section 5.1.

THEOREM 5.5 (Birational Greene's theorem). *Let $R = [r] \times [s]$, and choose $(i, j) \in R$ such that either $i = r$ or $j = s$. Then for any $x \in \mathbb{R}_{>0}^R$ and $1 \leq k \leq \min\{i, j\} + 1$,*

$$\prod_{t=0}^{k-1} RSK(x)_{i-t, j-t} = w_{[i] \times [j]}^{(k)}(x).$$

Proof. Let $x \in \mathbb{R}_{>0}^R$ and let $y = \phi^{-1}(x)$. By Proposition 5.4, since

$$\min\{r - (i - t), s - (j - t)\} = \min\{r - i, s - j\} + t = t,$$

we have

$$RSK(x)_{i-t, j-t} = \rho^{-t}(y)_{i-t, j-t}.$$

Theorem 3.4(a) then gives

$$\rho^{-t}(y)_{i-t, j-t} = \frac{W_{t+2, i+1}^{(j-t-1)}}{W_{t+1, i+1}^{(j-t)}}.$$

We then have the telescoping product

$$\prod_{t=0}^{k-1} RSK(x)_{i-t, j-t} = \prod_{t=0}^{k-1} \rho^{-t}(y)_{i-t, j-t} = \frac{W_{k+1, i+1}^{(j-k)}}{W_{1, i+1}^{(j)}} = w_{[i] \times [j]}^{(k)}(x)$$

by Corollary 3.3. \square

Note that Theorem 5.5 uniquely characterizes the map RSK in terms of the values of $w_{[i] \times [j]}^{(k)}(x)$ where $i = r$ or $j = s$. To see why this can be considered to be a birational version of Greene's theorem, consider the tropicalization of Theorem 5.5:

$$\sum_{t=0}^{k-1} RSK(x)_{i-t, j-t} = \max_{\mathcal{L} \in \mathcal{P}_{[i] \times [j]}^{(k)}} \sum_{z \in \mathcal{L}} x_z.$$

This corresponds directly to the description of $RSK(A)$ given in Section 5.1. We can therefore treat this as a proof that the map RSK as defined above is a birational analogue to the classical RSK correspondence.

EXAMPLE 5.6. Suppose $R = [4] \times [4]$ and $x \in \mathbb{R}_{>0}^R$. According to Theorem 5.5, we can compute the labels of $RSK(x)$ in terms of the weights of nonintersecting paths in x . For instance,

$$\begin{aligned} w_{[3] \times [4]}^{(1)}(x) &= RSK(x)_{34}, \\ w_{[3] \times [4]}^{(2)}(x) &= RSK(x)_{34} \cdot RSK(x)_{23}, \\ w_{[3] \times [4]}^{(3)}(x) &= RSK(x)_{34} \cdot RSK(x)_{23} \cdot RSK(x)_{12}, \end{aligned}$$

from which these three labels of $RSK(x)$ can easily be determined by taking quotients. Note that these three equations tropicalize to the description of the corresponding

entries of the classical RSK map using Greene’s theorem as in Section 5.1. (Compare to the bullet points in Example 5.1.)

This implies that we can compute the classical RSK map by using the piecewise-linear version of Definition 5.2. For instance, treating the matrix A in Figure 10 as a labeling of the rectangle $R = [4] \times [4]$, we can compute $RSK(A)$ as follows. First apply the inverse transfer map ϕ^{-1} :

$$A = \begin{bmatrix} 0 & 0 & 0 & 2 \\ 2 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix} \xrightarrow{\phi^{-1}} \begin{bmatrix} 0 & 0 & 0 & 2 \\ 2 & 2 & 3 & 3 \\ 2 & 3 & 4 & 5 \\ 3 & 4 & 4 & 5 \end{bmatrix}.$$

Then successively apply the partial rowmotion maps as indicated. (At each step, the red entries are toggled starting in the upper left corner.)

$$\begin{bmatrix} 0 & 0 & 0 & 2 \\ 2 & 2 & 3 & 3 \\ 2 & 3 & 4 & 5 \\ 3 & 4 & 4 & 5 \end{bmatrix} \xrightarrow{\rho_{[3] \times [3]}^{-1}} \begin{bmatrix} 0 & 0 & 2 & 2 \\ 0 & 1 & 2 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 5 \end{bmatrix} \xrightarrow{\rho_{[2] \times [2]}^{-1}} \begin{bmatrix} 0 & 1 & 2 & 2 \\ 1 & 2 & 2 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 5 \end{bmatrix} \xrightarrow{\rho_{[1] \times [1]}^{-1}} \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 2 & 2 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 5 \end{bmatrix}.$$

The resulting matrix $RSK(A)$ is the same as in Figure 10.

5.3. RELATION TO PRIOR WORK. Our construction of birational RSK using the octahedron recurrence is very similar to the approach taken by Danilov–Koshevoy [3] (although their construction differs from the standard one by a symmetry of A). Our construction is also necessarily equivalent to the description of “tropical RSK” (sometimes also “geometric RSK”) given by Noumi–Yamada [20] (motivated by Kirillov [16]) since their map is also constructed to satisfy Theorem 5.5. An analysis of the Noumi–Yamada construction via local toggles can also be found in [21].

In [6], a birational version of RSK is described using the following piecewise-linear analogue of RSK defined in [11, 22]. Let $x \in \mathcal{C}(R)$ be a point in the chain polytope of $R = [r] \times [s]$. We construct $y = rsk(x) \in \mathcal{O}(R)$ in the order polytope of R using the following procedure.

- (1) Set $y_{11} = x_{11}$.
- (2) Choose an element $p \in R$ such that y_q has been defined for all $q < p$. Set $y_p = x_p + \max_{q < p} y_q$, and then toggle y at all elements below p in the same file.
- (3) Repeat the previous step until all coordinates of y have been defined.

In step (2), the elements p can be considered in the order of any linear extension of R . It is straightforward to verify that the result is independent of the choice of linear extension. We can also extend rsk in the obvious way to all of $\mathbb{R}_{\geq 0}^R$.

EXAMPLE 5.7. Consider the matrix A from Examples 5.1 and 5.6. We compute $rsk(A)$ by considering elements in order of rank, taking the elements at each rank simultaneously (entries colored in red are toggled at the following step):

$$\begin{bmatrix} 0 \\ \\ \\ \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 0 \\ 2 \\ \\ \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 2 & 2 \\ 2 \\ \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 0 & 0 & 2 \\ 0 & 2 & 3 \\ 2 & 3 \\ 3 \end{bmatrix} \\ \rightarrow \begin{bmatrix} 0 & 0 & 2 & 2 \\ 0 & 1 & 3 & 3 \\ 1 & 3 & 4 \\ 3 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 & 2 & 2 \\ 1 & 1 & 2 & 3 \\ 1 & 2 & 4 & 5 \\ 3 & 4 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 2 & 2 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 5 \end{bmatrix}.$$

Note that the result agrees with the computations of $RSK(A)$ from earlier.

In fact, this map is the same as the piecewise-linear version of the map RSK defined in the previous section.

PROPOSITION 5.8. $RSK = rsk$.

Proof. When performing step (2) above, none of the toggled elements form a cover relation with any element that has yet to be considered. Since the value of each new y_p depends only on the values y_q for $q < p$, the toggles at the previous steps do not affect the initial value of y_p . Hence we can apply all the toggles after first assigning all y_p , which is equivalent to first applying ϕ^{-1} to x .

Without loss of generality, we may assume $r \leq s$. If $r = 1$, then the rectangle is a chain and no toggles are applied, so $rsk = \phi^{-1} = RSK$.

Suppose inductively that on $[r - 1] \times [s]$, $rsk = \rho_{[1] \times [s-r+2]}^{-1} \circ \cdots \circ \rho_{[r-2] \times [s-1]}^{-1} \circ \phi^{-1}$, and consider the rectangle $R = [r] \times [s]$ for $r \leq s$. After applying rsk on the order ideal $[r - 1] \times [s]$, to complete rsk for R we must apply toggles at all elements in $[r - 1] \times [s]$ that lie in files $2 - r, 3 - r, \dots, s - r$. Since toggles at elements that are not adjacent commute, we can toggle row by row instead of file by file. Write $I_i = \{i\} \times [s - r + i] \subseteq R$ for $i < r$. We then find that on R ,

$$\begin{aligned} rsk &= \left(\rho_{I_1}^{-1} \circ \rho_{I_2}^{-1} \cdots \circ \rho_{I_{r-1}}^{-1} \right) \circ \rho_{[1] \times [s-r+2]}^{-1} \circ \cdots \circ \rho_{[r-2] \times [s-1]}^{-1} \circ \phi^{-1} \\ &= \left(\rho_{I_1}^{-1} \right) \circ \left(\rho_{I_2}^{-1} \circ \rho_{[1] \times [s-r+2]}^{-1} \right) \circ \cdots \circ \left(\rho_{I_{r-1}}^{-1} \circ \rho_{[r-2] \times [s-1]}^{-1} \right) \circ \phi^{-1} \\ &= \rho_{[1] \times [s-r+1]}^{-1} \circ \rho_{[2] \times [s-r+2]}^{-1} \circ \cdots \circ \rho_{[r-1] \times [s-1]}^{-1} \circ \phi^{-1} \\ &= RSK. \end{aligned} \quad \square$$

5.4. RELATION TO CHAIN SUMS AND STANLEY–THOMAS WORDS. Since RSK is invertible, any $x \in \mathbb{R}_{>0}^R$ is uniquely determined by $RSK(x)$. From Theorem 5.5, it follows that the values of $w_I^{(k)}(x)$, where I has the form $[r] \times [j]$ or $[i] \times [s]$, determine x .

Perhaps even more relevant for our work with rowmotion, we can also characterize x using the types of chain sums that appear in Lemma 4.1 with $k = 1$.

THEOREM 5.9. Let $x \in \mathbb{R}_{>0}^R$. The chain sums $w_I^{(1)}(x)$, where I ranges over intervals of the form $[r] \times [u, v]$ and $[u, v] \times [s]$, uniquely determine x .

Proof. By the Lindström–Gessel–Viennot Lemma on R (using vertex weights), we can express any $w_{[r] \times [j]}^{(k)}$ as a determinant of a matrix with entries of the form $w_{[r] \times [u, v]}^{(1)}$, and similarly any $w_{[i] \times [s]}^{(k)}$ is determined by the values $w_{[u, v] \times [s]}^{(1)}$. By Theorem 5.5, each entry of $RSK(x)$ is a quotient of (or equal to) entries of the form $w_{[r] \times [j]}^{(k)}$ or $w_{[i] \times [s]}^{(k)}$, so the result follows.

(In fact, we do not even need $w_{[r] \times [u, s]}^{(1)}$ for $2 \leq u \leq s$ since $w_{[r] \times [s]}^{(k)}$ can be computed from the other chain sums.) \square

EXAMPLE 5.10. Let $R = [2] \times [3]$, and take $x \in \mathbb{R}_{>0}^R$ to be labeled as in Figure 8. From Theorem 5.9, x is uniquely determined by the following chain sums:

$$\begin{aligned} w_{[2] \times [1, 1]}^{(1)} &= ab, & w_{[2] \times [1, 2]}^{(1)} &= abd + acd, & w_{[2] \times [2, 2]}^{(1)} &= cd, \\ w_{[1, 1] \times [3]}^{(1)} &= ace, & w_{[1, 2] \times [3]}^{(1)} &= abdf + acdf + acef, & w_{[2, 2] \times [3]}^{(1)} &= bdf. \end{aligned}$$

Indeed, from these we can compute

$$w_{[2] \times [1,2]}^{(2)} = \begin{vmatrix} w_{[2] \times [1,1]}^{(1)} & w_{[2] \times [1,2]}^{(1)} \\ 0 & w_{[2] \times [2,2]}^{(1)} \end{vmatrix} = abcd,$$

$$w_{[1,2] \times [3]}^{(2)} = \begin{vmatrix} w_{[1,1] \times [3]}^{(1)} & w_{[1,2] \times [3]}^{(1)} \\ 0 & w_{[2,2] \times [3]}^{(1)} \end{vmatrix} = abcdef.$$

We can then compute $RSK(x)$ using Theorem 5.5:

$$RSK(x)_{11} = \frac{w_{[2] \times [1,2]}^{(2)}}{w_{[2] \times [1,2]}^{(1)}} = \frac{abcd}{abd + acd},$$

$$RSK(x)_{12} = \frac{w_{[1,2] \times [3]}^{(2)}}{w_{[1,2] \times [3]}^{(1)}} = \frac{abcdef}{abdf + acdf + acef},$$

$$RSK(x)_{13} = w_{[1,1] \times [3]}^{(1)} = ace,$$

$$RSK(x)_{21} = w_{[2] \times [1,1]}^{(1)} = ab,$$

$$RSK(x)_{22} = w_{[2] \times [1,2]}^{(1)} = abd + acd,$$

$$RSK(x)_{23} = w_{[1,2] \times [3]}^{(1)} = abdf + acdf + acef.$$

It follows that $RSK(x)$ and hence x are uniquely determined.

In particular, since these chain sums all appear in the generalized Stanley–Thomas words, the following corollary is immediate.

COROLLARY 5.11. *The map $\phi \circ \rho \circ \phi^{-1}$ is the unique function on $\mathbb{R}_{>0}^R$ that cyclically shifts the generalized Stanley–Thomas words ST_i and \overline{ST}_j .*

It follows that the cyclic shifting of the generalized Stanley–Thomas words could be used to prove periodicity of birational rowmotion on rectangles. Of course, presently our method of proving this cyclic shifting yielded a proof of periodicity more directly. However, the techniques in our follow-up work [13] can be used to derive this cyclic shifting in a manner that is more independent of periodicity.

By Lemma 4.1, most of the chain sums $w_I^{(1)}$ considered above get sent to other such sums. It follows that the chain shifting lemma determines “most” of rowmotion in the following sense.

COROLLARY 5.12. *The chain shifting property of $\phi \circ \rho^{-1} \circ \phi^{-1}$ (stated in Lemma 4.1) uniquely determines $RSK(\phi \circ \rho^{-1} \circ \phi^{-1}(x))_{ij}$, where $j - i \neq s - r$.*

Proof. To compute the (i, j) th coordinate of the RSK map where $j - i < s - r$, we need only compute the values of $w_{[r] \times [j+r-i]}^{(k)}$ for $k \leq r - i + 1$ by Theorem 5.5. Since $j + r - i < s$, by Lemma 4.1, $w_{[r] \times [j+r-i]}^{(k)}(\phi \circ \rho^{-1} \circ \phi^{-1}(x)) = w_{[r], [2, j+r-i+1]}^{(k)}(x)$.

The entries with $j - i > s - r$ can likewise be obtained using the reflected version of Lemma 4.1. \square

Acknowledgements. The authors would like to thank Sam Hopkins, Alexander Postnikov, Darij Grinberg, and Tom Roby for useful conversations, as well as the anonymous referees for their helpful comments.

REFERENCES

- [1] A. E. Brouwer and A. Schrijver, *On the period of an operator, defined on antichains*, Mathematisch Centrum, Afdeling Zuivere Wiskunde ZW 24/74, Mathematisch Centrum, Amsterdam, 1974.
- [2] P. J. Cameron and D. G. Fon-Der-Flaass, *Orbits of antichains revisited*, *European J. Combin.* **16** (1995), no. 6, 545–554.
- [3] V. I. Danilov and G. A. Koshevoy, *The octahedron recurrence and RSK-correspondence*, *Sém. Lothar. Combin.* **54A** (2005/07), article no. B54An (16 pages).
- [4] David Einstein and James Propp, *Piecewise-linear and birational toggling*, in 26th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2014), *Discrete Math. Theor. Comput. Sci. Proc.*, AT, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2014, pp. 513–524.
- [5] David Einstein and James Propp, *Combinatorial, piecewise-linear, and birational homomesy for products of two chains*, *Algebr. Comb.* **4** (2021), no. 2, 201–224.
- [6] Miriam Farber, Sam Hopkins, and Wuttisak Trongsirawat, *Interlacing networks: birational RSK, the octahedron recurrence, and Schur function identities*, *J. Combin. Theory Ser. A* **133** (2015), 339–371.
- [7] Ira Gessel and Gérard Viennot, *Determinants, paths, and plane partitions*, 1989, <https://people.brandeis.edu/~gessel/homepage/papers/pp.pdf>.
- [8] Curtis Greene, *An extension of Schensted's theorem*, *Advances in Math.* **14** (1974), 254–265.
- [9] Darij Grinberg and Tom Roby, *Iterative properties of birational rowmotion II: rectangles and triangles*, *Electron. J. Combin.* **22** (2015), no. 3, article no. 3.40 (49 pages).
- [10] Darij Grinberg and Tom Roby, *Iterative properties of birational rowmotion I: generalities and skeletal posets*, *Electron. J. Combin.* **23** (2016), no. 1, article no. 1.33 (40 pages).
- [11] Sam Hopkins, *RSK via local transformations*, 2014, <https://www.samuelhopkins.com/docs/rsk.pdf>.
- [12] Sam Hopkins, *Cyclic sieving for plane partitions and symmetry*, *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), article no. 130 (40 pages).
- [13] Joseph Johnson and Ricky Ini Liu, *Plane partitions and rowmotion on rectangular and trapezoidal posets*, 2023, <https://arxiv.org/abs/2311.07133>.
- [14] Michael Joseph and Tom Roby, *Birational and noncommutative lifts of antichain toggling and rowmotion*, *Algebr. Comb.* **3** (2020), no. 4, 955–984.
- [15] Michael Joseph and Tom Roby, *A birational lifting of the Stanley-Thomas word on products of two chains*, *Discrete Math. Theor. Comput. Sci.* **23** (2021), no. 1, article no. 17 (20 pages).
- [16] Anatol N. Kirillov, *Introduction to tropical combinatorics*, in *Physics and combinatorics*, 2000 (Nagoya), World Sci. Publ., River Edge, NJ, 2001, pp. 82–150.
- [17] C. Krattenthaler, *Growth diagrams, and increasing and decreasing chains in fillings of Ferrers shapes*, *Adv. in Appl. Math.* **37** (2006), no. 3, 404–431.
- [18] Bernt Lindström, *On the vector representations of induced matroids*, *Bull. London Math. Soc.* **5** (1973), 85–90.
- [19] Gregg Musiker and Tom Roby, *Paths to understanding birational rowmotion on products of two chains*, *Algebr. Comb.* **2** (2019), no. 2, 275–304.
- [20] Masatoshi Noumi and Yasuhiko Yamada, *Tropical Robinson-Schensted-Knuth correspondence and birational Weyl group actions*, in *Representation theory of algebraic groups and quantum groups*, *Adv. Stud. Pure Math.*, vol. 40, Math. Soc. Japan, Tokyo, 2004, pp. 371–442.
- [21] Neil O’Connell, Timo Seppäläinen, and Nikos Zygouras, *Geometric RSK correspondence, Whittaker functions and symmetrized random polymers*, *Invent. Math.* **197** (2014), no. 2, 361–416.
- [22] Igor Pak, *Hook length formula and geometric combinatorics*, *Sém. Lothar. Combin.* **46** (2001/02), article no. B46f (13 pages).
- [23] James Propp and Tom Roby, *Homomesy in products of two chains*, *Electron. J. Combin.* **22** (2015), no. 3, article no. 3.4 (29 pages).
- [24] Bruce E. Sagan, *The symmetric group. Representations, combinatorial algorithms, and symmetric functions*, second ed., *Graduate Texts in Mathematics*, vol. 203, Springer-Verlag, New York, 2001.
- [25] Richard P. Stanley, *Two poset polytopes*, *Discrete Comput. Geom.* **1** (1986), no. 1, 9–23.
- [26] Richard P. Stanley, *Promotion and evacuation*, *Electron. J. Combin.* **16** (2009), no. 2, Special volume in honor of Anders Björner, article no. 9 (24 pages).
- [27] Jessica Striker and Nathan Williams, *Promotion and rowmotion*, *European J. Combin.* **33** (2012), no. 8, 1919–1942.
- [28] H. Thomas and N. Williams, *Rowmotion in slow motion*, *Proc. Lond. Math. Soc. (3)* **119** (2019), no. 5, 1149–1178.

Rowmotion and the octahedron recurrence

JOSEPH JOHNSON, KTH Royal Institute of Technology, Department of Mathematics, Stockholm,
114 28 (Sweden)
E-mail : `josjohn@kth.se`

RICKY INI LIU, University of Washington, Department of Mathematics, Seattle, WA 98195 (USA)
E-mail : `riliu@uw.edu`