



ALGEBRAIC COMBINATORICS


Ayah Almousa, Anton Dochtermann & Ben Smith

Root polytopes, tropical types, and toric edge ideals

Volume 8, issue 1 (2025), p. 59-99.

<https://doi.org/10.5802/alco.404>

© The author(s), 2025.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*
www.tccpublishing.org www.centre-mersenne.org
e-ISSN: 2589-5486





Root polytopes, tropical types, and toric edge ideals

Ayah Almousa, Anton Dochtermann & Ben Smith

ABSTRACT We consider arrangements of tropical hyperplanes where the apices of the hyperplanes are taken to infinity in certain directions. Such an arrangement defines a decomposition of Euclidean space where a cell is determined by its ‘type’ data, analogous to the covectors of an oriented matroid. By work of Develin–Sturmfels and Fink–Rincón, these ‘tropical complexes’ are dual to regular subdivisions of root polytopes, which in turn are in bijection with mixed subdivisions of certain generalized permutohedra. Extending previous work with Joswig–Sanyal, we show how a natural monomial labeling of these complexes describes polynomial relations (syzygies) among ‘type ideals’ which arise naturally from the combinatorial data of the arrangement. In particular, we show that the cotype ideal is Alexander dual to a corresponding initial ideal of the lattice ideal of the underlying root polytope. This leads to novel ways of studying algebraic properties of various monomial and toric ideals, as well as relating them to combinatorial and geometric properties. In particular, our methods of studying the dimension of the tropical complex leads to new formulas for homological invariants of toric edge rings of bipartite graphs, which have been extensively studied in the commutative algebra community.

1. INTRODUCTION

The study of *tropical convexity* has its origins in optimization and related fields under the guise of max-plus linear algebra; see [10, 11, 34]. A geometric approach was initiated by Develin and Sturmfels in [17], where the notion of a *tropical polytope* was defined as the tropical convex hull of a finite set of points in the tropical torus $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. A tropical polytope can also be defined via an *arrangement of tropical hyperplanes* in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. Such an arrangement leads to a decomposition of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ called the *tropical complex*, and the subcomplex of bounded cells recovers the tropical polytope. In [17], it was shown that the combinatorial types of tropical complexes arising from n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ are in bijection with the set of regular subdivisions of a product of simplices $\Delta_{n-1} \times \Delta_{d-1}$. Via the *Cayley trick* these are in turn encoded by regular *mixed subdivisions* of the dilated simplex $n\Delta_{d-1}$.

An arrangement of tropical hyperplanes gives rise to combinatorial information captured by the *type* data of cells in the tropical complex, a tropical analogue of the

Manuscript received 7th March 2023, revised 28th March 2024 and 30th September 2024, accepted 7th October 2024.

KEYWORDS. tropical hyperplanes, free resolutions, toric edge ideals, root polytopes, monomial ideals.

ACKNOWLEDGEMENTS. The first author was partially supported by the NSF GRFP under Grant No. DGE-1650441 and by the RTG grant NSF/DMS-1745638.

The second author was partially supported by Simons Foundation Grant #964659.

The third author was supported by the Heilbronn Institute for Mathematical Research.

covector of an oriented matroid. In [18], the second author, Sanyal, and Joswig studied monomial ideals defined by fine and coarse type data and showed that various complexes arising from the tropical complex support minimal (co)cellular resolutions, leading to a number of combinatorial and algebraic applications. These results extend work of Block and Yu [9], who first exploited the connection between tropical polytopes and cellular resolutions. The ideas go further back to a paper of Novik, Postnikov, and Sturmfels [37] who studied ideals arising from (classical) oriented matroids. In [9] the authors also establish a connection between type ideals and Gröbner bases of initial ideals of toric ideals, based on results developed by Sturmfels in [48]. In particular, it is shown that the fine cotype ideal of a tropical hyperplane arrangement is Alexander dual to a certain initial ideal of the ideal of maximal minors of a $2 \times n$ matrix of indeterminates, where the term order is determined by the underlying arrangement.

1.1. GENERALIZATIONS TO ROOT POLYTOPES. A natural generalization of the product of simplices is given by the root polytopes introduced by Postnikov in [40]. For a finite graph G on vertex set $[n]$, the *root polytope* $Q_G \subset \mathbb{R}^n$ is the convex hull of $e_i - e_j$ for all edges $(i, j) \in E(G)$ with $i < j$. We will mostly be interested in root polytopes determined by *bipartite graphs* B . The root polytope Q_B associated to the complete bipartite graph $K_{d,n}$ is a product of simplices $\Delta_{d-1} \times \Delta_{n-1}$, and for general $B \subseteq K_{d,n}$ the root polytope Q_B is the convex hull of a subset of the vertices of $\Delta_{d-1} \times \Delta_{n-1}$. It follows that Q_B inherits unimodularity from the product of simplices.

A bipartite graph $B \subseteq K_{d,n}$ also gives rise to a polytope P_B obtained as a Minkowski sum of simplices, an example of a *generalized permutohedron* [40]. The Cayley embedding of the corresponding simplices naturally recovers the root polytope Q_B , and in particular the fine mixed subdivisions of P_B are in bijection with the triangulations of Q_B . For these (and other) reasons, the study of triangulations of Q_B is an active area of research, and for instance an axiomatic approach is introduced in [22]. In [21], Fink and Rincón show how the combinatorics of triangulations of root polytopes can be encoded in a generalized version of tropical hyperplane arrangements, where one is allowed infinite coordinates in the defining functionals. The finite entries in such an arrangement define a bipartite graph B , and in [21] it is shown how the combinatorics of the resulting tropical complex are described by the corresponding regular subdivision of the root polytope Q_B . This theory was also worked out from a more combinatorial perspective by Joswig and Loho in [32]. Once again the faces in the tropical complex are encoded by tropical covectors which give rise to fine/coarse types and associated monomial ideals.

1.2. OUR CONTRIBUTIONS. In this paper we apply methods from [18] to the more general setting of root polytopes and find surprising connections to other well-studied objects. We consider the fine and coarse (co)type ideals of a generalized tropical hyperplane arrangement and relate them to well-known algebraic objects, including determinantal ideals and toric edge ideals. Using this ‘tropical perspective’ we establish new results and provide novel and more compact proofs of several known theorems from the literature.

In what follows, we fix A to be a $d \times n$ matrix with entries in $\mathbb{R} \cup \{\infty\}$ (which we will hereby refer to as a *tropical matrix*), and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. The finite entries of A define a bipartite graph $B_A \subseteq K_{d,n}$. The values of these entries define a lifting of the vertices of the polytope Q_{B_A} , which in turn induces a regular subdivision $\mathcal{S}(A)$ of Q_{B_A} . We say that A is *sufficiently generic* if $\mathcal{S}(A)$ is a triangulation. The arrangement $\mathcal{H}(A)$ also defines a subdivision $\mathcal{C}(A)$ of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ called the *tropical complex*, and we

let $\mathcal{B}(A)$ denote the subcomplex consisting of the bounded cells of $\mathcal{C}(A)$. We refer to Example 1.6 and Figure 1 for illustrations of these constructions.

Our first result is a combinatorial formula for the dimension of $\mathcal{B}(A)$, which will be pivotal for our algebraic applications. For a bipartite graph B , denote by $\lambda(B)$ the *recession connectivity* of B , defined in terms of subgraphs of B that induce strongly connected directed graph structures (see Definition 3.5). Extending earlier work on tropical convexity, we establish the following.

THEOREM 1.1 (Proposition 3.7, Corollary 6.8). *Let $B \subset K_{d,n}$ be a connected bipartite graph. Then for any $d \times n$ tropical matrix A satisfying $B = B_A$ we have*

$$\dim(\mathcal{B}(A)) \leq \lambda(B_A) - 1.$$

Furthermore, if A is sufficiently generic then we have equality in the above expression.

1.2.1. Resolutions of type ideals. We next turn to algebraic applications. For a tropical hyperplane arrangement $\mathcal{H} = \mathcal{H}(A)$, the cells of the tropical complex $\mathcal{C}(A)$ (and hence of $\mathcal{B}(A)$) have a natural monomial labeling given by their fine/coarse type and cotype. These in turn define the fine/coarse type and cotype ideals; see Section 2. Our next collection of results describes how homological invariants of these ideals are encoded in the facial structure of the associated tropical complexes. In particular, by employing methods from [18], we establish the following.

THEOREM 1.2 (Propositions 4.3, 4.4, 4.5). *Let $\mathcal{H} = \mathcal{H}(A)$ be a sufficiently generic arrangement of tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. Then with the monomial labels given by fine type (respectively, coarse type), the complex $\mathcal{C}(A)$ supports a minimal cocellular resolution of the fine (resp. coarse) type ideal. Similarly, the labeled complex $\mathcal{B}(A)$ supports a minimal cellular resolution of the fine (resp. coarse) cotype ideal.*

Recall that the finite entries of the matrix A define a bipartite graph $B = B_A \subset K_{d,n}$. We let P_B denote the corresponding generalized permutohedron (sum of simplices), and note that the coarse type ideal is generated by monomials corresponding to the lattice points of P_B . An application of the Cayley trick therefore implies that P_B supports a minimal cellular resolution of the ideal generated by its lattice points; see Corollaries 4.6 and 4.7. We refer to Figure 2 for an illustration.

Our results involving cellular resolutions of type ideals allow us to interpret volumes of generalized permutohedra in terms of algebraic data. For instance, we compute the volume of P_B as well as other combinatorial data including *G-draconian sequences*, in terms of a computation in initial ideals (see Proposition 4.23). If $\mathcal{H} = \mathcal{H}(A)$ is a *graphic* tropical hyperplane arrangement, i.e., each column in the tropical matrix A has exactly two finite entries (see Definition 4.19), the associated polytope P_B is a graphical *zonotope*, and its volume is given by the dimension of the largest syzygy of the underlying type ideal (see Corollary 4.21).

The entries of the tropical matrix A naturally define a term order on the polynomial ring whose variables are given by the vertices of the associated root polytope Q_B . This in turn defines an initial ideal $\text{in}_A(L)$ of the toric ideal L defined by Q_B . Extending results of [9] and [18] we relate this ideal to the cotype ideal associated to the arrangement $\mathcal{H}(A)$.

THEOREM 1.3 (Proposition 4.14). *Let $\mathcal{H} = \mathcal{H}(A)$ be any arrangement of tropical hyperplanes, and let L be the lattice ideal of the associated root polytope Q_{B_A} . Then the fine cotype ideal associated to \mathcal{H} is the Alexander dual of $M(\text{in}_A(L))$, where $M(\text{in}_A(L))$ is the largest monomial ideal contained in $\text{in}_A(L)$.*

In the case that every entry of A is finite (so that each hyperplane in \mathcal{H} has full support) we have that $B = K_{d,n}$ is the complete bipartite graph. In this case we see

that Q_B is a product of simplices $\Delta_{d-1} \times \Delta_{n-1}$, $P_B = n\Delta_{d-1}$ is a dilated simplex, and the toric ideal L is generated by the 2-minors of an $d \times n$ matrix of indeterminates.

1.2.2. *Toric edge rings.* Our primary application of type ideals will be their connection to *toric edge rings*. Fix a field \mathbb{k} , and let G be a finite graph G on vertex set V and edge set E . The *edge ring* $\mathbb{k}[G]$ is the \mathbb{k} -algebra generated by the monomials $x_i x_j$ where $\{i, j\} \in E$. Define a monomial map $\pi : \mathbb{k}[E] \rightarrow \mathbb{k}[V]$ by $e_j \mapsto v_{j_1} v_{j_2}$, where $e_j = \{v_{j_1}, v_{j_2}\}$. The *toric edge ideal* of G is $J_G = \ker \pi$, so that

$$\mathbb{k}[G] \cong \mathbb{k}[E]/J_G.$$

Homological properties of toric edge rings have been intensively studied in recent years; see Section 5 for references. Such ideals are homogeneous and are known to be generated by binomials corresponding to even closed walks in G (see [51]). We apply our results regarding type ideals to provide new insights into homological properties of toric edge ideals of bipartite graphs. For instance, because type ideals always have linear resolutions, we give a short proof that toric edge rings of bipartite graphs are Cohen-Macaulay, see Proposition 6.5. Alexander duality also provides us with the following geometric interpretation of the regularity of toric edge rings.

THEOREM 1.4 (Theorem 6.6, Corollary 5.5). *Suppose $B \subset K_{d,n}$ is a connected bipartite graph. Let A be a sufficiently generic tropical matrix satisfying $B = B_A$, and let $\mathcal{B}(A)$ denote the bounded subcomplex of the arrangement $\mathcal{H}(A)$. Then the regularity and Krull dimension of the toric edge ring $\mathbb{k}[B]$ are given by*

$$(1) \quad \begin{aligned} \operatorname{reg}(\mathbb{k}[B]) &= \dim(\mathcal{B}(A)); \\ \dim(\mathbb{k}[B]) &= d + n - 1. \end{aligned}$$

This geometric perspective on the regularity of toric edge rings of bipartite graphs leads to a number of further results. For instance, in Theorem 6.11 we observe a monotonicity property for regularity, in the sense that if $B' \subset B$ is a connected subgraph then

$$\operatorname{reg}(\mathbb{k}[B']) \leq \operatorname{reg}(\mathbb{k}[B]).$$

This was proved for the case of *induced* subgraphs (in the not necessarily bipartite context) by Hà, Kara, and O’Keefe in [25].

We apply our combinatorial characterization of $\dim(\mathcal{B}(A))$ from Theorem 1.1 to obtain new bounds on the regularity of toric edge rings. For instance, in Theorem 6.13 we prove a bound on $\operatorname{reg}(\mathbb{k}[B])$ for all connected bipartite graphs, in terms of the bipartition of B . This extends a result of Biermann, O’Keefe, and Van Tuyl from [8], who established the bound for *chordal bipartite* graphs. Our Theorem 1.1 also immediately implies a result of Herzog and Hibi (see [26]), who bound $\operatorname{reg}(\mathbb{k}[B])$ by one less than the *matching number* of the graph B . It is clear that $\lambda(B) \leq \operatorname{mat}(B)$ and hence we obtain a strengthening of this result, see Corollary 6.14.

For our next result recall that $\mathbb{k}[B]$ has a *q-linear resolution* if the Betti numbers satisfy $\beta_{i,j} = 0$ for all i and all $j \neq q + i - 1$. Note that in this case we have $\operatorname{reg}(\mathbb{k}[B]) = q - 1$ and J_B is generated by homogeneous polynomials of degree q . Our techniques lead to the following.

THEOREM 1.5 (Theorem 6.19). *Suppose $B \subset K_{d,n}$ is a finite connected bipartite graph. Then we have*

- $\mathbb{k}[B]$ has a 2-linear resolution if and only if B is obtained by appending trees to the vertices of $K_{2,d}$.
- If $q \geq 3$ and $\mathbb{k}[B]$ has a q -linear resolution then $\mathbb{k}[B]$ is a hypersurface.

The first statement was established by Ohsugi and Hibi in [39, Theorem 4.6] and the second statement recovers a result of Tsuchiya from [50]. Our short proofs of these results demonstrates how tropical and discrete geometric methods can be brought to bear.

In recent years we have seen a number of mathematicians working on the combinatorics of subdivisions of root polytopes, and a mostly disjoint group studying the homological properties of toric edge ideals. As far as we know there has been no dialogue between these communities and it is our hope that the work presented here will lead to more fruitful connections and applications. We explore some potential projects in Section 7.

1.3. RUNNING EXAMPLE. We end this section with an extended example to illustrate some of the main results described above. These particular constructions will be used throughout the paper.

EXAMPLE 1.6. Consider the 3×4 tropical matrix

$$A = \begin{bmatrix} 0 & 3 & \infty & 0 \\ 0 & 0 & 2 & 3 \\ 0 & \infty & 0 & \infty \end{bmatrix}.$$

The matrix A defines the bipartite graph B_A , the arrangement $\mathcal{H}(A)$ of four tropical hyperplanes in $\mathbb{R}^3/\mathbb{R}\mathbf{1}$, as well as the bounded subcomplex $\mathcal{B}(A)$ depicted in Figure 1.




FIGURE 1. The bipartite graph B_A , hyperplane arrangement $\mathcal{H} = \mathcal{H}(A)$, and bounded complex $\mathcal{B}(A)$ associated to the tropical matrix A from Example 1.6. The points in $\mathbb{R}^3/\mathbb{R}\mathbf{1}$ are labelled by their representative with $p_1 = 0$.

One can check that the graph B_A has recession connectivity $\lambda(B_A) = 3$, and hence Theorem 1.1 tells us that $\dim(\mathcal{B}(A)) = 2$, as indicated in the figure. The natural labeling of cells in the decomposition of $\mathbb{R}^3/\mathbb{R}\mathbf{1}$ determined by $\mathcal{H}(A)$ defines the fine/coarse type ideals, as well as the fine/coarse cotype ideals. Theorem 1.2 says that the type ideals have cellular resolutions supported on the tropical complex $\mathcal{C}(A)$, whereas the cotype ideals have resolutions supported on the bounded complex $\mathcal{B}(A)$. See Figure 1 for an illustration of the resolution of the fine cotype ideal.

The bipartite graph B also defines a generalized permutohedron P_B , and if $B = B_A$ for some tropical matrix A we obtain a mixed subdivision of P_B that is dual to the arrangement $\mathcal{H}(A)$. A corollary to Theorem 1.2 (see Corollaries 4.6 and 4.7) says that this mixed subdivision supports a minimal cellular resolution of the ideal generated by the lattice points of P_B (which coincides with the coarse type ideal associated to A). See Figure 2 for an illustration.




FIGURE 2. The generalized permutohedron P_B along with the mixed subdivision dual to the hyperplane arrangement in Figure 1. The monomial ideal generated by the vertices is minimally supported by this complex.

Next recall that any graph B on edge set E defines a toric edge ring $\mathbb{k}[B]$ a toric edge ideal $J_B \subset \mathbb{k}[E]$. For the bipartite graph $B = B_A$ depicted in Figure 1 we have

$$\tilde{S} = \mathbb{k}[E] = \mathbb{k}[x_{11}, x_{12}, x_{14}, x_{21}, x_{22}, x_{23}, x_{24}, x_{31}, x_{33}],$$

and the corresponding toric edge ideal is given by

$$J_B = \langle x_{11}x_{22} - \underline{x_{12}x_{21}}, \underline{x_{11}x_{24}} - x_{14}x_{21}, \underline{x_{12}x_{24}} - x_{14}x_{22}, x_{21}x_{33} - \underline{x_{23}x_{31}} \rangle.$$

Note that the binomial generators are describe by the minimal closed even walks in B .

Our observations imply that the ideal J_B corresponds to the lattice ideal of the root polytope Q_B . Furthermore, the entries of the tropical matrix A provides a monomial term order on \tilde{S} , which in turn defines an initial ideal $(\text{in}_A(J_B))$. In our example $(\text{in}_A(J_B))$ is generated by the underlined monomials above. The Alexander dual of this ideal (in the polynomial ring \tilde{S}) is given by

$$(\text{in}_A(J_B))^\vee = \langle x_{12}x_{23}x_{24}, x_{11}x_{12}x_{23}, x_{11}x_{31}x_{12}, x_{31}x_{12}x_{24}, x_{21}x_{31}x_{24}, x_{21}x_{23}x_{24} \rangle.$$

Our Theorem 1.3 then says that $(\text{in}_A(J_B))^\vee$ coincides with the fine cotype ideal associated to the arrangement $\mathcal{H} = \mathcal{H}(A)$, which has a minimal resolution supported on the bounded complex \mathcal{B} (again see Figure 1). Also by construction the arrangement $\mathcal{H}(A)$ depicted in Figure 1 satisfies $B = B_A$, and from Theorem 1.4 we conclude that

$$\text{reg}(\mathbb{k}[B]) = \dim(\mathcal{B}(A)) = 2.$$

Note that in this case J_B is generated by binomials of degree 2 but $\text{reg}(\mathbb{k}[B]) = 2$. Hence this ideal does not have a linear resolution, which can also be deduced from our Theorem 1.5.

1.4. ORGANIZATION AND NOTATION. The rest of the paper is organized as follows. In Section 2, we establish fundamental notation and terminology for the remainder of the paper surrounding tropical hyperplane arrangements with not necessarily full support, root polytopes, and generalized permutohedra. In Section 3, we study the topology of the bounded complex obtained from such a tropical hyperplane arrangement. We also define the *recession graph* (Definition 3.5) of a bipartite graph, whose connectivity properties can be used to detect bounded cells (Proposition 3.2) and bound the dimension of the bounded complex (Proposition 3.7).

In Section 4, we introduce the fine and coarse *type* and *cotype* ideals associated to an arrangement (Definition 4.2) and show that the bounded complex supports a minimal cellular resolution of cotype ideals (Proposition 4.5), and that for sufficiently generic arrangements, the tropical complex supports a minimal, linear, cocellular resolution

of type ideals (Proposition 4.4). These results imply that the fine mixed subdivision of the generalized permutohedron associated to the arrangement supports a cellular resolution of the type ideal (Corollary 4.6), which we apply to bound the (homological) types of type ideals in the case of graphic arrangements (Corollary 4.21) and give a novel way of computing volumes of generalized permutohedra (Proposition 4.23).

In Section 5, we establish that the toric edge ideal of a bipartite graph B is exactly the toric lattice ideal of the associated root polytope Q_B , and observe conditions on the root polytope such that its lattice ideal corresponds to well-studied ideals in commutative algebra such as ladder determinantal ideals and Ferrers ideals. This crucial observation drives the results of Section 6, where we relate combinatorial and topological properties of the bounded complex associated to an arrangement with homological properties of the associated toric edge ideal. In particular, our methods recover and strengthen several results in the literature bounding regularity of toric edge ideals of bipartite graphs; see Theorem 5.7 and Corollary 6.14. In addition, we give a novel proof characterizing toric edge ideals of bipartite graphs with linear resolution (Theorem 6.19). Finally, we conclude the paper with Section 7, where we collect questions and potential directions for further exploration.

1.5. NOTATION. Throughout the paper we make use of several mathematical objects that may not be familiar to the reader. For convenience we collect notations here.

- $\mathbb{T} = (\mathbb{R} \cup \{-\infty\}, \oplus, \odot)$ is the *(max)-tropical semiring* (Definition 2.1);
- $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ is the tropical torus;
- A is a $d \times n$ matrix with entries in $\mathbb{R} \cup \{\infty\}$ (a *tropical matrix*);
- $\mathcal{H}(A)$ is the tropical hyperplane arrangement associated to A (Definition 2.2);
- $\mathcal{C}(A)$ is the tropical complex associated to $\mathcal{H}(A)$, $\mathcal{B}(A)$ is the bounded sub-complex (Definition 2.12);
- $I_{\mathcal{T}(\mathcal{H})}$ and $I_{\overline{\mathcal{T}}(\mathcal{H})}$ are the fine type and cotype ideal, and $I_{\mathbf{t}(\mathcal{H})}$ and $I_{\overline{\mathbf{t}}(\mathcal{H})}$ are the coarse type and cotype ideal, associated to \mathcal{H} (Definition 4.2);
- $B \subseteq K_{d,n}$ is a bipartite graph, B_A is the bipartite graph associated to A ;
- Q_B is the root polytope associated to B (Definition 2.17);
- P_B is the generalized permutohedron (sum of simplices) associated to B (Definition 2.19);
- $\mathbb{k}[B]$ is the toric edge ring and J_B is the toric edge ideal associated to B with underlying field \mathbb{k} (Definition 5.2);
- $R(S; B)$ is the recession graph of S inside B (Definition 3.1);
- $\lambda(B)$ is the *recession connectivity* of B (Definition 3.5);
- $\text{LD}(B)$ is the *left-degree vector* of B and $\text{RD}(B)$ is its *right-degree vector* (see Definition 2.5).

2. TROPICAL HYPERPLANES, ROOT POLYTOPES, AND GENERALIZED PERMUTOHEDRA

In this section, we review relevant notions from tropical convexity and establish notation that will be used in the rest of the paper. We discuss tropical hyperplane arrangements and corresponding tropical complexes, and how their combinatorial data is encoded in both fine and coarse types and cotypes. We then discuss the connection to regular subdivisions of root polytopes and mixed subdivisions of generalized permutohedra. For proofs and further reading, we refer to [31].

2.1. TROPICAL HYPERPLANE ARRANGEMENTS.

DEFINITION 2.1 (Tropical Semiring). Let $\mathbb{T} = (\mathbb{R} \cup \{-\infty\}, \oplus, \odot)$ be the (max)-tropical semiring where

$$x \oplus y := \max(x, y) \quad \text{and} \quad x \odot y := x + y.$$

One can also define the (min)-tropical semiring by replacing $-\infty$ with ∞ , and the operation \max with \min ; the two semi-rings are isomorphic via

$$-\max(x, y) = \min(-x, -y).$$

Unless stated otherwise, we shall work with the max-tropical semiring, and note that all statements can be rephrased in terms of the min-tropical semiring. If we need to work with both simultaneously, we shall differentiate between them by denoting them respectively by \mathbb{T}_{\min} and \mathbb{T}_{\max} .

We can extend the operations on \mathbb{T} to addition and scalar multiplication on \mathbb{T}^d , giving it the structure of a semimodule:

$$\mathbf{x} \oplus \mathbf{y} = (x_1 \oplus y_1, \dots, x_d \oplus y_d), \quad c \odot \mathbf{x} = (c \odot x_1, \dots, c \odot x_d), \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{T}^d, c \in \mathbb{T}.$$

Restricting \mathbb{T} to \mathbb{R} gives a similar semimodule structure on \mathbb{R}^d . This will be convenient when considering tropical convexity in Section 3.2.

Following [21], we generalize the notion of a tropical hyperplane arrangement by allowing some of the hyperplanes to be “taken to infinity” in some directions.

DEFINITION 2.2 (Tropical Hyperplane Arrangement). Let $\mathbf{a} \in (\mathbb{R} \cup \{\infty\})^d$ with at least one finite entry. The (max)-tropical hyperplane H with apex \mathbf{a} is the set

$$H(\mathbf{a}) := \{ \mathbf{p} \in \mathbb{R}^d \mid (-a_1 \odot p_1) \oplus (-a_2 \odot p_2) \oplus \dots \oplus (-a_d \odot p_d) \text{ is attained at least twice} \},$$

that is, it is the vanishing locus of a max-tropical homogeneous linear polynomial whose coefficients are $-\mathbf{a}$.

A $d \times n$ matrix $A = (\mathbf{a}_1, \dots, \mathbf{a}_n)$ with entries $\in \mathbb{R} \cup \{\infty\}$ such that each \mathbf{a}_i has at least one finite entry will be called a tropical matrix. Such a matrix gives rise to an (ordered) (max)-tropical hyperplane arrangement $\mathcal{H}(A) = (H_1, \dots, H_n)$ in \mathbb{R}^d whose i th hyperplane $H_i := H(\mathbf{a}_i)$ is the tropical hyperplane with apex \mathbf{a}_i .

An important observation is that if $H(\mathbf{a})$ is a max-tropical hyperplane, its apex \mathbf{a} has entries in \mathbb{T}_{\min} , and vice-versa. Unless stated otherwise, we shall work with max-tropical hyperplanes, so that their apices will be elements of \mathbb{T}_{\min}^d . Although we allow tropical hyperplanes with infinite apices, we will only consider their vanishing locus in \mathbb{R}^d .

Observe that for any point $\mathbf{p} \in H(\mathbf{a})$, the point $\mathbf{p} + \lambda \cdot \mathbf{1}$ is also contained in $H(\mathbf{a})$, where $\mathbf{1} \in \mathbb{R}^d$ is the vector of all ones. For this reason, one typically views tropical objects in the tropical torus $\mathbb{R}^d / \mathbb{R}\mathbf{1}$, by definition the quotient of the Euclidean space \mathbb{R}^d by the linear subspace $\mathbb{R}\mathbf{1}$. By interpreting this quotient in the category of topological spaces, $\mathbb{R}^d / \mathbb{R}\mathbf{1}$ inherits a natural topology which is homeomorphic to the usual topology on \mathbb{R}^{d-1} . While $\mathbb{R}^d / \mathbb{R}\mathbf{1}$ is a convenient space to view geometric objects, we must define said objects in \mathbb{R}^d first, as the addition operation of the tropical semiring is not well defined in $\mathbb{R}^d / \mathbb{R}\mathbf{1}$.

DEFINITION 2.3 (Support of a Hyperplane). The support of a tropical hyperplane $H(\mathbf{a})$ is $\text{supp}(H) = \{k \in [d] \mid a_k \text{ finite}\}$. The support of a hyperplane arrangement $\mathcal{H}(A) = (H_1, \dots, H_n)$ is the set

$$\text{supp}(A) = \{(i, j) \in [d] \times [n] \mid i \text{ is in the support of } H_j\},$$

so that $\text{supp}(\mathcal{H}(A))$ is equal to the support of the matrix A .

The combinatorics of supports can be described by a bipartite graph in the following way.

DEFINITION 2.4 (Bipartite graphs from tropical matrices). *Suppose A is a $d \times n$ tropical matrix and let $\mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. Define $B_A \subseteq K_{d,n}$ to be the bipartite graph whose edges encode the support of A , i.e.,*

$$(i, j) \in E(B_A) \iff (i, j) \in \text{supp}(A).$$

We refer to Figure 1 for an illustration of $\mathcal{H}(A)$ and B_A described in Definitions 2.2 and 2.4. One can encode information about a tropical hyperplane arrangement $\mathcal{H}(A)$ via left and right degree vectors of the associated bipartite graph B_A .

DEFINITION 2.5 (Left/right degree vectors). *For a bipartite graph $B \subset K_{d,n}$, the left degree vector $\text{LD}(B) = (l_1, \dots, l_d)$ is the tuple of node degrees of vertices $1, \dots, d$, and the right degree vector $\text{RD}(B) = (r_1, \dots, r_n)$ is the tuple of node degrees of vertices $1, \dots, n$.*

EXAMPLE 2.6. We continue with our running example

$$A = \begin{bmatrix} 0 & 3 & \infty & 0 \\ 0 & 0 & 2 & 3 \\ 0 & \infty & 0 & \infty \end{bmatrix}.$$

The corresponding bipartite graph B_A and tropical hyperplane arrangement $\mathcal{H}(A)$ is depicted in Figure 1. The tropical hyperplane $H(\mathbf{a}_1)$ has full support $\{1, 2, 3\}$, hyperplanes $H(\mathbf{a}_2)$ and $H(\mathbf{a}_4)$ have support set $\{1, 2\}$, and hyperplane $H(\mathbf{a}_3)$ has support set $\{2, 3\}$. This support data can be combined to give the corresponding bipartite graph B_A (see Figure 1), with left degree vector $\text{LD}(B_A) = (3, 4, 2)$ and right degree vector $\text{RD}(B_A) = (3, 2, 2, 2)$.

2.2. TYPES AND COTYPES. By definition, a point $\mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1}$ is contained in $H(\mathbf{a})$ if and only if $\max_{j \in [d]}(p_j - a_j)$ is attained at least twice. The points that obtain this maximum value in one particular term form a polyhedral *sector*, and a tropical hyperplane decomposes $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ into these sectors.

DEFINITION 2.7 (Sector). *The i -th (closed) sector of the (max)-tropical hyperplane $H(\mathbf{a})$ is*

$$S_i(\mathbf{a}) = \{\mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \mid p_i - a_i \geq p_j - a_j \text{ for all } j \in [d]\}.$$

The definition of a min-tropical sector is identical except with the inequality reversed, as the corresponding tropical linear functional is looking to minimize.

Given an arrangement \mathcal{H} of tropical hyperplanes, to each point in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ we can record which sector it sits in with respect to each hyperplane. This leads to the notion of *types* and *cotypes*.

DEFINITION 2.8 (Fine type/cotype). *Let $A = (\mathbf{a}_1, \dots, \mathbf{a}_n)$ be a $d \times n$ tropical matrix, and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. The fine type of a point $\mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1}$ with respect to \mathcal{H} is the table $T_{\mathcal{H}}(\mathbf{p}) \in \{0, 1\}^{d \times n}$ with*

$$T_{\mathcal{H}}(\mathbf{p})_{ik} = 1 \text{ if and only if } \mathbf{p} \in S_i(\mathbf{a}_k)$$

for $(i, k) \in \text{supp}(\mathcal{H})$. The fine cotype $\bar{T}_{\mathcal{H}}(\mathbf{p}) \in \{0, 1\}^{d \times n}$ is defined by $\bar{T}_{\mathcal{H}}(\mathbf{p})_{ik} = 1 - T_{\mathcal{H}}(\mathbf{p})_{ik}$ for $(i, k) \in \text{supp}(\mathcal{H})$, and zero for $(i, k) \notin \text{supp}(\mathcal{H})$.

DEFINITION 2.9 (Coarse type/cotype). *Let $\mathcal{H} = \mathcal{H}(A)$ be an arrangement of n max-tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. The coarse type of a point $\mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1}$ with respect to \mathcal{H} is given by $\mathbf{t}_{\mathcal{H}}(\mathbf{p}) = (t_1, t_2, \dots, t_d) \in \mathbb{N}^d$ with*

$$t_i = \sum_{k=1}^n T_{\mathcal{H}}(\mathbf{p})_{ik}$$

for $i \in [d]$.

NOTATION 2.10. We will often view fine (co)types as (complements of) bipartite subgraphs of the support graph B_A described in Definition 2.4. Explicitly, a fine type $T \in \{0, 1\}^{d \times n}$ encodes a subgraph $B_T \subset B_A$ whose edges are (i, k) whenever T has a 1 in the (i, k) -th entry. Similarly, a fine cotype \bar{T} encodes a subgraph $B_{\bar{T}} \subset B_A$ given by the complement of B_T inside B_A . As a bipartite graph, the corresponding coarse (co)type is the left degree vector $\text{LD}(B_T)$. As they are equivalent, we will only explicitly use B_T when graph properties are required.

Observe that not all tables or bipartite graphs can arise as fine types, as every point in $\mathbb{R}^d/\mathbb{R}\mathbb{1}$ is contained in at least one sector of a tropical hyperplane. Each column of a viable table must have at least one non-zero entry, and viable bipartite graphs must have no disconnected nodes in $[n]$. These tables and graphs are called *type tables* and *type graphs*.

REMARK 2.11. Tropical types are also referred to as *tropical covectors* in the literature (see e.g. [21]). As we will work with both types and cotypes, we choose to follow the terminology in [18].

As discussed in [17] and [32], the type data of points $\mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbb{1}$ gives rise to a polyhedral complex associated to a tropical hyperplane arrangement as follows.

DEFINITION 2.12 (Tropical complex, bounded subcomplex). *Let A be a $d \times n$ tropical matrix and let $\mathcal{H}(A)$ denote the corresponding arrangement in $\mathbb{R}^d/\mathbb{R}\mathbb{1}$. For a fixed fine type $T = T_{\mathcal{H}(\mathbf{p})}$, the set*

$$C_T^\circ := \{\mathbf{q} \in \mathbb{R}^d/\mathbb{R}\mathbb{1} \mid T(\mathbf{q}) = T\}$$

is a relatively open subset of $\mathbb{R}^d/\mathbb{R}\mathbb{1} = \mathbb{R}^{d-1}$ called the relatively open cell of type T . The set of all relatively open cells $C^\circ = C^\circ(A)$ partitions the tropical torus $\mathbb{R}^d/\mathbb{R}\mathbb{1}$.

Denote by C_T the closed cell of type T , the set of all points $\mathbf{q} \in \mathbb{R}^d/\mathbb{R}\mathbb{1}$ such that $T(\mathbf{q}) \supseteq T$. The collection of all closed cells yields a polyhedral subdivision $\mathcal{C} = \mathcal{C}(A)$ of $\mathbb{R}^d/\mathbb{R}\mathbb{1}$ called the tropical complex. The cells which are bounded form the bounded subcomplex $\mathcal{B} = \mathcal{B}(A)$.

EXAMPLE 2.13. Consider the running example discussed in Example 2.6. Figure 3 shows the tropical complex and bounded complex, along with certain distinguished cells whose types are given in Figure 4.

The combinatorics of tropical types of an arrangement $\mathcal{H} = \mathcal{H}(A)$ reflect the geometry of the tropical complex $\mathcal{C}(A)$. In the following we consider fine types as bipartite graphs, and use $c(G)$ to denote the number of connected components of a graph G .

LEMMA 2.14 ([17], [32]). *Suppose A is a $d \times n$ tropical matrix with corresponding arrangement $\mathcal{H}(A)$, and let $\mathcal{C}(A)$ be the tropical complex. Let \mathcal{T} be the collection of fine types that appear. For all $S, T \in \mathcal{T}$, we have:*

- (1) $C_S \cap C_T = C_{S \cup T}$ (where $B_{S \cup T} = B_S \cup B_T$),
- (2) $C_S \subseteq C_T \iff B_S \supseteq B_T$,
- (3) $\dim(C_T) = c(B_T) - 1$.

Proof. See Remark 28 and Corollary 38 of [32]. □

Tropical hyperplanes arrangements enjoy a natural notion of duality that we will need in our work. Explicitly, an arrangement $\mathcal{H}(A)$ of n hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbb{1}$ is “dual” to $\mathcal{H}(A^T)$, the arrangement of d hyperplanes in $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ obtained by taking the transpose of the tropical matrix. Moreover, despite being in different ambient spaces, these arrangements have isomorphic bounded complexes; this follows from [21, Proposition 6.2].




FIGURE 3. The tropical complex and bounded complex corresponding to the tropical hyperplane arrangement in Figure 1.




FIGURE 4. The tropical types corresponding to the tropical complex and distinguished cells in Figure 3.

PROPOSITION 2.15. *Let A be a $d \times n$ tropical matrix. There exists a piecewise linear bijection between the bounded complexes $\mathcal{B}(A)$ and $\mathcal{B}(A^T)$ that induces an isomorphism between their face posets.*

EXAMPLE 2.16. Consider the tropical matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 0 & 2 \end{bmatrix}.$$

The tropical hyperplane arrangements $\mathcal{H}(A) \subseteq \mathbb{R}^3/\mathbb{R}\mathbf{1}$ and $\mathcal{H}(A^T) \subseteq \mathbb{R}^2/\mathbb{R}\mathbf{1}$ are given in Figure 5. In particular, both bounded complexes consist of two edges and three vertices.

2.3. SUBDIVISIONS OF ROOT POLYTOPES AND GENERALIZED PERMUTOHEDRA. As discussed in [21], an arrangement $\mathcal{H} = \mathcal{H}(A)$ of n hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ defined by a $d \times n$ tropical matrix A gives rise to a regular subdivision of a root polytope associated to A . An application of the Cayley trick then provides a regular mixed subdivision of a certain generalized permutohedron. For the case of hyperplanes with full support, this recovers the connection to subdivisions of products of simplices and mixed divisions of dilated simplices, as worked out in [17]. We first recall some definitions.

DEFINITION 2.17 (Root Polytopes). *Let $B \subseteq K_{d,n}$ be a bipartite graph and denote by $E(B)$ its edge set. Define Q_B to be the root polytope*

$$Q_B = \text{conv}\{\mathbf{e}_i - \mathbf{e}_j \mid (i, j) \in E(B)\} \subset \mathbb{R}^{d+n},$$

where $\mathbf{e}_1, \dots, \mathbf{e}_d, \mathbf{e}_{\bar{1}}, \dots, \mathbf{e}_{\bar{n}}$ are the coordinate vectors in \mathbb{R}^{d+n} .




FIGURE 5. The tropical hyperplane arrangements $\mathcal{H}(A)$ and $\mathcal{H}(A^T)$. Their bounded complexes, shown in bold, are isomorphic.

EXAMPLE 2.18. A well-studied class of root polytopes arises when $B = K_{d,n}$ is a complete bipartite graph. In this case, the root polytope $Q_{K_{d,n}}$ is the product of simplices $\Delta_{d-1} \times \Delta_{n-1}$. For other bipartite graphs, Q_B is the convex hull of some subset of vertices of $\Delta_{d-1} \times \Delta_{n-1}$.

The root polytope Q_B of a bipartite graph $B \subset K_{d,n}$ is also closely related to another convex body obtained as a sum of (not necessarily full dimensional) simplices in \mathbb{R}^d . Such polytopes are examples of the *generalized permutohedra* studied by Postnikov in [40].

DEFINITION 2.19 (Generalized permutohedra). *Let $B \subseteq K_{d,n}$ be a bipartite graph and let $N_j(B) \subseteq [d]$ denote the set of neighbours of $j \in [n]$. The generalized permutohedron associated to B is the polytope*

$$P_B = \sum_{j=1}^n \Delta_{N_j(B)} \subset \mathbb{R}^d,$$

where $\Delta_{N_j(B)}$ denotes the subsimplex of Δ_{d-1} whose vertices are labelled by $N_j(B)$.

DEFINITION 2.20 (Mixed subdivision). *Suppose $P_B = \sum_{j=1}^n \Delta_{N_j(B)} \subset \mathbb{R}^d$ is a generalized permutohedron of dimension k . A Minkowski cell in this sum is a polytope of dimension k of the form $\Delta_{I_1} + \dots + \Delta_{I_n}$, where $I_j \subset N_j(B)$. A mixed subdivision \mathcal{M} of P_B is a decomposition into a union of Minkowski cells, with the property that the intersection of any two cells is their common face.*

REMARK 2.21. Our definition of a mixed subdivision of a sum of simplices is a special case of the corresponding concept for any Minkowski sum of polytopes. In the more general case, stricter conditions must be satisfied that are always satisfied for simplices; see [44] for further details.

The *Cayley trick* [45] provides a geometric correspondence between the generalized permutohedron P_B and the root polytope Q_B . In particular we recover P_B by intersecting Q_B with the affine subspace $\mathbb{R}^d \times \{-\sum_{j=1}^n \mathbf{e}_j\}$. Under this intersection, polyhedral subdivisions of Q_B are in one-to-one correspondence with mixed subdivisions of P_B [30]. In our context a $d \times n$ tropical matrix A induces a *regular subdivision* $\mathcal{S}(A)$ of the root polytope Q_{B_A} by lifting the vertex $\mathbf{e}_i - \mathbf{e}_j$ to height v_{ij} and projecting the lower faces of the resulting polytope back to \mathbb{R}^{d+n} . Via the Cayley trick,

this gives rise to a corresponding regular mixed subdivision $\mathcal{M}(A)$ of the generalized permutohedron $P_{B_A} = \sum_{j=1}^n \Delta_{N_j(B_A)} \subseteq \mathbb{R}^d$.

NOTATION 2.22. We say that a $d \times n$ tropical matrix A is *sufficiently generic* if the corresponding subdivision $\mathcal{S}(A)$ of the root polytope Q_{B_A} is a triangulation.

In [17] it was shown that the tropical complex associated to an arrangement of tropical hyperplanes (with full support) is dual to the corresponding mixed subdivision of a dilated simplex. Furthermore, this duality is described by fine type data and descriptions as mixed cells. This was generalized to the setting of arbitrary tropical matrices by Fink and Rincon as follows.

PROPOSITION 2.23. [21, Proposition 4.1] *Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of tropical hyperplanes. Then the tropical complex $\mathcal{C}(A)$ is dual to the mixed subdivision $\mathcal{M}(A)$ of P_{B_A} . A face of $\mathcal{C}(A)$ labelled by fine type $T \subseteq B_A$ is dual to the cell $\mathcal{M}(A)$ obtained as the sum $\sum_{j=1}^n \Delta_{N_j(T)}$.*

We refer to Figure 6 for an illustration of this duality in the context of our running example.




FIGURE 6. The tropical hyperplane arrangement $\mathcal{H}(A)$, where A is from Example 1.6, along with the corresponding mixed subdivision of a generalized permutohedron.

3. THE BOUNDED COMPLEX AND TOPOLOGY OF TYPES

In this section, we extend existing techniques in tropical convexity to derive new results regarding the bounded complex and the topology of type data associated to tropical hyperplane arrangements. These results will be interpreted in the algebraic setting later in the paper.

3.1. THE RECESSION CONE AND DIMENSION OF THE BOUNDED COMPLEX. For the case of a tropical matrix A with only finite entries, the bounded complex $\mathcal{B}(A)$ can be recovered as the *min*-tropical polytope $\text{tconv}(A)$, i.e. the smallest tropically convex set containing the columns of A ; see Definition 3.10. In this case there is a very simple combinatorial characterization for when a cell is contained in $\text{tconv}(A)$: the cell C_T is contained in $\mathcal{B}(A)$ if and only if its type T has no isolated left nodes [17].

For a general tropical matrix A , this characterization of $\text{tconv}(A)$ still holds (see [32]), but if A has infinite entries then the corresponding tropical polytope will have unbounded cells. In particular, the bounded complex $\mathcal{B}(A)$ will be a subcomplex of $\text{tconv}(A)$, and so T having no isolated left nodes becomes a necessary but not




FIGURE 7. A tropical complex with two distinguished cells, $C_T, C_{T'}$, along with their corresponding recession graphs. Thin edges are directed only from $[d]$ to $[n]$, while thick edges are bidirected.

sufficient condition for C_T being contained in $\mathcal{B}(A)$. In this section we give a precise combinatorial condition on T that characterizes when C_T is contained in $\mathcal{B}(A)$ for a general tropical matrix A .

DEFINITION 3.1. Let $B \subset K_{d,n}$ be a finite simple bipartite graph and $S \subseteq B$ a subgraph. We define the recession graph $R = R(S; B)$ to be the directed bipartite graph with vertices $V(B) = [d] \sqcup [n]$ and with edges $e \in E(B) \setminus E(S)$ directed from $[d]$ to $[n]$, and $e' \in E(S)$ bidirected, i.e. directed from $[d]$ to $[n]$ and $[n]$ to $[d]$.

If $B = B_A$ for a tropical matrix A and $S = B_T$ for some fine type T , we will also write $R(T; A)$ for the recession graph.

See Example 3.3 and Figure 7 for an example of recession graphs arising from a tropical hyperplane arrangement.

The recession graph is so named as it describes the *recession cone* of a cell. Recall that the recession cone of a polyhedron P is the cone

$$\text{rec}(P) = \{y \in \mathbb{R}^d \mid x + \lambda y \in P \forall x \in P, \lambda \geq 0\},$$

which describes the directions in which P is unbounded. The following proposition shows the relation between the recession cone and the recession graph. Recall that a digraph is *strongly connected* if one can find a directed path from any vertex to any other vertex.

PROPOSITION 3.2. [32] Let A be a tropical matrix, and suppose C_T is a cell in the tropical complex $\mathcal{C}(A)$. Then the dimension of the recession cone of C_T in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ is equal to the number of strongly connected components of $R(T; A)$, minus one.

In particular, the cell C_T is contained in the bounded complex $\mathcal{B}(A)$ if and only if the graph $R(T; A)$ is strongly connected.

Proof. This can be deduced from a number of results from [32] via weighted digraph polyhedra, all results cited within this proof are contained there. By Lemma 7 and Proposition 30, the cell C_T is affinely isomorphic to a weighted digraph polyhedron. In particular, the recession cone of C_T is the weighted digraph polyhedron $Q(R(T; A))$. By Proposition 9, the dimension of Q in $\mathbb{R}^{d+n}/\mathbb{R}\mathbf{1}$ is equal to the number of strongly connected components of $R(T; A)$ minus one, which extends to C_T in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ by affine isomorphism. Therefore C_T is bounded if and only if $R(T; A)$ is strongly connected. \square

EXAMPLE 3.3. Consider the tropical complex depicted in Figure 7, along with the recession graphs for the cells C_T and $C_{T'}$. The cell C_T is bounded, as reflected by the fact that the recession graph $R(T; A)$ is strongly connected. The cell $C_{T'}$ is unbounded, since in the recession graph $R(T'; A)$, the edge $(2, 3)$ can only be traversed from left to right. Hence there is no path from left node 3 to any other left node that respects edge orientation.

We get the following as an immediate corollary.

COROLLARY 3.4. *The bounded complex $\mathcal{B}(A)$ is non-empty if and only if B_A is connected.*

Proof. The bounded complex is non-empty if and only if it contains a 0-cell, as any bounded k -cell must contain a 0-cell. By Lemma 2.14, a cell C_T is zero dimensional if and only if $B_T \subseteq B_A$ is connected. This immediately shows $\mathcal{B}(A)$ non-empty implies B_A is connected.

Conversely if B_A is connected, the root polytope Q_{B_A} is full-dimensional [40, Lemma 12.5] and the maximal cells of its regular subdivision $\mathcal{S}(A)$ correspond to 0-cells of $\mathcal{C}(A)$ by Proposition 2.23. As $\mathcal{C}(A)$ contains a 0-cell, the bounded complex must be non-empty. \square

Inspired by these observations we define the following invariant of a bipartite graph. Recall that we use $c(B)$ to denote the number of connected components of a graph B .

DEFINITION 3.5. *For a bipartite graph B define the recession connectivity of B to be the integer*

$$\lambda(B) = \max_{S \subseteq B} \{c(S) \mid R(S; B) \text{ is strongly connected}\}.$$

REMARK 3.6. Note that for any bipartite graph B we have

$$\lambda(B) \leq \text{mat}(B),$$

where $\text{mat}(B)$ denotes the *matching number* of B . Indeed, any $S \subseteq B$ realizing $\lambda(B)$ contains a matching of size $c(S)$.

PROPOSITION 3.7. *Let B be a finite simple connected bipartite graph. For any tropical matrix A satisfying $B = B_A$, we have $\dim(\mathcal{B}(A)) \leq \lambda(B_A) - 1$. Furthermore, there exists a generic tropical matrix A with $\dim(\mathcal{B}(A)) = \lambda(B_A) - 1$.*

Proof. The first part of this statement is an immediate corollary of Proposition 3.2, as the dimension of any bounded cell is bounded above by $\lambda(B_A) - 1$. For the second part, let $S \subseteq B$ be a subgraph realizing $\lambda(B)$, i.e. $R(S; B)$ is strongly connected, and has $\lambda(B)$ connected components. We can assume that S is a forest, as removing an arbitrary edge from any cycle in S does not strongly disconnect $R(S; B)$ nor does it affect the number of connected components of S .

It remains to show that we can find a tropical matrix A and fine type T such that $B = B_A$ and $S = B_T$. We define the matrix A by setting $a_{ij} = 0$ for all $(i, j) \in S$, sufficiently large generic values for $(i, j) \in B \setminus S$, and ∞ otherwise. Then $B = B_A$, and the regular subdivision of Q_B induced by A is a triangulation that contains the cell Q_S . After applying the Cayley trick, Q_S corresponds to a mixed cell dual to the cell C_T in the tropical complex. By Proposition 3.2, C_T is contained in the bounded complex, and its dimension is $\lambda(B) - 1$. Furthermore, as $\lambda(B) - 1$ is the maximum possible dimension of a bounded cell, this shows $\dim(\mathcal{B}(A)) = \lambda(B) - 1$. \square

LEMMA 3.8. *For the complete bipartite graph $K_{d,n}$ we have $\lambda(K_{d,n}) = \min\{d, n\}$.*

Proof. Without loss of generality assume that $d \leq n$ and denote the vertices of $B = K_{d,n}$ as $\{v_1, \dots, v_d\} \cup \{w_1, \dots, w_n\}$. It is clear that $\lambda(K_{d,n}) \leq d$. For the reverse inequality, define S to be the graph with edges $v_1w_1, \dots, v_dw_d, v_dw_{d+1}, \dots, v_dw_n$. Then S has d connected components and $R(S; B)$ is strongly connected. The result follows. \square

The following lemma shows that λ is monotone on subgraphs, and will be useful at numerous points.

LEMMA 3.9. *Let B be a finite simple connected bipartite graph and $B' \subseteq B$ a connected subgraph. Then $\lambda(B') \leq \lambda(B)$.*

Proof. Let $S' \subseteq B'$ be a graph realizing $\lambda(B')$ i.e. $c(S') = \lambda(B')$ and $R(S'; B')$ is strongly connected. Note that we can assume S' is a forest: removing an edge from any cycle in S' does not disconnect $R(S'; B')$ as we can traverse the cycle in the opposite direction. Let $T_{B'}$ be a spanning tree of B' that contains S' . We then extend this to T_B , a spanning tree of B that contains $T_{B'}$. We define $S \subseteq B$ as the graph

$$S = S' \cup (T_B \setminus T_{B'}) .$$

We claim that $R(S; B)$ is strongly connected and that $c(S) \geq c(S')$.

We first show strong connectivity. Consider $v, v' \in V(R(S; B))$. If $v, v' \in V(B')$, the same path in $R(S'; B')$ connects them in $R(S; B)$. If $v, v' \in V(B \setminus B')$, the edges of $T_B \setminus T_{B'}$ give a path between them. If $v \in V(B \setminus B'), v' \in V(B')$, the edges of $T_B \setminus T_{B'}$ give a path into B' , where we can use the same path in $R(S'; B')$.

Finally we show that $c(S) \geq c(S')$. Let $S' = S_1 \sqcup \dots \sqcup S_k$ be the decomposition of S' into connected components. Suppose that there exist some set of edges $E \subseteq T_B \setminus T_{B'}$ that connect two components S_i and S_j . As $T_{B'}$ is a spanning tree, there exist edges $E' \subseteq T_{B'}$ that connect S_i, S_j that are disjoint from E . This implies $S_i \cup S_j \cup E \cup E'$ contains a cycle, and is contained in T_B , a contradiction as T_B a tree. Therefore S has at least as many components as S' . \square

From the previous two lemmas we observe that if $B \subset K_{d,n}$ is any connected bipartite graph then $\lambda(B) \leq \min\{d, n\}$.

3.2. TOPOLOGY OF TYPES. In this section, we investigate topological properties of certain subsets of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ induced by a tropical hyperplane arrangement. In particular, our goal is to show that these subsets are contractible, which will be central to applications to (co)cellular resolutions in Section 4. Note that none of our results in this section require \mathcal{H} to be generic nor have full support. The results here mimic those in [18, Section 2.3], but require some additional considerations to account for not necessarily full support.

We begin by recalling the notion of tropical convexity. It was established in [17, Theorem 2] that tropically convex sets are contractible.

DEFINITION 3.10 (Tropical convexity). *A subset S of \mathbb{R}^d is tropically convex if the set S contains the point $\mathbf{p} = c \odot \mathbf{x} \oplus d \odot \mathbf{y}$ for all $\mathbf{x}, \mathbf{y} \in S$ and $c, d \in \mathbb{R}$.*

If the distinction is important, we shall specify that a set is *max-tropically convex* (resp. *min*) if the tropical addition operation is \max (resp. \min). Note that all tropically convex sets contain $\mathbb{R}\mathbf{1}$, therefore we often view them as subsets of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. When we refer to a tropically convex set S in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$, we mean that $S + \mathbb{R}\mathbf{1}$ is tropically convex; recall that \oplus is not well defined on $\mathbb{R}^d/\mathbb{R}\mathbf{1}$.

Given a tropical matrix $A = (\mathbf{a}_1, \dots, \mathbf{a}_n)$ with entries in \mathbb{T}_{\min} , we can define the min-tropical polytope $\text{tconv}(A)$ to be the set of points of \mathbb{R}^d obtainable as a min-tropically convex combination of columns of A , i.e.

$$\text{tconv}(A) = \left\{ \mathbf{p} \in \mathbb{R}^d \mid \exists \lambda_i \in \mathbb{R} \text{ s.t. } \mathbf{p} = \bigoplus_{i=1}^n \lambda_i \odot \mathbf{a}_i \right\}.$$

NOTATION 3.11. For two fine types $T, T' \in \{0, 1\}^{n \times d}$, we write $T \leq T'$ if $T_{ij} \leq T'_{ij}$ for all i, j . This is equivalent to saying that the bipartite graph B_T is a subgraph of $B_{T'}$. We let $\min(T, T')$ and $\max(T, T')$ denote the table with entries given by the component-wise minimum and maximum, respectively. Equivalently, $B_{\min(T, T')} := B_T \cap B_{T'}$ is the largest subgraph of B_A contained in both T and T' , and $B_{\max(T, T')} := B_T \cup B_{T'}$ is the smallest subgraph of B_A containing both T and T' .

The following theorem is the main theorem of this subsection. It says that downsets of cells partially ordered by (either fine or coarse) type are tropically convex and hence contractible.

THEOREM 3.12. Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. Let $U \in \{0, 1\}^{n \times d}$ and $\mathbf{u} \in \mathbb{N}^d$. With labels determined by fine (resp. coarse) type, the following subsets of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ are max-tropically convex and hence contractible:

$$\begin{aligned} (\mathcal{C}(A), T)_{\leq U} &:= \{ \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \mid T_{\mathcal{H}}(\mathbf{p}) \leq U \} = \bigcup \{ C \in \mathcal{C}(A) \mid T_{\mathcal{H}}(C) \leq U \} \\ (\mathcal{C}(A), \mathbf{t})_{\leq \mathbf{u}} &:= \{ \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \mid \mathbf{t}_{\mathcal{H}}(\mathbf{p}) \leq \mathbf{u} \} = \bigcup \{ C \in \mathcal{C}(A) \mid \mathbf{t}_{\mathcal{H}}(C) \leq \mathbf{u} \}. \end{aligned}$$

Analogously, the following subsets of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ with labels determined by fine (respectively, coarse) cotype are min-tropically convex and hence:

$$\begin{aligned} (\mathcal{C}(A), \bar{T})_{\leq U} &:= \{ \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \mid \bar{T}_{\mathcal{H}}(\mathbf{p}) \leq U \} = \bigcup \{ C \in \mathcal{C}(A) \mid \bar{T}_{\mathcal{H}}(C) \leq U \} \\ (\mathcal{C}(A), \bar{\mathbf{t}})_{\leq \mathbf{u}} &:= \{ \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \mid \bar{\mathbf{t}}_{\mathcal{H}}(\mathbf{p}) \leq \mathbf{u} \} = \bigcup \{ C \in \mathcal{C}(A) \mid \bar{\mathbf{t}}_{\mathcal{H}}(C) \leq \mathbf{u} \}. \end{aligned}$$

The following lemma gives the first step towards proving Theorem 3.12.

LEMMA 3.13. Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$, and let $\mathbf{p}, \mathbf{q} \in \mathbb{R}^d/\mathbb{R}\mathbf{1}$. Then

$$\min(T_{\mathcal{H}}(\mathbf{p}), T_{\mathcal{H}}(\mathbf{q})) \leq T_{\mathcal{H}}(\mathbf{r}) \leq \max(T_{\mathcal{H}}(\mathbf{p}), T_{\mathcal{H}}(\mathbf{q}))$$

and

$$\mathbf{t}_{\mathcal{H}}(\mathbf{r}) \leq \max(\mathbf{t}_{\mathcal{H}}(\mathbf{p}), \mathbf{t}_{\mathcal{H}}(\mathbf{q}))$$

for $\mathbf{r} \in \text{tconv}^{\max}\{\mathbf{p}, \mathbf{q}\}$ on the max-tropical line segment between \mathbf{p} and \mathbf{q} .

Proof. To allow us to take tropical sums of points, we will prove the result in \mathbb{R}^d . Recall that hyperplane sectors and tropically convex sets are closed under scalar addition, and so the result will also hold in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$.

Let $\mathbf{r} = (\lambda \odot \mathbf{p}) \oplus (\mu \odot \mathbf{q}) = \max\{\lambda + \mathbf{p}, \mu + \mathbf{q}\}$ for $\lambda, \mu \in \mathbb{R}$, and fix some $k \in [d]$. Without loss of generality, assume that $r_k = \lambda + p_k \geq \mu + q_k$.

First we show that $\min(T_{\mathcal{H}}(\mathbf{p}), T_{\mathcal{H}}(\mathbf{q})) \leq T_{\mathcal{H}}(\mathbf{r})$. If at least one of the points \mathbf{p} and \mathbf{q} are not in the k th sector of some hyperplane $H(\mathbf{v}_i)$, then $\min(T_{\mathcal{H}}(\mathbf{p})_{ki}, T_{\mathcal{H}}(\mathbf{q})_{ki}) = 0$ and certainly $T_{\mathcal{H}}(\mathbf{r})_{ki} \geq 0$. Now suppose that both \mathbf{p} and \mathbf{q} are in the k th sector for some hyperplane $H(\mathbf{v})$, so $p_k - p_i \geq v_k - v_i$ and $q_k - q_i \geq v_k - v_i$ for all $i \in \text{supp}(H(\mathbf{v}))$. Fix some $j \in [d]$. If $r_j = \lambda + p_j \geq \mu + q_j$, then $r_j - r_k = p_j - p_k$ and r is in the k th sector of $H(\mathbf{v})$; similarly, if $r_j = \mu + q_j \geq \lambda + p_j$, then $r_j - r_k = q_j - q_k$ and r again lies in the k th sector of $H(\mathbf{v})$.

Now we show that $T_{\mathcal{H}}(\mathbf{r}) \leq \max(T_{\mathcal{H}}(\mathbf{p}), T_{\mathcal{H}}(\mathbf{q}))$. We show that if r is in the k th sector of some hyperplane $H(\mathbf{v})$, then at least one of \mathbf{p} and \mathbf{q} must be, too. Assume

that $r_k - r_j \geq v_k - v_j$ for all $j \in [d]$, and also assume without loss of generality that $r_k = \lambda + p_k \geq \mu + q_k$. Then $\lambda + p_k \geq v_k - v_j + r_j$ for all $j \in [d]$. We also know that $r_j \geq \lambda + p_j$ by the definition of r , so $\lambda + p_k \geq v_k - v_j + \lambda + p_j$. This implies that $p_k - p_j \geq v_k - v_j$, so \mathbf{p} is in sector k of the hyperplane $H(\mathbf{v})$.

The last claim follows from the definition of coarse types. □

With this lemma in hand, we may now prove the main theorem of this subsection.

Proof of Theorem 3.12. The max-tropical convexity of the sets $(\mathcal{C}(A), T)_{\leq U}$ and $(\mathcal{C}(A), \mathbf{t})_{\leq \mathbf{u}}$ follows from Lemma 3.13. Reasoning similar to that of the proof of Lemma 3.13 shows that if \mathbf{r} is a point in the min-tropical line segment between \mathbf{p} and \mathbf{q} , then $T_{\mathcal{H}}(\mathbf{r}) \geq \min(T_{\mathcal{H}}(\mathbf{p}), T_{\mathcal{H}}(\mathbf{q}))$; passing to complements, we obtain that $\overline{T}_{\mathcal{H}}(\mathbf{r}) \leq \max(\overline{T}_{\mathcal{H}}(\mathbf{p}), \overline{T}_{\mathcal{H}}(\mathbf{q}))$ and $\overline{\mathbf{t}}_{\mathcal{H}}(\mathbf{r}) \leq \max(\overline{\mathbf{t}}_{\mathcal{H}}(\mathbf{p}), \overline{\mathbf{t}}_{\mathcal{H}}(\mathbf{q}))$. This gives the claim for cotypes. □

We will also need to extend Theorem 3.12 to downsets of cotypes restricted to the bounded complex.

PROPOSITION 3.14. *The following subsets of $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ with labels determined by fine (respectively, coarse) cotype are contractible:*

$$\begin{aligned} (\mathcal{B}(A), \overline{T})_{\leq U} &:= \bigcup \{C \in \mathcal{B}(A) \mid \overline{T}_{\mathcal{H}}(C) \leq U\} \\ (\mathcal{B}(A), \overline{\mathbf{t}})_{\leq \mathbf{u}} &:= \bigcup \{C \in \mathcal{B}(A) \mid \overline{\mathbf{t}}_{\mathcal{H}}(C) \leq \mathbf{u}\}. \end{aligned}$$

Proof. We prove the theorem for downsets of fine cotypes, the proof for coarse cotypes is identical. If $(\mathcal{B}(A), \overline{T})_{\leq U}$ is empty then we are done, so assume it is non-empty.

We first note that every cell $C \subseteq \mathcal{C}(A)$ is a (min)-tropical polytope. To see this, observe that we can write C_T as the solution set of the linear inequalities:

$$\{x_i - a_{ik} \geq x_j - a_{jk} \mid T_{ik} = 1, a_{ik}, a_{jk} \neq \infty\}.$$

Each of these inequalities can also be written as min-tropical inequalities, and so C_T is the solution set of finitely many min-tropical inequalities. By the Tropical Minkowski-Weyl Theorem [31, Theorem 7.11], this implies that C_T is a min-tropical polytope.

For each cell $C \in (\mathcal{C}(A), \overline{T})_{\leq U}$, let V_C denote a finite generating set such that $C = \text{tconv}(V_C)$ and $V = \bigcup V_C$ the union of these finite generating sets. For any $p \in (\mathcal{C}(A), \overline{T})_{\leq U}$, we can write p as a min-tropical convex combination of elements of V , and so $(\mathcal{C}(A), \overline{T})_{\leq U} \subseteq \text{tconv}(V)$. However, the set $(\mathcal{C}(A), \overline{T})_{\leq U}$ is min-tropically convex itself by Theorem 3.12, and so $(\mathcal{C}(A), \overline{T})_{\leq U} = \text{tconv}(V)$.

Letting $m = |V|$, one can view V as a $d \times m$ tropical matrix. By [32, Theorem 33], $\text{tconv}(V)$ is obtainable as the upper hull of the *envelope* of the tropical matrix V (see also [31, Section 6.1])

$$\mathcal{E}(V) = \{(y, z) \in \mathbb{R}^d \times \mathbb{R}^m \mid y_i - z_j \leq v_{ij}\}$$

orthogonally projected onto \mathbb{R}^d . Moreover, this projection is an affine isomorphism when restricted to the bounded cells of $\mathcal{E}(V)$ and $\text{tconv}(V)$ respectively. Note that the bounded cells of $\text{tconv}(V)$ are precisely $(\mathcal{B}(A), \overline{T})_{\leq U}$.

As $(\mathcal{B}(A), \overline{T})_{\leq U}$ is non-empty, $\text{tconv}(V)$ has at least one bounded cell and therefore so does $\mathcal{E}(V)$. This makes $\mathcal{E}(V)$ a pointed polyhedron, a polyhedron with no lineality space. By [31, Example 10.54], $\mathcal{E}(V)$ is projectively equivalent to a polytope with a distinguished *face at infinity* and the set of faces of $\mathcal{E}(V)$ that are disjoint from the face at infinity are precisely the bounded cells of $\mathcal{E}(V)$. Therefore by [35, Theorem 4.17] the bounded part of $\mathcal{E}(V)$ is contractible. As this is affinely isomorphic to $(\mathcal{B}(A), \overline{T})_{\leq U}$, this implies that $(\mathcal{B}(A), \overline{T})_{\leq U}$ is also contractible. □




FIGURE 8. An arrangement of four degenerate tropical hyperplanes whose bounded complex is not tropically convex. The tropical line segment (blue) between any two points in the interior of different maximal cells must leave the bounded complex at some point.

The proof of Proposition 3.14 is much cleaner for arrangements with full support. When all entries of A are finite, the bounded complex $\mathcal{B}(A)$ agrees with the tropical polytope $\text{tconv}(A)$ and hence is min-tropically convex. This immediately implies $(\mathcal{B}(A), \overline{T})_{\leq U}$ is min-tropically convex as an intersection of min-tropically convex sets, and therefore contractible. When A has infinite entries, $\mathcal{B}(A)$ is only a subset of $\text{tconv}(A)$ and no longer tropically convex in general. Figure 8 shows an example of such a complex consisting of two maximal cells. For any pair of points in the interior of these two cells, the tropical line segment between them must leave the bounded complex.

4. IDEALS AND RESOLUTIONS FROM TROPICAL ARRANGEMENTS

In [18] the authors introduce monomial ideals arising from the type data associated to a tropical hyperplane arrangement, and prove that the facial structure of the various tropical complexes describes algebraic relations (syzygies) among the generators. In this section we extend these constructions and results to the case of tropical hyperplanes with not necessarily full support. In particular, we show that the cellular complexes from Definition 2.12 support (co-)cellular resolutions of certain ideals arising naturally from the combinatorics of the arrangement.

4.1. IDEALS FROM THE ARRANGEMENT. The ideals that we study will live in polynomial rings defined by the finite entries of a tropical matrix.

NOTATION 4.1. Suppose A is a $d \times n$ tropical matrix. We let X_A denote the $d \times n$ generic matrix with entries

$$X_A(i, j) = \begin{cases} x_{i,j} & \text{if } (i, j) \in \text{supp}(A) \\ 0 & \text{otherwise} \end{cases}$$

and for a field \mathbb{k} define $\tilde{S} = \mathbb{k}[X_A]$ to be the polynomial ring in the indeterminates in X_A . Note that if $B = B_A$ is the bipartite graph associated to A , then the number of indeterminates of \tilde{S} is given by $|E(B)|$, the number of edges of B . We let S denote the polynomial ring $S = \mathbb{k}[x_1, \dots, x_d]$.

Let the weight of the variable x_{ij} in \tilde{S} be v_{ij} and the weight of a monomial $\mathbf{x}^{\mathbf{a}} = \prod x_{ij}^{a_{ij}} \in \tilde{S}$ be $\sum a_{ij}v_{ij}$. The initial form $\text{in}_A(f)$ of a polynomial $f = \sum c_i \mathbf{x}^{\mathbf{a}_i}$ is defined to be the sum of terms $c_i \mathbf{x}^{\mathbf{a}_i}$ such that $\mathbf{x}^{\mathbf{a}_i}$ has maximal weight.

The fine and coarse (co)-type data from Definitions 2.8 and 2.9 naturally give rise to monomial ideals. These ideals were first introduced in the full-support case in [18] as a means of giving a tropical analogue of the *oriented matroid ideals* studied by Novik, Postnikov, and Sturmfels in [37].

DEFINITION 4.2 (Type/cotype ideals). *Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. The fine type ideal and fine cotype ideal associated to \mathcal{H} are the squarefree monomial ideals*

$$I_{T(\mathcal{H})} = \langle \mathbf{x}^{T(\mathbf{p})} \mid \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \rangle \subset \tilde{S}$$

$$I_{\bar{T}(\mathcal{H})} = \langle \mathbf{x}^{\bar{T}(\mathbf{p})} \mid \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \rangle \subset \tilde{S}$$

where $\mathbf{x}^{T(\mathbf{p})} = \prod \{x_{ij} \mid T(\mathbf{p})_{ij} = 1\}$. Analogously, the coarse type ideal and coarse cotype ideal associated to \mathcal{H} are given by

$$I_{\mathbf{t}(\mathcal{H})} = \langle \mathbf{x}^{\mathbf{t}(\mathbf{p})} \mid \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \rangle \subset S$$

$$I_{\bar{\mathbf{t}}(\mathcal{H})} = \langle \mathbf{x}^{\bar{\mathbf{t}}(\mathbf{p})} \mid \mathbf{p} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \rangle \subset S$$

where $\mathbf{x}^{\mathbf{t}(\mathbf{p})} = x_1^{t_1} \dots x_n^{t_n}$ with $\mathbf{t}(\mathbf{p}) = (t_1, \dots, t_d)$.

We will see in the following sections that an arrangement \mathcal{H} gives rise to complexes that support minimal free resolutions of these ideals. This provides a means to understand the projective dimension and regularity of our type ideals. Later in the paper we will use our techniques to obtain a formula for the Krull dimension as well, see Section 5.1.

4.2. CELLULAR RESOLUTIONS OF TYPE AND COTYPE IDEALS. Next we show that the tropical complex and bounded complex of a tropical hyperplane arrangement support minimal (co)cellular resolutions of the various type and cotype ideals. This provides a tropical analogue of the work of [37] and generalizes results from [9] and [18]. For a reference on cellular resolutions we refer the reader to [35, Chapter 4]; for details on cocellular resolutions, see [35, Section 5.3]. The main criteria we use in this section to check whether a resolution is cellular is [35, Prop. 4.5], and the analogous co-cellular version which follows from the same proof techniques.

PROPOSITION 4.3. *Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. For every cell $C \in \mathcal{C}(A)$ of codimension at least 1,*

$$T_{\mathcal{H}}(C) = \max\{T_{\mathcal{H}}(D) \mid C \subset D\}, \text{ and}$$

$$\mathbf{t}_{\mathcal{H}}(C) = \max\{\mathbf{t}_{\mathcal{H}}(D) \mid C \subset D\}.$$

Therefore, both the fine type and the coarse type yield a monomial colabeling for the complex $\mathcal{C}(A)$.

Similarly, for every cell $D \in \mathcal{C}(A)$ of dimension ≥ 1 we have

$$\bar{T}_{\mathcal{H}}(D) = \max\{\bar{T}_{\mathcal{H}}(C) \mid C \subset D\}, \text{ and}$$

$$\bar{\mathbf{t}}_{\mathcal{H}}(D) = \max\{\bar{\mathbf{t}}_{\mathcal{H}}(C) \mid C \subset D\}.$$

Thus, both the fine and coarse cotypes yield monomial labelings for the complex $\mathcal{C}(A)$.

The proof is essentially identical to that of [18, Propositions 3.4 and 3.8].

PROPOSITION 4.4. *Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding tropical hyperplane arrangement in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ with tropical complex $\mathcal{C}(A)$. Then with labels given by fine (respectively coarse) type, the labeled complex $\mathcal{C}(A)$ supports a minimal cocellular resolution of the fine (respectively coarse) type ideal $I_{T_{\mathcal{H}}}$ (respectively $I_{\bar{T}_{\mathcal{H}}}$).*

Proof. Proposition 4.3 implies that the polyhedral complex $\mathcal{C}(A)$ is colabeled by both fine and coarse types. Combining Theorem 3.12 and a cocellular version of the criterion in [35, Prop. 4.5] gives the result. Minimality follows from the fact that if C is strictly contained in D , then $T_{\mathcal{H}}(D) < T_{\mathcal{H}}(C)$; see e.g. [17, Corollary 13]. \square

In a similar argument to that above, one can combine Proposition 4.3, Proposition 3.14, and [35, Prop. 4.5] to obtain the following result.

PROPOSITION 4.5. *Suppose A is a $d \times n$ tropical matrix, let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of tropical hyperplanes, and let $\mathcal{B}(A)$ be the subcomplex of bounded cells of $\mathcal{C}(A)$ (if it exists). Then $\mathcal{B}(A)$, with labels given by the fine (resp. coarse) cotype, supports a minimal cellular resolution of the fine (resp. coarse) cotype ideal.*

4.3. RESOLUTIONS SUPPORTED ON MIXED SUBDIVISIONS OF GENERALIZED PERMUTOHEDRA. Recall from Definition 2.19 that the support of a tropical matrix A also gives rise to a generalized permutohedron P_{B_A} , given by

$$P_{B_A} = \sum_{j=1}^n \Delta_{N_j(B_A)} \subseteq \mathbb{R}^d.$$

Furthermore, the entries of the matrix A gives rise to a regular mixed subdivision $\mathcal{M}(A)$ of P_{B_A} . We define a ‘fine monomial labeling’ on $\mathcal{M}(A)$ (with monomials in \tilde{S}) in the following way. Given a face $F = \sum_{j=1}^n \Delta_{I_j}$, where $I_j \subseteq N_{B_A}(j)$, we assign to F the monomial

$$(2) \quad \prod_{j \in [n]} \prod_{i \in I_j} x_{ij}.$$

Similarly we obtain a ‘coarse monomial labeling’ of P_{B_A} (with monomials in S) by specializing the monomials above to the second index $x_{ij} \mapsto x_j$. Note that under this coarse labeling a 0-cell v of P_{B_A} is labeled by a monomial whose exponent vector is precisely the coordinates of v . See Figure 2 for an example. Using Proposition 2.23, it turns out these monomial labelings are dual to the type labelings of the corresponding tropical complex. From this we observe the following.

COROLLARY 4.6. *For any $d \times n$ tropical matrix A , the monomial labeled mixed subdivision $\mathcal{M}(A)$ of the generalized permutohedron P_{B_A} , with labels given by fine (respectively coarse) type, supports a minimal cellular resolution of the ideal $I_{T(\mathcal{H})}$ (respectively $I_{\mathfrak{t}(\mathcal{H})}$).*

Proof. From Proposition 2.23 we have that the tropical complex $\mathcal{C}(A)$ is dual to the mixed subdivision $\mathcal{M}(A)$ of P_{B_A} , and in addition the fine (respectively coarse) monomial labeling of a cell in P_{B_A} has the same fine (respectively coarse) type of the corresponding cell in $\mathcal{C}(A)$. The result then follows from Proposition 4.4. \square

COROLLARY 4.7. *For a sufficiently generic $d \times n$ tropical matrix A , the coarse type ideal $I_{\mathfrak{t}(\mathcal{H})}$ is generated by monomials corresponding to the lattice points of P_{B_A} .*

Proof. Corollary 4.6 gives that the generators are the labels of the 0-dimensional cells of $\mathcal{M}(A)$. It suffices to show this is all lattice points when A is sufficiently generic, i.e. when $\mathcal{M}(A)$ is a fine mixed subdivision.

Every 0-dimensional cell is a sum $\sum_{j=1}^n \mathbf{e}_{k_j}$ for some $k_j \in N_j(B_A)$, so is a lattice point. Conversely, let \mathbf{p} be a lattice point and $C = \Delta_{I_1} + \cdots + \Delta_{I_n}$ the smallest fine mixed cell containing \mathbf{p} . By [40, Lemma 14.12], we can write $\mathbf{p} = \sum_{j=1}^n \mathbf{e}_{k_j}$ where $k_j \in I_j$. If $C \neq \{\mathbf{p}\}$, then \mathbf{p} is contained in the relative interior of C , and so we can perturb \mathbf{p} in any direction in the affine span of C . The mixed subdivision structure

implies that there exists some $k, l \in [d]$ such that $\mathbf{p} \pm (\mathbf{e}_k - \mathbf{e}_l) \in C$. In particular, this means $k, l \in I_s \cap I_t$ for distinct $s, t \in [n]$. This contradicts C being a fine mixed cell, as $\dim(\Delta_{I_s} + \Delta_{I_t}) \neq \dim(\Delta_{I_s}) + \dim(\Delta_{I_t})$. Therefore $C = \{\mathbf{p}\}$, and so is a cell of $\mathcal{M}(A)$. \square

Recall that a minimal free resolution of any finitely generated \tilde{S} -module is unique up to isomorphism. The fact that any regular fine mixed subdivision of P_{B_A} supports a minimal resolution of $I_{T(\mathcal{H})}$ allows us to recover the strong equidecomposability of such complexes, generalizing [18, Corollary 5].

COROLLARY 4.8. *Suppose $B \subset K_{d,n}$ is a bipartite graph and let P_B denote the generalized permutohedron associated to B . Then for any fine mixed subdivision of P_B , the collection of coarse types $\mathbf{t}(C)$ for cells C with $\dim C = k$, counted with multiplicities, is the same. In particular, any fine mixed subdivision of P_B has the same f -vector.*

4.4. ALEXANDER DUALITY AND INITIAL IDEALS OF LATTICE IDEALS. In the case of arrangements of tropical hyperplanes with full support, Block and Yu [9] observed that the fine cotype ideal is Alexander dual to a corresponding initial ideal of a certain toric ideal. Here we establish a similar result for the case of arbitrary A , which will be important for our applications to toric edge ideals studied in Section 5. We first recall some relevant notions.

DEFINITION 4.9 (Lattice ideal). *Let \mathcal{A} be a set of vectors in \mathbb{Z}^n . Then the lattice ideal associated to \mathcal{P} is the ideal $I_{\mathcal{A}} = \ker \varphi_{\mathcal{A}}$, where*

$$\begin{aligned} \varphi_{\mathcal{A}} : \mathbb{k}[x_{\mathbf{p}} \mid \mathbf{p} \in \mathcal{A}] &\longrightarrow \mathbb{k}[x_1, \dots, x_n] \\ x_{\mathbf{p}} &\longmapsto x_1^{p_1} \cdots x_n^{p_n}. \end{aligned}$$

In the case of a root polytope Q_B associated to a bipartite graph B , we have $\mathcal{A} = \{e_i - e_{\bar{j}} \mid (i, \bar{j}) \in E(B)\}$ and the lattice ideal of Q_B will be the kernel of the map

$$(3) \quad \begin{aligned} \varphi_{\mathcal{A}} : \mathbb{k}[x_{ij} \mid (i, \bar{j}) \in E(B)] &\longrightarrow \mathbb{k}[x_1, \dots, x_d, y_1^{-1}, \dots, y_n^{-1}] \\ x_{ij} &\longmapsto x_i y_j^{-1}. \end{aligned}$$

REMARK 4.10. Since Q_B is unimodular, it is known that monomial initial ideals of the lattice ideal of Q_B are in bijection with regular triangulations of Q_B . As we have seen, regular triangulations of Q_B are encoded by sufficiently generic tropical hyperplane arrangements $\mathcal{H}(A)$ where A satisfies $B = B_A$.

EXAMPLE 4.11. Let Q_B be the root polytope corresponding to the bipartite graph in our running example (see Figure 1), where

$$A = \begin{bmatrix} 0 & 3 & \infty & 0 \\ 0 & 0 & 2 & 3 \\ 0 & \infty & 0 & \infty \end{bmatrix}.$$

Then the lattice ideal $L = I_{\mathcal{A}}$ of Q_B is the ideal generated by

$$(4) \quad \langle \underline{x_{11}x_{22}} - \underline{x_{12}x_{21}}, \underline{x_{11}x_{24}} - x_{14}x_{21}, \underline{x_{12}x_{24}} - x_{14}x_{22}, x_{21}x_{33} - \underline{x_{23}x_{31}} \rangle.$$

Recall from Notation 4.1 that a $d \times n$ tropical matrix A gives rise to a monomial term order on \tilde{S} and hence an initial ideal $\text{in}_A(I)$ for any ideal $I \subset \tilde{S}$. The underlined terms correspond to the generators of the leading term ideal of L with respect to the monomial term order on \tilde{S} induced by A .

DEFINITION 4.12 (Crosscut complex). *Let $\mathcal{S}(A)$ the subdivision of Q_{B_A} induced by A . Define the crosscut complex $\text{CrossCut}(\mathcal{S}(A))$ to be the unique simplicial complex with the same vertices-in-facet incidences as the polyhedral complex $\mathcal{S}(A)$.*

REMARK 4.13. If $\mathcal{S}(A)$ is a triangulation of Q_{B_A} , then $\text{Crosscut}(\mathcal{S}(A)) = \mathcal{S}(A)$.

PROPOSITION 4.14. Let A be a $d \times n$ tropical matrix and let $\mathcal{H} = \mathcal{H}(A)$ denote the associated arrangement of tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. Let L denote the lattice ideal of Q_{B_A} (or, equivalently, the toric edge ideal of B_A). Then the fine cotype ideal $I_{\overline{\mathcal{T}}(\mathcal{H})}$ associated to \mathcal{H} is the Alexander dual of $M(\text{in}_A(L))$, where $M(\text{in}_A(L))$ is the largest monomial ideal contained in $\text{in}_A(L)$.

Proof. Let J be the Stanley–Reisner ideal of $\text{Crosscut}(\mathcal{S}(A))$. Then J is the Alexander dual of the fine cotype ideal associated to \mathcal{H} .

It remains to check that $J = M(\text{in}_A(L))$. Because \mathcal{A} is unimodular, one has that $\text{in}_A(L)$ is squarefree and therefore $M(\text{in}_A(L))$ is radical. The proof of [18, Lemma 6.7] applies directly here, so one has

$$M(\text{in}_A(L)) = \text{Rad}(M(\text{in}_A(L))) = J$$

as desired. □

Combining Propositions 4.5 and 4.14 gives the following result. Here for a squarefree monomial ideal I we use I^\vee to denote its Alexander dual.

COROLLARY 4.15. For any tropical matrix A the bounded complex $\mathcal{B}(A)$ of the corresponding tropical hyperplane arrangement supports a minimal cellular resolution of the ideal $M(\text{in}_A(L))^\vee$. In particular, if $\mathcal{H}(A)$ is sufficiently generic, then $\text{in}_A(L)$ is a monomial ideal and $\mathcal{B}(A)$ supports a minimal free resolution of $(\text{in}_A(L))^\vee$.

Moreover, if A is sufficiently generic, one can say more about the resolution of the (fine or coarse) cotype ideal.

COROLLARY 4.16. If A is sufficiently generic, then $(\text{in}_A(L))^\vee$ has a linear minimal free resolution.

Proof. If A is sufficiently generic, then $\mathcal{S}(A)$ is a triangulation of Q_{B_A} corresponding to the Stanley–Reisner complex of the (squarefree) monomial ideal $\text{in}_A(L)$ [48, Theorem 8.3]. In this case, $\mathcal{B}(A)$ supports a minimal linear free resolution of $(\text{in}_A(L))^\vee$. This follows from the following observations:

- (i) If A is sufficiently generic, then the labels on the vertices of $\mathcal{B}(A)$ are all of the same degree: each vertex is “outside of” the same number of regions as every other vertex.
- (ii) Let D be a cell in $\mathcal{B}(A)$. For any cell $C \subset D$ with $\dim C = \dim D - 1$, the cotype $T_{\mathcal{H}}(C)$ differs from $T_{\mathcal{H}}(D)$ in exactly one spot if A is sufficiently generic. Therefore, the monomial labeling of the cell D is exactly one degree higher than the monomial labeling of any cell $C \subset D$ such that $\dim C = \dim D - 1$.

□

EXAMPLE 4.17. Consider the tropical hyperplane arrangement of Figure 1. The bounded complex supports a minimal cellular resolution of the fine cotype ideal

$$I_{\overline{\mathcal{T}}(\mathcal{H})} = \langle x_{12}x_{23}x_{24}, x_{11}x_{12}x_{23}, x_{11}x_{31}x_{12}, x_{31}x_{12}x_{24}, x_{21}x_{31}x_{24}, x_{21}x_{23}x_{24} \rangle.$$

The Alexander dual of this ideal in the polynomial ring $\mathbb{k}[x_{ij} \mid (i, j) \in \text{supp}(A)]$ is

$$\langle x_{21}x_{12}, x_{31}x_{23}, x_{11}x_{24}, x_{12}x_{24} \rangle$$

which is exactly the ideal $\text{in}_A(L)$ corresponding to the underlined leading terms of the ideal L from (4).

EXAMPLE 4.18. Now consider the tropical hyperplane arrangement associated to the tropical matrix

$$A' = \begin{bmatrix} 0 & 0 & \infty & 0 \\ 0 & 0 & 2 & 3 \\ 0 & \infty & 0 & \infty \end{bmatrix}.$$

This arrangement corresponds to the same one as in Figure 1, except that now the hyperplane corresponding to v_2 lies on top of the hyperplane corresponding to v_1 , so it is no longer generic. However, the bipartite graph for A and A' coincide, and therefore so do their lattice ideals L . In this case, the fine cotype ideal is

$$I_{\overline{T}(\mathcal{H}(A'))} = \langle x_{23}x_{24}, x_{24}x_{31}, x_{11}x_{31}x_{12}, x_{11}x_{12}x_{23} \rangle$$

and the Alexander dual of the fine cotype ideal has generators

$$\langle x_{11}x_{24}, x_{12}x_{24}, x_{23}x_{31} \rangle.$$

This coincides with the largest monomial ideal inside of the initial ideal of L in (4) with respect to the term order induced by A' , which is

$$\text{in}_{A'}(L) = \langle x_{11}x_{22} - x_{12}x_{21}, x_{11}x_{24}, x_{12}x_{24}, x_{23}x_{31} \rangle.$$

4.5. VOLUMES AND SYZYGIES. Recall that a bipartite graph $B \subset K_{d,n}$ encodes a generalized permutohedron P_B obtained as a sum of simplices, as described in Definition 2.19. For any tropical matrix A satisfying $B = B_A$ we have from Corollary 4.6 that the corresponding mixed subdivision of P_B supports a minimal cellular resolution of the resulting fine type ideal $I_{T(\mathcal{H})}$ as well as the coarse type ideal $I_{\mathfrak{t}(\mathcal{H})}$. In particular, the Betti numbers of these ideals can be read off from the f -vectors of the underlying subdivision. In many cases the volume of these cells can also be understood in terms of combinatorial data, and in this section we explore this connection.

4.5.1. *Graphic tropical hyperplane arrangements.* The most straightforward application arises in the case of tropical arrangements where the apex of each hyperplane has exactly two finite coordinates. This inspires the following definition.

DEFINITION 4.19. *An arrangement $\mathcal{H}(A)$ of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ is graphic if each column \mathbf{a}_i in the underlying $d \times n$ matrix $A = (\mathbf{a}_1, \dots, \mathbf{a}_n)$ has exactly two finite coordinates.*

We explain the choice of terminology. First note that a graphic tropical hyperplane arrangement $\mathcal{H}(A)$ in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ consists of Euclidean subspaces that are each dual to a line segment, so that $\mathcal{H}(A)$ is in fact a collection of (ordinary) affine hyperplanes in $(d - 1)$ -dimensional Euclidean space. In fact if a column $\mathbf{a}_k \in A$ has finite entries a_{ik} and a_{jk} then the corresponding hyperplane $H(\mathbf{a}_k)$ consists of the set

$$\{\mathbf{x} \in \mathbb{R}^d/\mathbb{R}\mathbf{1} \mid x_i - x_j = a_{jk} - a_{ik}\}.$$

We can now think of the rows of A as vertices of a graph $\mathcal{G}(A)$, where each column of A describes an edge. One can then see that the tropical hyperplane arrangement $\mathcal{H}(A)$ corresponds to a perturbation of the classical (central) *graphic hyperplane arrangement* associated to the graph $\mathcal{G}(A)$.

For a graphic tropical hyperplane arrangement $\mathcal{H}(A)$, the corresponding generalized permutohedron P_{B_A} is the (*graphic*) *zonotope* $\mathcal{Z}_{\mathcal{G}(A)}$ given by the sum of line segments of the form $[\mathbf{e}_i, \mathbf{e}_j]$, where i and j are the finite entries in some column of A (or in other words some edge in the graph $\mathcal{G}(A)$). A generic perturbation of the central graphic hyperplane arrangement corresponds to a fine mixed subdivision (in this case a *zonotopal tiling*) of $\mathcal{Z}_{\mathcal{G}(A)}$. In our case the subdivision we obtain is dual to the subdivision of tropical space $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ induced by the sufficiently generic tropical hyperplane arrangement $\mathcal{H}(A)$ discussed above.

EXAMPLE 4.20. As an example, we can take A to be the tropical matrix

$$A = \begin{bmatrix} 0 & \infty & \infty & 0 \\ 0 & 1 & \infty & \infty \\ \infty & 2 & 4 & \infty \\ \infty & \infty & 6 & 9 \end{bmatrix}$$

One can see that $\mathcal{G}(A)$ is a 4-cycle, and the corresponding zonotope $\mathcal{Z}_{\mathcal{G}(A)}$ is depicted in Figure 9 as a Minkowski sum of edges of the simplex. That fine mixed subdivision (zonotopal tiling) of $\mathcal{Z}_{\mathcal{G}(A)}$ induced by A is also depicted, computed via `Polymake` [23].




FIGURE 9. The zonotope $\mathcal{Z}_{\mathcal{G}(A)}$ and its fine mixed subdivision from Example 4.20.

Now suppose $\mathcal{H} = \mathcal{H}(A)$ is a sufficiently generic graphic tropical hyperplane arrangement with underlying graph $\mathcal{G} = \mathcal{G}(A)$. Combining our results with well-known properties of zonotopes provides enumerative interpretations of certain Betti numbers of our type ideals.

COROLLARY 4.21. *Suppose $\mathcal{H} = \mathcal{H}(A)$ is a graphic tropical hyperplane arrangement with underlying connected graph $\mathcal{G} = \mathcal{G}(A)$. Then the number of generators of the fine type ideal $I_T(\mathcal{H})$, as well as the coarse type ideal $I_{\mathbf{t}(\mathcal{H})}$, is given by the number of forests in \mathcal{G} . The rank of the highest syzygy module of both $I_T(\mathcal{H})$ and $I_{\mathbf{t}(\mathcal{H})}$ is given by the number of spanning trees of \mathcal{G} .*

Proof. In [40, Proposition 2.4] it is shown that for a connected graph \mathcal{G} the volume of the graphic zonotope $\mathcal{Z}(\mathcal{G})$ equals the number of spanning trees of \mathcal{G} , and also that the number of lattice points of $\mathcal{Z}(\mathcal{G})$ equals the number of forests in \mathcal{G} . Our first claim then follows from the fact that $\mathcal{Z}(\mathcal{G})$ supports a minimal resolution of both $I_T(\mathcal{H})$ and $I_{\mathbf{t}(\mathcal{H})}$, whose generators in each case are in bijection with the lattice points of $\mathcal{Z}(\mathcal{G})$ (see Corollaries 4.6 and 4.7). The second claim follows from the fact that any maximal cell in a zonotopal tiling (in this case a parallelepiped corresponding to a spanning tree of \mathcal{G}) has volume 1. \square

REMARK 4.22. We observe that the type ideals coming from graphic tropical hyperplane arrangements can be realized as *oriented matroid ideals* studied by Novik, Postnikov, and Sturmfels in [37]. As we have seen, a graphic tropical hyperplane arrangement can be realized as a perturbation of the corresponding central graphic

hyperplane arrangement. The oriented matroid ideal of the resulting affine arrangement can be seen to coincide with the underlying fine type ideal.

On the other hand, the authors of [37] study a different oriented matroid ideal arising from a graph \mathcal{G} . Namely, they use the affine oriented matroid \mathcal{A} obtained by intersecting the central graphic hyperplane arrangement of \mathcal{G} by a generic affine hyperplane. It is proved that the resulting oriented matroid ideal $O_{\mathcal{A}}$ is Cohen-Macaulay, and in [37] the authors provide formulas and interpretations for the Möbius invariants and coinvariants of both the complete graph K_n and complete bipartite graph $K_{m,n}$.

In particular, our methods provide a different way to obtain an oriented matroid ideal from a graph \mathcal{G} . It is not clear to us how these two ideals relate (note they live in different polynomial rings).

Graphic tropical hyperplane arrangements also have connections to G -parking function ideals and other concepts coming from the theory of chip-firing on a graph G . Such ideals M_G were first introduced by Postnikov and Shapiro in [41], where connections to power ideals were studied. In [19] Dochtermann and Sanyal proved that a minimal cellular resolution of M_G is supported on a certain affine slice of the graphic hyperplane arrangement associated to G . One can recover this ideal as a certain coarse type ideal associated to the underlying graphic tropical hyperplane arrangement, leading to a number of applications and generalizations. See Section 7 for more discussion.

4.5.2. *Computing volumes via Alexander duality.* For P_B a generalized permutohedron obtained as a sum of simplices Postnikov [40, Theorem 9.3] provides a formula for the volume of P_B in terms of what he calls G -draconian sequences (where $G = B$ in our context). Given a sufficiently generic $d \times n$ tropical matrix A with $B_A = B$, one may also obtain the volume of P_B by studying the types associated to the vertices of the tropical complex.

In what follows, recall that a cell in the bounded complex $\mathcal{B}(A)$ can be described by its type T . Also recall from Definition 2.5 that for any fine type T we denote by $\text{RD}(T) = (a_1, \dots, a_n)$ the right degree vector of T .

PROPOSITION 4.23. *Let A be a sufficiently generic $d \times n$ tropical matrix with corresponding tropical hyperplane arrangement $\mathcal{H}(A)$, and set $B = B_A \subseteq K_{d,n}$. Let \mathcal{S} be the set of fine types associated to the vertices of the bounded complex $\mathcal{B}(A)$. Then the volume of the associated generalized permutohedron is*

$$\text{Vol } P_B = \sum_{T \in \mathcal{S}} \frac{1}{(\text{RD}(T)_1 - 1)! \cdots (\text{RD}(T)_n - 1)!}$$

Proof. Note that every element $T \in \mathcal{S}$ is in bijection with a full-dimensional fine mixed cell M_T in $\mathcal{M}(A)$. We can read off the volume of M_T in the following way. Let B_T denote its fine type graph and $\text{RD}(B_T) = (a_1, a_2, \dots, a_n)$ its right degree sequence. Then one can see that

$$\text{Vol}(M_T) = \prod_{i=1}^n \frac{1}{(a_i - 1)!},$$

as the volume of M_T is the product of the volumes of the simplices in the Minkowski sum. Summing over the elements of \mathcal{S} provides the desired volume of P_B . \square

Comparing this with Postnikov’s formula for volumes of generalized permutohedra (see [40, Theorem 9.3]) yields the following.

COROLLARY 4.24. *Let A be a sufficiently generic $d \times n$ tropical matrix with corresponding hyperplane arrangement $\mathcal{H}(A)$, and set $B = B_A \subseteq K_{d,n}$. Let \mathcal{S} be the set of*

fine types associated to the vertices of the bounded complex associated to \mathcal{H} , and let $\mathbb{1} = (1, \dots, 1) \in \mathbb{Z}^n$. Then the B -draconian sequences correspond to the set

$$\{\text{RD}(T) - \mathbb{1} \mid T \in \mathcal{S}\}.$$

Given a bipartite graph $B \subseteq K_{d,n}$, this gives a novel method to compute and prove statements about volumes of generalized permutohedra P_B :

- (1) Compute the lattice ideal L of the associated root polytope Q_B .
- (2) Find a sufficiently generic tropical matrix A such that $B_A = B$ and compute $\text{in}_A L$.
- (3) Compute the Alexander dual $(\text{in}_A L)^\vee = I_{\overline{T}(\mathcal{H}(A))}$ (see Proposition 4.14).
- (4) Take the complements of the monomial generators of $I_{\overline{T}(\mathcal{H}(A))}$ to obtain the set \mathcal{S} described above.
- (5) Compute P_B using the formula in Proposition 4.23.

EXAMPLE 4.25. Consider the generalized permutohedron defined by $B = K_{3,2}$, so that P_B is the sum of two triangles. To compute its volume we choose the sufficiently

generic tropical matrix $A = \begin{bmatrix} 0 & 0 \\ 0 & 2 \\ 0 & 1 \end{bmatrix}$. This defines an initial ideal

$$\langle \underline{x_{11}x_{22}} - x_{12}x_{21}, \underline{x_{11}x_{32}} - x_{12}x_{31}, x_{21}x_{32} - \underline{x_{31}x_{22}} \rangle,$$

generated by the underlined monomials. The Alexander dual of this initial ideal is given by the fine cotype ideal

$$I_{\overline{T}(\mathcal{H}(A))} = \langle x_{11}x_{22}, x_{11}x_{31}, x_{22}x_{32} \rangle.$$

Each of these generators defines a fine cotype, and the right degree sequences of the corresponding types is given by $(2, 2)$, $(1, 3)$, $(3, 1)$. This provides cells of volume 1, $1/2$, $1/2$, implying that P_B has volume 2. Note also that the G -draconian sequences for the case $G = K_{3,2}$ are given by $(1, 1)$, $(0, 2)$, and $(2, 0)$.

5. LATTICE IDEALS OF ROOT POLYTOPES AS TORIC EDGE IDEALS

In this section we describe a connection between cotype ideals associated to tropical hyperplane arrangements and the class of *toric edge ideals*. The key observation here is that the lattice ideal of a root polytope Q_B for a *bipartite* graph B is exactly the toric edge ideal of the corresponding graph B , see Lemma 5.4. With this we are able to describe the generators of Q_B using a well-known result of Villarreal, and describe certain classes of lattice ideals which are well-studied by commutative algebraists. We then use results from Section 4 to show that the regularity of a toric edge ideal of a bipartite graph B coincides with the dimension of a corresponding bounded tropical complex. This leads to new proofs of existing bounds on homological invariants as well as new results. We begin by establishing notation and recalling some basic results regarding toric edge ideals.

NOTATION 5.1. Suppose $G = (V, E)$ is a finite simple graph with vertex set $V = \{v_1, \dots, v_n\}$ and edge set $E = \{e_1, \dots, e_k\}$. If \mathbb{k} is any field, let $\mathbb{k}[E]$ and $\mathbb{k}[V]$ denote the polynomial rings in these indeterminates.

DEFINITION 5.2. Suppose $G = (V, E)$ is a finite simple graph. The edge ring $\mathbb{k}[G]$ is the \mathbb{k} -algebra generated by the monomials $x_i x_j$ where $\{i, j\} \in E(G)$.

We define a monomial map

$$\begin{aligned} \pi : \mathbb{k}[E] &\rightarrow \mathbb{k}[V] \\ e_j &\mapsto v_{j_1} v_{j_2} \end{aligned}$$

where $e_j = \{v_{j_1}, v_{j_2}\}$. The toric edge ideal of G is defined to be the ideal $J_G = \ker \pi$, so that

$$\mathbb{k}[G] \cong \mathbb{k}[E]/J_G.$$

An effective way to study the edge ring of a graph G is via a certain polytope associated to G called the *edge polytope*.

DEFINITION 5.3. Let G be a finite simple graph with vertex set $V = [n]$. For an edge $e = \{i, j\}$ in G define $\rho(e) \in \mathbb{R}^n$ by $\rho(e) = \mathbf{e}_i + \mathbf{e}_j$, where \mathbf{e}_i is the i th unit coordinate vector in \mathbb{R}^n . The edge polytope \mathcal{P}_G is the convex hull of all such vectors

$$\mathcal{P}_G = \text{conv}\{\rho(e) \mid e \in E(G)\} \subset \mathbb{R}^n.$$

One can see that for any graph G , the toric edge ideal J_G is recovered as the toric ideal of the edge polytope \mathcal{P}_G . For this note that $\mathcal{P}_G \cap \mathbb{Z}^n = \{\rho(e) : e \in E(G)\}$ and also that the vertices of \mathcal{P}_G coincide with $\mathcal{P}_G \cap \mathbb{Z}^n$ (see [27, Lemma 5.2]). For the case that $G = B \subset K_{d,n}$ is *bipartite* the ideal J_G can be recovered in another way. The proof of the following lemma is straightforward.

LEMMA 5.4. If $B \subset K_{d,n}$ is a bipartite graph, then the toric edge ideal J_B is equal to the lattice ideal of the root polytope Q_B .

Recall that the Krull dimension of a toric ring associated to a matrix M is given by $\text{rank}(M)$ [27, Proposition 3.1]. The toric edge ring $\mathbb{k}[B]$ of a bipartite graph $B \subset K_{d,n}$ is the toric edge ring of its incidence matrix, which is known to have rank $d + n - 1$. Hence we get the following.

COROLLARY 5.5. For a bipartite graph $B \subset K_{d,n}$ the Krull dimension of $\mathbb{k}[B]$ is given by

$$\dim(\mathbb{k}[B]) = d + n - 1.$$

Toric edge rings are a particular example of a construction called *special fiber rings*, which are blow-up algebras that give insight into the *Rees algebra* of an ideal but which are often more tractable to compute. In general, there is interest in bounding the regularity of the Rees algebra of an ideal I because this provides information about the regularity of powers of I . Finding the defining ideal of the special fiber ring is a challenging problem and is still open for many classes of ideals. However, a famous result of Villarreal [51] tells us that we can write down a minimal generating set for the special fiber of the edge ideal of a bipartite graph by studying its minimal cycles.

The following proposition gives an explicit generating set for the toric edge ideal of a graph.

PROPOSITION 5.6. [51, Proposition 3.1] Let G be a finite simple graph. For any even closed walk $\mathbf{w} = (i_1, j_1, \dots, i_k, j_k)$ in G , define a binomial

$$f_{\mathbf{w}} := x_{i_1 j_1} x_{i_2 j_2} \dots x_{i_k j_k} - x_{j_1 i_2} x_{j_2 i_3} \dots x_{j_{k-1} i_k} x_{j_k i_1}.$$

Then the toric edge ideal J_G is generated by all $f_{\mathbf{w}}$ where \mathbf{w} is an even closed walk in G . In particular, if G is bipartite then J_G is generated by all $f_{\mathbf{w}}$ where \mathbf{w} is a minimal even cycle in G .

A bipartite graph B is *chordal bipartite* if it contains no induced cycles of length longer than four. In [38], the following algebraic characterization of the toric edge rings of such graphs is given.

THEOREM 5.7. [38] Suppose B is a finite connected bipartite graph. Then the following are equivalent.

- (1) B is chordal bipartite;
- (2) The toric edge ideal J_B admits a quadratic binomial Gröbner basis;
- (3) The edge ring $\mathbb{k}[B]$ is Koszul;
- (4) The toric edge ideal J_B is generated by quadratic binomials.

Toric edge ideals of chordal bipartite graphs also have a connection to certain *determinantal* ideals. For this recall that for a $d \times n$ tropical matrix A we use X_A to denote the matrix associated to the finite entries A , see Notation 4.1. We will let L_A denote the lattice ideal associated to the root polytope Q_{B_A} . Using the language established in Laura Ballard’s thesis [6], a 2-minor of X_A is called a *distinguished minor* if it is a 2-minor involving only nonzero entries of X_A . The following result also appeared in [6].

PROPOSITION 5.8. *Suppose A is a tropical matrix with bipartite graph B_A . If B_A is chordal bipartite, then the lattice ideal associated to the root polytope Q_{B_A} is generated by all distinguished minors of X_A .*

Observe that in the case that the tropical matrix A has full support, L_A is generated by all 2-minors of a generic $d \times n$ matrix as in [9].

EXAMPLE 5.9 (Ladder determinantal ideals, Ferrers graphs). Suppose the tropical matrix A satisfies the following property:

$$\text{if } (i, j), (h, k) \in \text{Supp}(A) \text{ and } i \leq h, j \leq k, \text{ then } (i, k), (h, j) \in \text{Supp}(A).$$

Then B_A is chordal bipartite and L_A is called a *ladder determinantal ideal*. Ladder determinantal ideals have been well-studied in commutative algebra and algebraic geometry; see work of Conca [12, 13], Gorla [24], and more recently Rajchgot, Robichaux, and Weigandt [42]. One particular case of this example is the case of toric ideals associated to *Ferrers graphs*. A Ferrers graph F is a bipartite graph such that if (i, j) is an edge of F , then so is (p, q) for $1 \leq p \leq i$ and $1 \leq q \leq j$. In addition, $(1, n)$ and $(d, 1)$ are required to be edges of F . The monomial edge ideals and toric edge ideals of these graphs were studied by Corso and Nagel in [15, 16].

EXAMPLE 5.10. We return to our running Example 1.6, whose corresponding tropical hyperplane arrangement is depicted in Figure 1. In this case the matrix X_A is

$$X_A = \begin{bmatrix} x_{11} & x_{12} & 0 & x_{14} \\ x_{21} & x_{22} & x_{23} & x_{24} \\ x_{31} & 0 & x_{33} & 0 \end{bmatrix}.$$

The lattice ideal L_A is generated by all the distinguished 2-minors of this matrix:

$$\ker \psi = \langle x_{11}x_{22} - x_{12}x_{21}, x_{21}x_{33} - x_{23}x_{31}, x_{11}x_{24} - x_{14}x_{21}, x_{12}x_{24} - x_{14}x_{22} \rangle.$$

Note that one can realize L_A as a determinantal ideal by rearranging the columns of A .

EXAMPLE 5.11. For an example of a lattice ideal that is not a ladder determinantal ideal, consider the tropical hyperplane arrangement corresponding to the tropical matrix

$$A = \begin{bmatrix} 0 & 1 & \infty \\ 0 & \infty & 0 \\ \infty & 0 & 1 \end{bmatrix}.$$

Then the corresponding lattice ideal is $\langle x_{11}x_{23}x_{32} - x_{21}x_{33}x_{12} \rangle$.

In the case that B has no cycles, we easily recover a result of Postnikov [40, Lemma 12.5].

COROLLARY 5.12. *Suppose that $B = B_A$ is a forest. Then Q_B is a simplex.*

Proof. If B is a forest, then it contains no cycles and the generating set of L_A is empty. Therefore, Q_B has no nontrivial triangulations and must itself be a simplex. \square

5.1. KRULL DIMENSION OF TYPE IDEALS. Most of our results involve applying methods from tropical geometry to say something about toric edge rings, this is the focus of Section 6. We can also apply results in the other direction to understand algebraic properties of type ideals.

PROPOSITION 5.13. *Suppose $d, n \geq 2$. Let A be a sufficiently generic $d \times n$ tropical matrix with corresponding bipartite graph $B = B_A$ and tropical hyperplane arrangement $\mathcal{H}(A)$. Then the Krull dimension of the fine cotype ideal satisfies*

$$\dim(I_{\overline{T}(\mathcal{H})}) \leq |E(B)| - 2,$$

with equality if and only if B is bipartite chordal.

Proof. First recall that $I_{\overline{T}(\mathcal{H})}$ is a squarefree monomial ideal in the polynomial ring \tilde{S} with $|E(B)|$ indeterminates (see Notation 4.1 and Definition 4.2). Let $\Delta_{\mathcal{H}}$ denote the associated Stanley-Reisner simplicial complex, so that $\dim(I_{\overline{T}(\mathcal{H})}) = \dim(\Delta_{\mathcal{H}})$. To see that $\dim(I_{\overline{T}(\mathcal{H})}) \leq |E(B)| - 2$, note that since $d, n \geq 2$ we have that a generic element in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ must sit in some region of 2 distinct tropical hyperplanes. Hence $\Delta_{\mathcal{H}}$ has a missing edge, from which it follows that $\dim(\Delta_{\mathcal{H}}) \leq 2$. This proves the first claim.

For the case of equality, recall from Theorem 5.7 that B is bipartite chordal if and only if J_B admits a Gröbner basis consisting of quadratic binomials. The term order underlying such a basis corresponds (via its weight vector) to a regular triangulation of the root polytope Q_B . Let A be the corresponding $d \times n$ tropical matrix (with finite entries corresponding to the height of the lifted vertices of Q_B , and with infinite coordinates otherwise). For this term order \preceq_A we have that the initial ideal $\text{in}_{\preceq_A}(J_B)$ is a quadratic monomial ideal. If we let Γ_A denote the Stanley-Reisner complex of this ideal, it then follows that the minimal nonfaces of Γ_A have cardinality 2 (so that Γ_A is a so-called *flag* complex).

This implies that the facets of the Alexander dual complex Γ_A^\vee has facets of cardinality $|E(B)| - 2$ (recall that the vertex set of $\Delta_{\mathcal{H}}$ is given by $E(B)$). But from Proposition 4.14 we gave that $\Delta_{\mathcal{H}} = \Gamma_A^\vee$, and the claim follows. \square

6. HOMOLOGICAL PROPERTIES OF BIPARTITE TORIC EDGE IDEALS

In this section we use results from above to investigate homological invariants of toric edge ideals of bipartite graphs. We take advantage of the fact that for a bipartite graph B , the lattice ideal of the root polytope Q_B is exactly the toric edge ideal of B . We can then utilize results of Section 4. We first define the invariants that we will be interested in.

DEFINITION 6.1. *Suppose $R = \mathbb{k}[x_1, \dots, x_m]$ and M is a graded finitely generated R -module. The (Castelnuovo-Mumford) regularity and projective dimension of M are given by the values*

$$\begin{aligned} \text{reg}_R(M) &= \max\{j \mid \beta_{i,i+j}(M) \neq 0 \text{ for some } i\}, \\ \text{pdim}_R(M) &= \max\{i \mid \beta_{i,i+j}(M) \neq 0 \text{ for some } j\}, \end{aligned}$$

where $\beta_{i,k}(M)$ are the Betti numbers of M .

Regularity corresponds to the highest degree that appears in any of the matrices in a minimal free resolution, and projective dimension measures the number of steps needed in constructing the resolution. Hence one may think of these invariants as measuring the “complexity” of the module.

DEFINITION 6.2. *A finitely generated graded R -module M admits a q -linear free resolution if its Betti numbers satisfy $\beta_{i,j}(M) = 0$ for each $1 \leq i \leq \text{pdim}(M)$ and for each $j \neq q + i - 1$.*

For the case of $M = R/I$ this means that I is generated by homogeneous polynomials of degree q and $\text{reg}(R/I) = q - 1$. It turns out that regularity and projective dimension of R/I for a squarefree monomial ideal I are related via Alexander duality, and a key tool for us will be the following result of Terai.

THEOREM 6.3. [49, Theorem 0.2] *For $I \subset R$ a squarefree monomial ideal, the projective dimension and regularity satisfy*

$$\text{pdim}(R/I) = \text{reg}(I^\vee).$$

In particular, this implies that R/I has a linear resolution if and only if R/I^\vee is Cohen-Macaulay, a fact first observed by Eagon and Reiner in [20]. We will also need the following recent result of Conca and Varbaro.

THEOREM 6.4 ([14]). *Let I be a homogeneous ideal of a polynomial ring R such that $\text{in}(I)$ is a square-free monomial ideal for some term order. Denote by $h^{ij}(M)$ the dimension of the degree j component of its i th local cohomology module $H_{\mathfrak{m}}^i(M)$ supported on the maximal ideal \mathfrak{m} . Then*

$$h^{ij}(R/I) = h^{ij}(R/\text{in}(I)).$$

In particular, the depth, regularity, and projective dimension of S/I and $S/\text{in}(I)$ coincide.

Recall that an R -module M is *Cohen-Macaulay* if $\text{depth}(M) = \dim(M)$, where $\dim(M)$ is the Krull dimension of M . If B is a bipartite graph, it is known that the toric edge ring $\mathbb{k}[B]$ is Cohen-Macaulay. This follows by combining the fact that toric rings of bipartite graphs are *normal* (by a result of Villarreal [51]), along with a celebrated result of Hochster [29] that says all normal semigroup rings are Cohen-Macaulay. We can easily recover this property by applying our results from previous sections.

PROPOSITION 6.5. *For any connected bipartite graph B the toric edge ideal J_B is Cohen-Macaulay.*

Proof. The root polytope Q_B corresponds to a (sufficiently generic) hyperplane arrangement $\mathcal{H}(A)$ where A is a tropical matrix satisfying $B = B_A$. By Corollary 4.16 the bounded complex $\mathcal{B}(A)$ supports a linear minimal free resolution of the Alexander dual of $\text{in}_A(J_B)$. It follows from Theorem 6.3 that $\tilde{S}/\text{in}_A(J_B)$ is Cohen-Macaulay. Applying Theorem 6.4, one has $\text{depth}(\tilde{S}/J_B) = \text{depth}(\tilde{S}/\text{in}_A(J_B))$, and the result follows. \square

We will see that our knowledge of a minimal resolution of the fine type ideal allows for a novel way to compute regularity of toric edge rings of bipartite graphs. The main tool in our study is the following result, which provides a tropical geometric interpretation for the regularity of a toric edge ring.

THEOREM 6.6. *Suppose A is a sufficiently generic $d \times n$ tropical matrix. Let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$ with bounded complex $\mathcal{B}(A)$. If $B = B_A$ is the bipartite graph corresponding to finite entries of A we have*

$$\text{reg}(\mathbb{k}[B]) = \dim(\mathcal{B}(A)).$$

Proof. By Lemma 5.4, the toric edge ideal J_B is exactly the lattice ideal of the root polytope Q_B . By Corollary 4.15, the bounded complex $\mathcal{B}(A)$ supports a minimal free resolution of \tilde{S}/I , where I is the Alexander dual of the ideal $\text{in}_A(J_B)$. Hence we have that $\text{pdim}(\tilde{S}/I) = \dim(\mathcal{B}(A)) + 1$. From Theorem 6.3 we also have $\text{reg}(\text{in}_A(J_B)) = \text{pdim}(\tilde{S}/I)$. Moreover, $\text{in}_A(J_B)$ is a *squarefree* initial ideal of J_B , so the result follows from Theorem 6.4 above and the fact that $\text{reg}(\mathbb{k}[B]) = \text{reg}(J_B) - 1$. \square

While Theorem 6.6 gives a precise formula for the regularity of J_B , a priori it depends on the matrix A . Proposition 3.7 gives a sufficiently generic A such that the regularity can be characterized in purely in terms of graph properties of B . In what follows recall that $\lambda(B)$ denotes *recession connectivity* of a bipartite graph B , see Definition 3.5.

COROLLARY 6.7. *Let B be a finite simple connected bipartite graph, then $\text{reg}(\mathbb{k}[B]) = \lambda(B) - 1$.*

As the regularity of $\mathbb{k}[B]$ is invariant under any choice of sufficiently generic A satisfying $B = B_A$, we get that the dimension of the bounded complex is also invariant of B .

COROLLARY 6.8. *Let A be a sufficiently generic tropical matrix with corresponding tropical hyperplane arrangement $\mathcal{H}(A)$ and bipartite graph B_A . Then $\dim(\mathcal{B}(A)) = \lambda(B_A) - 1$.*

6.1. WARM-UP: COMPLETE BIPARTITE GRAPHS. We first study the case of toric edge ideals J_B in the case that $B = K_{d,n}$ is a complete bipartite graph. These ideals are very well understood but we will see that the tropical perspective lends further insight into their homological properties. We first note that for $B = K_{d,n}$ the corresponding arrangement consists of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$, each of which has full support. The algebraic properties of the corresponding type and cotype ideals were studied in [18].

PROPOSITION 6.9. *Suppose A is a sufficiently generic tropical matrix with only finite entries, and let $\mathcal{H} = \mathcal{H}(A)$ denote the corresponding arrangement of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbf{1}$. Then the fine cotype ideal $I_{\overline{\mathcal{T}}(\mathcal{H})}$ satisfies*

$$(5) \quad \begin{aligned} \text{reg}(I_{\overline{\mathcal{T}}(\mathcal{H})}) &= (d - 1)(n - 1), \\ \text{pdim}(I_{\overline{\mathcal{T}}(\mathcal{H})}) &= \min\{d, n\} - 1. \end{aligned}$$

Proof. Recall from Proposition 4.5 that a minimal resolution of the fine type cotype ideal $I_{\overline{\mathcal{T}}(\mathcal{H})}$ is supported on the bounded tropical complex $\mathcal{B}(A)$. From Proposition 3.7 and Lemma 3.8 we see that $\dim(\mathcal{B}_A) = \min\{d, n\} - 1$ and hence the formula for projective dimension follows.

For the regularity claim, note that since A is sufficiently generic we have from Corollary 4.16 that the fine cotype ideal $I_{\overline{\mathcal{T}}(\mathcal{H})}$ has a linear resolution. Hence the regularity of $I_{\overline{\mathcal{T}}(\mathcal{H})}$ is equal to the (common) degree of its generators. Note that if v is the apex of any hyperplane of $\mathcal{H}(A)$ then it is not contained in $d - 1$ regions of the remaining $n - 1$ tropical hyperplanes. But v corresponds to a 0-cell of the bounded complex, and hence a generator of $I_{\overline{\mathcal{T}}(\mathcal{H})}$ of degree $(n - 1)(d - 1)$, as desired. \square

Applying Proposition 6.9 and Theorem 6.6, we recover a result from the literature regarding the toric edge rings of complete bipartite graphs (see for instance [25, Lemma 3.10]).

COROLLARY 6.10. For $B = K_{d,n}$ a complete bipartite graph, the toric edge ring $\mathbb{k}[B]$ satisfies

$$\begin{aligned} \text{further } \text{reg}(\mathbb{k}[B]) &= \min\{d, n\} - 1, \\ \text{pdim}(\mathbb{k}[B]) &= (d - 1)(n - 1). \end{aligned}$$

Note that in this case we have $\text{reg}(\mathbb{k}[B]) = \text{mat}(B) - 1$, where B is the *matching number* of B . We discuss this connection in further detail below.

6.2. REGULARITY BOUNDS FOR GENERAL BIPARTITE GRAPHS. We next consider toric edge ideals of arbitrary connected bipartite graphs. Most of our results will be applications of Proposition 6.6, where the regularity of a toric edge ring is described in terms of the dimension of a certain tropical complex. Our first observation is subgraph monotonicity in the regularity of toric edge rings of bipartite graphs.

THEOREM 6.11. Suppose $B \subseteq K_{d,n}$ is a connected bipartite graph and let $B' \subseteq B$ be a connected subgraph. Then we have

$$\text{reg}(\mathbb{k}[B']) \leq \text{reg}(\mathbb{k}[B]).$$

Proof. This follows directly from Lemma 3.9 and Corollary 6.7. \square

In [25] the authors consider a similar question in the context of arbitrary (not necessarily bipartite) graphs. They prove in [25, Theorem 3.7] that if G is any graph with *induced* subgraph G' then we have $\text{reg}(\mathbb{k}[G']) \leq \text{reg}(\mathbb{k}[G])$. We note that the induced property is necessary here: for instance if G' is the graph on 7 vertices consisting of a pair of triangles connected by a path of length 2 (for a total of 8 edges) then we have $\text{reg}(\mathbb{k}[G']) = 3$, whereas $\text{reg}(\mathbb{k}[G]) = 2$ for any graph on 7 vertices with 19 edges. We thank Tina O'Keefe for providing us with this example. In this example we see that $\mathbb{k}[G']$ is not normal. Recall that the toric edge ring of any bipartite graph is normal, and hence a natural question that arises in the following.

QUESTION 6.12. Suppose G is a connected graph with connected subgraph G' such that both $\mathbb{k}[G]$ and $\mathbb{k}[G']$ are normal. Is it true that $\text{reg}(\mathbb{k}[G']) \leq \text{reg}(\mathbb{k}[G])$?

After posting a version of this paper to the arXiv, Akiyoshi Tsuchiya provided a positive answer to our question. Here we include a proof based on his ideas.

Proof. As $\mathbb{k}[G]$ is normal, it follows that the Hilbert function of $\mathbb{k}[G]$ is equal to the Ehrhart function of the edge polytope \mathcal{P}_G , see [27, Lemma 4.22(b)]. Explicitly, the Hilbert series of $\mathbb{k}[G]$ is given by

$$H(\mathbb{k}[G], t) = \frac{h_0^* + h_1^*t + \dots + h_d^*t^d}{(1 - t)^d}$$

where $\dim(\mathcal{P}_G) = d - 1$ and $(h_0^*, h_1^*, \dots, h_d^*) \in \mathbb{Z}_{\geq 0}^{d+1}$ is the h^* -vector of \mathcal{P}_G . Let $(h_0^{*'}, h_1^{*'}, \dots, h_d^{*'})$ be the h^* -vector of $\mathcal{P}_{G'}$, note that $\mathcal{P}_{G'}$ is a subpolytope of \mathcal{P}_G . Therefore by Stanley's monotonicity theorem [47], we have $h_i^{*'} \leq h_i^*$ for all $0 \leq i \leq d$. Since $\mathbb{k}[G]$ is normal, we have from [7, Lemma 2.5] that $\text{reg}(\mathbb{k}[G]) = \max\{s \mid h_s^* \neq 0\}$. The result follows. \square

We next use tropical methods to establish new regularity bounds on toric edge rings.

THEOREM 6.13. Suppose B is a bipartite graph with bipartition $V = \{v_1, \dots, v_n\} \cup \{w_1, \dots, w_d\}$ for $n, d \geq 2$. Let $r = |\{v_i : \deg(v_i) = 1\}|$ and $s = |\{w_j : \deg(w_j) = 1\}|$. Then

$$\text{reg}(\mathbb{k}[B]) \leq \min\{n - r, d - s\} - 1.$$

Proof. Let A be a sufficiently generic tropical matrix satisfying $B = B_A$. Let $\mathcal{H}(A)$ denote the corresponding arrangement of tropical hyperplanes with bounded complex $\mathcal{B}(A)$. By Theorem 6.6, it suffices to show that $\dim(\mathcal{B}(A)) \leq \min\{n - r, d - s\} - 1$. By Proposition 2.15, the bounded complex is a subcomplex of both $\mathcal{H}(A)$ and $\mathcal{H}(A^T)$, where $B^T := B_{A^T}$ is just B where the vertices have swapped sides. The degree 1 vertices of B and B^T correspond to hyperplanes comprised of a single unbounded sector and hence cannot contribute to the dimension of the bounded complex. Hence the dimension of the bounded complex is at most $\min\{n - r, d - s\} - 1$. \square

We note that Theorem 6.13 was established for the case of *chordal* bipartite graphs by Biermann, O’Keefe, and Van Tuyl in [8]. An alternative proof can be given via recession graphs: we observe that if $R(S; B)$ is strongly connected then the subgraph $S \subseteq B$ must contain all leaves of B . Moreover, these leaves cannot form their own connected component, and hence removing them does not decrease recession connectivity. Note that we can repeatedly apply this result to remove certain trees from the graph without decreasing the recession connectivity.

Next we use our tropical tools to recover a regularity result of Herzog and Hibi [26]. Recall that $\text{mat}(B)$ denotes the *matching number* of a graph B .

COROLLARY 6.14. [26, Theorem 1] *If B is a connected bipartite graph then*

$$\text{reg}(\mathbb{k}[B]) \leq \text{mat}(B) - 1.$$

Proof. This follows from Proposition 3.7, Theorem 6.6, and Remark 3.6 \square

We next observe that our bound $\mathbb{k}[B] = \lambda(B) - 1$ can be arbitrarily stronger than the matching number bound. For this consider the following families of examples.

EXAMPLE 6.15. Let B be the path on $2n$ vertices. As B contains no cycles, the edge ring $\mathbb{k}[B]$ is just a polynomial ring and has regularity zero. This can be seen from the recession connectivity, as $R(S; B)$ is strongly connected if and only if $S = B$, and so trivially we have $\lambda(B) = 1$. However, B has a maximal matching of size n , and hence the bound in Corollary 6.7 is as bad as possible.

EXAMPLE 6.16. Let B be the 4-cycles C_1, \dots, C_k with an edge between C_i and C_{i+1} for each $1 \leq i \leq k$, as shown in Figure 10. Taking two disjoint edges in each C_i gives a maximal matching of size $2k$, however the recession connectivity of B is $k + 1$. This is as any subgraph S such that $R(S; B)$ is strongly connected must contain all edges between cycles; Figure 10 gives an example of such a subgraph.

We note that adding a single edge to a graph can have a huge impact on the regularity of the edge ring. Consider the bipartite graph B' obtained by adding a single edge to B between the cycles C_1 and C_k , as depicted in Figure 10. B' has the same matching number as B , but setting S to be a maximal matching now ensures $R(S; B')$ is strongly connected, and the bound in Corollary 6.7 is an equality in this example. To see this note that S no longer needs to contain the edges between cycles, as one can now walk directly from C_k to C_1 without having to backtrack through earlier cycles.

6.3. LINEAR RESOLUTIONS. We next use our tools to investigate toric edge rings of bipartite graphs that have linear resolutions (see Definition 6.2). In [39, Theorem 4.6] Ohsugi and Hibi prove that a toric edge ring $\mathbb{k}[G]$ has a 2-linear resolution if and only if $G = K_{2,n}$, in which case $\mathbb{k}[G]$ is isomorphic to the polynomial ring over a Segre product of $\mathbb{k}[x_1, x_2]$ and $\mathbb{k}[x_1, \dots, x_n]$. In [28] Hibi, Matsuda, and Tsuchiya prove that if $\mathbb{k}[G]$ is the edge ring of a finite connected simple graph with a 3-linear resolution then $\mathbb{k}[G]$ is a hypersurface. Moreover they conjecture that the same is true for any




FIGURE 10. The bipartite graphs B and B' discussed in Example 6.16. Both graphs have matching number $2k$, but very different recession connectivity: $\lambda(B) = k + 1$ while $\lambda(B') = 2k$. Examples of subgraphs that realize the recession connectivity are highlighted in bold.

$\mathbb{k}[G]$ with a q -linear resolution. In [50] Tsuchiya establishes this conjecture for the case of finite connected bipartite graphs, using combinatorial and algebraic methods. Very recently Mori, Ohsugi, and Tsuchiya [36] have established the conjecture in full generality.

EXAMPLE 6.17. Examples of toric edge rings with q -linear resolutions are given by the case of $B = C_{2q}$, cycles of length $2q$. In this case the toric edge ideal has a single generator (of degree q) given by the binomial corresponding to the cycle itself. The corresponding tropical hyperplane arrangement consists of q tropical hyperplanes, each of which has only two finite coordinates (an example of a *graphical tropical hyperplane arrangement* from Section 4.5). In this case the bounded complex $\mathcal{B}(A)$ is a simplex of dimension $q - 1$, and hence by Theorem 6.6 we have $\text{reg}(R/J_{C_{2q}}) = q - 1$.

We are able to use our machinery to provide an alternative proof of these results using the geometry of tropical complexes. Our first observation relates the q -linear resolution property of toric edge rings to the geometry of tropical complexes.

LEMMA 6.18. *Let B be a connected finite simple bipartite graph with no cycles of length $< 2q$, and at least two cycles of length $2q$ where $q \geq 3$. Then $\text{reg}(\mathbb{k}[B]) \geq q$.*

Proof. Let C, C' be the two $2q$ -cycles, the proof will be split into cases depending on the structure of C, C' . In each case, we shall consider a minimal connected subgraph $B' \subseteq B$ such that $C \cup C' \subseteq B'$. We then build a subgraph $A \subseteq B'$ with at least $q + 1$ connected components such that $R(A; B')$ is strongly connected, showing that $\lambda(B') \geq q + 1$ for $q \geq 3$. As $\lambda(B) \geq \lambda(B')$ by Lemma 3.9, combining with Corollary 6.7 gives the desired result that $\text{reg}(\mathbb{k}[B]) \geq q$.

We first distinguish the cases of interest. Let $\widehat{V} = V(C) \cap V(C')$ be the set of shared vertices and $\widehat{E} = E(C) \cap E(C')$ the set of shared edges. We claim we have one of the following three cases:

- (1) $|\widehat{V}| \leq 1, \widehat{E} = \emptyset,$
- (2) $\widehat{V} = \{v, v'\}, \widehat{E} = \emptyset$ where v, v' are distance q apart,
- (3) \widehat{E} is a path of length $|\widehat{V}| - 1 \leq q.$

The proof of this claim comes from the following case analysis. If $|\widehat{V}| \leq 1$, clearly there are no shared edges.

Suppose that $|\widehat{V}| > 1$, and that there exist $v, v' \in \widehat{V}$ that are not connected by edges in \widehat{E} . Let P and P' be the shortest paths between v and v' in C and C' respectively; the union $P \cup P'$ contains at most $2q$ edges and must contain a cycle as $P \neq P'$. As B contains no cycles less than length $2q$, this can only happen if P, P' are disjoint and exactly length q . Furthermore, $\widehat{V} = \{v, v'\}$ as all other vertices of $C \cup C'$ are less than distance q from either v or v' .

Suppose all vertices \widehat{V} are connected by edges \widehat{E} . As \widehat{E} is given by the intersection of cycles, it must be a path. Furthermore, if it is of length $> q$, the edges $C \cup C' \setminus \widehat{E}$ form a cycle of length $\leq 2q$, a contradiction.

We now complete the proof by constructing $A \subseteq B' \subseteq B$ such that $R(A; B')$ is strongly connected and $c(A) \geq q + 1$ for each of the cases. An example for the construction in each case is given in Figure 11.

Case 1: Let $B' \subseteq B$ be the union of C, C' along with a minimal path between them—such a path exists as B is connected. We let A be the subgraph of B' whose edges form a matching in each cycle, along with all edges in the minimal path between the cycles. The recession graph $R(A; B')$ is strongly connected: adding a matching from a cycle to A allows one to move between any vertices in the cycle, and adding the minimal path allows one to move between cycles. Furthermore, A is a forest with precisely $2q - 1$ connected components: q disconnected edges in each cycle, minus one as the path between cycles connects an edge from each cycle. As a result, $\lambda(B') \geq 2q - 1 \geq q + 1$ when $q \geq 2$.

Case 2: Let $B' \subseteq B$ be the union $C \cup C'$ and let A consist of a matching from each cycle. As each cycle is covered by a maximal matching, the recession graph $R(A; B')$ is strongly connected. Furthermore, A consists of $2q - 2$ connected components, coming from the $2q$ edges that are disconnected everywhere, except at v, v' where the two matchings share a vertex. As a result, $\lambda(B') \geq 2q - 2 \geq q + 1$ when $q \geq 3$.

Case 3: Let $B' \subseteq B$ be the union $C \cup C'$. Let A be the union of maximal matchings in C, C' that agree on \widehat{E} , and the matching on \widehat{E} is also maximal. As each cycle is covered by a maximal matching, the recession graph $R(A; B')$ is strongly connected. The number of edges of A is

$$2q - \left\lfloor \frac{|\widehat{E}|}{2} \right\rfloor.$$

The only vertices where multiple edges may meet are the endpoints of \widehat{E} . If $|\widehat{E}|$ is odd, the maximal matching on \widehat{E} covers its endpoints, and so all edges of A are disconnected. If $|\widehat{E}|$ is even, one endpoint of \widehat{E} is not covered by the maximal matching. Hence there are two edges meeting this endpoint, giving one connected component with two edges. This means the number of connected components of A is

$$c(A) = \begin{cases} 2q - \left\lfloor \frac{|\widehat{E}|}{2} \right\rfloor & \widehat{E} \text{ odd} \\ 2q - \left\lfloor \frac{|\widehat{E}|}{2} \right\rfloor - 1 & \widehat{E} \text{ even} \end{cases}$$

For $|\widehat{E}| \leq q$ and $q \geq 3$, this is greater than or equal to $q + 1$. This can be checked for small values of q , after which the $2q$ term becomes dominant. \square

THEOREM 6.19. *Suppose $B \subset K_{d,n}$ is a finite connected bipartite graph. Then we have*

- $\mathbb{k}[B]$ has a 2-linear resolution if and only if B is obtained by appending (possibly no) trees to the vertices of $K_{2,n}$.
- If $q \geq 3$ and $\mathbb{k}[B]$ has a q -linear resolution then $\mathbb{k}[B]$ is a hypersurface.

Proof. For the first part assume that B is a bipartite graph as described. Note that from Proposition 5.6 we have $\mathbb{k}[B] \cong \mathbb{k}[K_{2,n}]$, so we can in fact assume that $B = K_{2,n}$. Now, to obtain a tropical hyperplane arrangement $\mathcal{H}(A)$ satisfying $B = B_A$ we can take a collection of d 0-dimensional hyperplanes (points) on the real line $\mathbb{R}^2/\mathbb{R}\mathbf{1} \cong \mathbb{R}$. Then the bounded complex $\mathcal{B}(A)$ has dimension 1, which implies $\mathbb{k}[B]$ has a 2-linear resolution.




FIGURE 11. The graph B' depicted each of the three cases. The collection of bold edges give the bidirected subgraph A .

Next suppose that $B \subset K_{d,n}$ is a connected bipartite graph such that $\mathbb{k}[B]$ has a 2-linear resolution. Let B' denote the graph obtained from B by removing all cut edges (or, equivalently, all edges not contained in any cycle). Because $\mathbb{k}[B] = \mathbb{k}[B']$, it suffices to show that $B' = K_{2,n}$. Since J_B must be generated by quadratics, we know that B' is chordal bipartite with a nonempty edge set. Notice that B' must also be connected, since otherwise the resolution of $\mathbb{k}[B']$ would be the tensor product of the resolutions of $\mathbb{k}[B_1]$ and $\mathbb{k}[B_2]$ and therefore could not be linear.

Assume, seeking contradiction, that B' has at least 3 vertices in each vertex class. Since B' is a nonempty chordal bipartite graph with no cut edges, it must contain a 4-cycle C . Since B is connected, let e be an edge incident to C . Then e itself is contained in another 4-cycle C' (since again B is chordal bipartite and B' has no cut edges). This leaves us with three cases, since C and C' can intersect in no edges, a single edge, or two edges. If C and C' intersect in a single edge, then there is a 6 cycle on 6 vertices contained in B' , so $\lambda(B')$ is at least 3 (by monotonicity of regularity, Theorem 6.11) and therefore $\text{reg}(\mathbb{k}[B'])$ is at least 2 by Corollary 6.7, giving a contradiction. Similarly, if C and C' share no edges, then there is an 8 cycle in B' on 7 vertices and a similar argument shows that $\lambda(B')$ is at least 3 and the regularity is too high. The only case remaining is if for any edge e incident to C , the only cycles C' containing e share two edges with C . In this case B' is exactly $K_{2,n}$, proving the claim.

For the second part we assume $q \geq 3$ and that $B \subset K_{d,n}$ is a connected bipartite graph such $\mathbb{k}[B]$ has a q -linear resolution. To prove the claim, it suffices to show that if B has no even cycles of length $< 2q$ and has at least two even cycles of length $2q$ then $\text{reg}(\mathbb{k}[B]) \geq q$. This follows from Lemma 6.18. \square

Recall that J_G has a q -linear resolution if and only if a square-free initial ideal $\text{in}(J_G)$ has this property, which is the case exactly when the $(\text{in}(J_G))^\vee$ (in this case corresponding to the underlying fine cotype ideal) is Cohen-Macaulay. Hence we have a characterization of which tropical hyperplane arrangements give rise to Cohen-Macaulay cotype ideals. Note that since type ideals have linear resolutions (see Corollary 4.16) this implies that such ideals are in fact *bi-Cohen-Macaulay*.

7. FURTHER COMMENTS AND QUESTIONS

In this section we address some open questions and directions for future research. As we have seen, the connection between subdivisions of root polytopes (and the associated theory of tropical hyperplane arrangements and generalized permutohedra) and the study of toric edge ideals have lead to new insights in both contexts. We expect that further applications of these ideas can be pursued, and in this section we collect some potential directions.

7.1. GENERALIZED PERMUTOHEDRA. Recall from Definition 2.19 that a bipartite graph B gives rise to a generalized permutohedron P_B obtained as a sum of simplices. In Corollary 4.6 we proved that any regular fine mixed subdivision of P_B supports a minimal cellular resolution of the monomial ideal generated by the lattice points of P_B . However, not all generalized permutohedra arise as sums of simplices; instead, it is known that any generalized permutohedron can be realized as a Minkowski sum and *difference* of simplices (see for instance [5]). It is natural to wonder whether one can extend the algebraic theory we developed here and describe minimal resolutions of the monomial ideal generated by lattice points of generalized permutohedra. A first step in this direction would be to understand our constructions for the case of Minkowski differences. This involves developing a good notion of mixed subdivision in this setting, which is not something we have seen in the literature.

One particularly well-studied class of generalized permutohedra is the class of *matroid* or *matroid basis* polytopes, which are generalized permutohedra whose vertices are $\{0, 1\}$ -vectors. The complex M supports a (non-minimal) cellular resolution of a monomial ideal I_M known as a *matroidal ideal*. Furthermore, one can check that the Betti numbers of the ideal I_M are Möbius invariants of certain deletions of M [Alex Fink, private communication]. A natural question that arises is the following.

QUESTION 7.1. *For a matroid M can one describe a minimal cellular resolution of the matroidal ideal I_M ? What about for certain classes of matroids?*

One of the simplest examples of a Minkowski difference is the case of a *trimmed permutohedron*,

$$P_B^- = P_B - \Delta_{[d]},$$

where $P_B = \sum_k \Delta_{I_k}$ is the generalized permutohedron associated to some bipartite graph $B \subset K_{d,n}$ and where for all k we have $\Delta_{I_k} \subset \mathbb{R}^n$. One can check [40, Theorem 12.9] that for any triangulation $\{\Delta_{T_1}, \dots, \Delta_{T_s}\}$ of the root polytope Q_B , the set of shifted left degree vectors $\{\text{LD}(T_1) - 1, \dots, \text{LD}(T_s) - 1\}$ is exactly the set of lattice points of P_B^- .

QUESTION 7.2. *Does a (subdivision of a) trimmed permutohedron P_B^- support a resolution of any naturally occurring ideal?*

7.2. h -VECTORS. Stanley famously conjectured that the h -vector of a graded Cohen-Macaulay integral domain is log-concave in [46]. It is known that Stanley’s conjecture is not true in general, but it is still open for special cases. For instance, Conca and Herzog conjectured that the h -vector of a ladder determinantal ring is always log concave and Rubey proved this in the special case of 2×2 ladders [43]. As far as the authors are aware, the question is still open for the case of larger minors and for toric edge ideals of bipartite graphs.

In our situation, we know that the following polynomials coincide:

- the h -polynomial of the toric edge ideal J_B of a bipartite graph B .
- the h -polynomial of any *initial* ideal in A J_B with respect to any weight matrix A (such as a tropical matrix).
- the h -polynomial of the Stanley-Reisner complex of any in A J_B , which is a shellable unimodular triangulation of the root polytope Q_B .
- the Ehrhart polynomial of the root polytope Q_B .
- the interior polynomial of either hypergraph of Q_B (see [33]).

Our hope is that our methods can provide some headway in proving Stanley’s conjecture in the case of ladder determinantal ideals, or perhaps even for toric edge ideals of all bipartite graphs.

7.3. POLARIZATIONS. An arrangement $\mathcal{H} = \mathcal{H}(A)$ of n tropical hyperplanes in $\mathbb{R}^d/\mathbb{R}\mathbb{1}$ gives rise to both a fine type ideal $I_{T(\mathcal{H})} \subset \tilde{S} = \mathbb{k}[x_{11}, \dots, x_{dn}]$ as well as a coarse type ideal $I_{\mathbf{t}(\mathcal{H})} \subset S = \mathbb{k}[x_1, \dots, x_d]$. There is a natural map $I_{T(\mathcal{H})} \rightarrow I_{\mathbf{t}(\mathcal{H})}$ given by ‘forgetting the label’ on each hyperplane. This map is a flat deformation and is an example of *polarization* of monomial ideals. For example, in the case of hyperplanes with full support, we obtain a family of polarizations of the ideal $\langle x_1, \dots, x_n \rangle^d$, studied for instance in [4].

On the other hand, an axiomatic approach to triangulations of products of simplices (and more generally of root polytopes) was initiated by Galashin, Nenashev, and Postnikov in [22]. Here the notion of a *trianguloid* is defined as an edge colored graph satisfying simple local axioms, and in [22] it is shown that trianguloids are in bijection with triangulations of root polytopes. One can check that requiring a subset of these axioms gives rise to a polarization of the underlying ideal. The connection between trianguloids and polarizations first appeared in the first author’s thesis [1] and is further explored by the authors in a follow up paper [3].

7.4. GRAPHIC TROPICAL HYPERPLANE ARRANGEMENTS AND G -PARKING FUNCTION IDEALS. In Subsection 4.5.1, we saw a connection between graphic tropical hyperplane arrangements and classical hyperplane arrangements. In particular, the tropical complex associated to such an arrangement can be recovered as the subdivision of \mathbb{R}^{d-1} obtained from a perturbation of the underlying graphic arrangement. As discussed in Section 4.5, the fine type ideals of these arrangements can be recovered as the oriented matroid ideals of [37].

In [19] it is shown that an affine slice of a graphic hyperplane arrangement associated to G defines a cellular resolution of the underlying G -parking function ideal M_G . Here M_G is a monomial ideal depending on a graph G and a choice of root vertex $v \in G$, and whose standard monomial ideals corresponding to G -parking functions. These ideals were introduced in [41] and have connections to power ideals and chip-firing on graphs.

One can see that M_G can be obtained as a certain coarse type ideal associated to a graphic tropical hyperplane. An application of tropical convexity recovers the main result of [19], and also places the theory in a broader context where, for instance, a choice of root vertex is not necessary. In particular, the tropical theory allows one to define parking functions for arbitrary root polytopes. We explore these ideas in [2].

Acknowledgements. We would like to thank Laura Ballard, Michael Joswig, Suho Oh, Tina O’Keefe, Vic Reiner, Raman Sanyal, and Adam Van Tuyl for helpful conversations. We also thank Akiyoshi Tsuchiya for providing us a positive answer to Question 6.12 and Kieran Bhaskara for pointing out an error in earlier version of the statement of Theorem 6.13. We are grateful to an anonymous referee for helpful comments.

REFERENCES

- [1] Ayah Almousa, *Combinatorial Characterizations of Polarizations of Powers of the Graded Maximal Ideal*, Ph.D. thesis, Cornell University, 2021.
- [2] Ayah Almousa, Anton Dochtermann, and Ben Smith, *Tropical types, parking function ideals, and root polytopes*, 2025, work in progress.
- [3] Ayah Almousa, Anton Dochtermann, and Ben Smith, *Tropical types, trianguloids, and polarizations*, 2025, work in progress.
- [4] Ayah Almousa, Gunnar Fløystad, and Henning Lohne, *Polarizations of powers of graded maximal ideals*, J. Pure Appl. Algebra **226** (2022), no. 5, article no. 106924 (33 pages).
- [5] Federico Ardila, Carolina Benedetti, and Jeffrey Doker, *Matroid polytopes and their volumes*, Discrete Comput. Geom. **43** (2010), no. 4, 841–854.

- [6] Laura E Ballard, *Properties of the Toric Rings of a Chordal Bipartite Family of Graphs*, Ph.D. thesis, Syracuse University, 2020.
- [7] Bruno Benedetti and Matteo Varbaro, *On the dual graphs of Cohen-Macaulay algebras*, Int. Math. Res. Not. IMRN **2015** (2015), no. 17, 8085–8115.
- [8] Jennifer Biermann, Augustine O’Keefe, and Adam Van Tuyl, *Bounds on the regularity of toric ideals of graphs*, Adv. in Appl. Math. **85** (2017), 84–102.
- [9] Florian Block and Josephine Yu, *Tropical convexity via cellular resolutions*, J. Algebraic Combin. **24** (2006), no. 1, 103–114.
- [10] Peter Butkovič, *Max-linear systems: theory and algorithms*, Springer Monographs in Mathematics, Springer-Verlag London, Ltd., London, 2010.
- [11] Guy Cohen, Stéphane Gaubert, and Jean-Pierre Quadrat, *Duality and separation theorems in idempotent semimodules*, Linear Algebra Appl. **379** (2004), 395–422.
- [12] Aldo Conca, *Ladder determinantal rings*, J. Pure Appl. Algebra **98** (1995), no. 2, 119–134.
- [13] Aldo Conca, *Gorenstein ladder determinantal rings*, J. London Math. Soc. (2) **54** (1996), no. 3, 453–474.
- [14] Aldo Conca and Matteo Varbaro, *Square-free Gröbner degenerations*, Invent. Math. **221** (2020), no. 3, 713–730.
- [15] Alberto Corso and Uwe Nagel, *Specializations of Ferrers ideals*, J. Algebraic Combin. **28** (2008), no. 3, 425–437.
- [16] Alberto Corso and Uwe Nagel, *Monomial and toric ideals associated to Ferrers graphs*, Trans. Amer. Math. Soc. **361** (2009), no. 3, 1371–1395.
- [17] Mike Develin and Bernd Sturmfels, *Tropical convexity*, Doc. Math. **9** (2004), 1–27.
- [18] Anton Dochtermann, Michael Joswig, and Raman Sanyal, *Tropical types and associated cellular resolutions*, J. Algebra **356** (2012), 304–324.
- [19] Anton Dochtermann and Raman Sanyal, *Laplacian ideals, arrangements, and resolutions*, J. Algebraic Combin. **40** (2014), no. 3, 805–822.
- [20] John A. Eagon and Victor Reiner, *Resolutions of Stanley-Reisner rings and Alexander duality*, J. Pure Appl. Algebra **130** (1998), no. 3, 265–275.
- [21] Alex Fink and Felipe Rincón, *Stiefel tropical linear spaces*, J. Combin. Theory Ser. A **135** (2015), 291–331.
- [22] Pavel Galashin, Gleb Nenashev, and Alexander Postnikov, *Trianguloids and triangulations of root polytopes*, J. Combin. Theory Ser. A **201** (2024), article no. 105802 (28 pages).
- [23] Evgenij Gawrilov and Michael Joswig, *polymake: a framework for analyzing convex polytopes*, in Polytopes—combinatorics and computation (Oberwolfach, 1997), DMV Sem., vol. 29, Birkhäuser, Basel, 2000, pp. 43–73.
- [24] Elisa Gorla, *Mixed ladder determinantal varieties from two-sided ladders*, J. Pure Appl. Algebra **211** (2007), no. 2, 433–444.
- [25] Huy Tài Hà, Selvi Kara Beyarslan, and Augustine O’Keefe, *Algebraic properties of toric rings of graphs*, Comm. Algebra **47** (2019), no. 1, 1–16.
- [26] Jürgen Herzog and Takayuki Hibi, *The Regularity of Edge Rings and Matching Numbers*, Mathematics **8** (2020), no. 1, 103–114.
- [27] Jürgen Herzog, Takayuki Hibi, and Hidefumi Ohsugi, *Binomial ideals*, Graduate Texts in Mathematics, vol. 279, Springer, Cham, 2018.
- [28] Takayuki Hibi, Kazunori Matsuda, and Akiyoshi Tsuchiya, *Edge rings with 3-linear resolutions*, Proc. Amer. Math. Soc. **147** (2019), no. 8, 3225–3232.
- [29] M. Hochster, *Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes*, Ann. of Math. (2) **96** (1972), 318–337.
- [30] Birkett Huber, Jörg Rambau, and Francisco Santos, *The Cayley trick, lifting subdivisions and the Bohne-Dress theorem on zonotopal tilings*, J. Eur. Math. Soc. (JEMS) **2** (2000), no. 2, 179–198.
- [31] Michael Joswig, *Essentials of tropical combinatorics*, Graduate Studies in Mathematics, vol. 219, American Mathematical Society, Providence, RI, 2021.
- [32] Michael Joswig and Georg Loho, *Weighted digraphs and tropical cones*, Linear Algebra Appl. **501** (2016), 304–343.
- [33] Tamás Kálmán and Alexander Postnikov, *Root polytopes, Tutte polynomials, and a duality theorem for bipartite graphs*, Proc. Lond. Math. Soc. (3) **114** (2017), no. 3, 561–588.
- [34] G. L. Litvinov, V. P. Maslov, and G. B. Shpiz, *Idempotent functional analysis. An algebraic approach*, Mat. Zametki **69** (2001), no. 5, 758–797.
- [35] Ezra Miller and Bernd Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics, vol. 227, Springer-Verlag, New York, 2005.

- [36] Kenta Mori, Hidefumi Ohsugi, and Akiyoshi Tsuchiya, *Edge rings with q -linear resolutions*, J. Algebra **593** (2022), 550–567.
- [37] Isabella Novik, Alexander Postnikov, and Bernd Sturmfels, *Syzygies of oriented matroids*, Duke Math. J. **111** (2002), no. 2, 287–317.
- [38] Hidefumi Ohsugi and Takayuki Hibi, *Koszul bipartite graphs*, Adv. in Appl. Math. **22** (1999), no. 1, 25–28.
- [39] Hidefumi Ohsugi and Takayuki Hibi, *Toric ideals generated by quadratic binomials*, J. Algebra **218** (1999), no. 2, 509–527.
- [40] Alexander Postnikov, *Permutohedra, associahedra, and beyond*, Int. Math. Res. Not. IMRN **2009** (2009), no. 6, 1026–1106.
- [41] Alexander Postnikov and Boris Shapiro, *Trees, parking functions, syzygies, and deformations of monomial ideals*, Trans. Amer. Math. Soc. **356** (2004), no. 8, 3109–3142.
- [42] Jenna Rajchgot, Colleen Robichaux, and Anna Weigandt, *Castelnuovo-Mumford regularity of ladder determinantal varieties and patches of Grassmannian Schubert varieties*, J. Algebra **617** (2023), 160–191.
- [43] Martin Rubey, *The h -vector of a ladder determinantal ring cogenerated by 2×2 minors is log-concave*, J. Algebra **292** (2005), no. 2, 303–323.
- [44] Francisco Santos, *Triangulations of oriented matroids*, Mem. Amer. Math. Soc. **156** (2002), no. 741, viii+80.
- [45] Francisco Santos, *The Cayley trick and triangulations of products of simplices*, in Integer points in polyhedra—geometry, number theory, algebra, optimization, Contemp. Math., vol. 374, Amer. Math. Soc., Providence, RI, 2005, pp. 151–177.
- [46] Richard P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, in Graph theory and its applications: East and West (Jinan, 1986), Ann. New York Acad. Sci., vol. 576, New York Acad. Sci., New York, 1989, pp. 500–535.
- [47] Richard P. Stanley, *A monotonicity property of h -vectors and h^* -vectors*, European J. Combin. **14** (1993), no. 3, 251–258.
- [48] Bernd Sturmfels, *Gröbner bases and convex polytopes*, University Lecture Series, vol. 8, American Mathematical Society, Providence, RI, 1996.
- [49] Naoki Terai, *Generalization of Eagon-Reiner theorem and h -vectors of graded rings*, 1997, unpublished preprint.
- [50] Akiyoshi Tsuchiya, *Edge rings of bipartite graphs with linear resolutions*, J. Algebra Appl. **20** (2021), no. 9, article no. 2150163 (8 pages).
- [51] Rafael H. Villarreal, *Rees algebras of edge ideals*, Comm. Algebra **23** (1995), no. 9, 3513–3524.

AYAH ALMOUSA, University of South Carolina, Dept. of Mathematics, 1523 Greene Street, LeConte College Rm 417, Columbia, SC 29208 (USA)
E-mail : aalmousa@sc.edu
Url : <https://sites.google.com/view/ayah-almousa>

ANTON DOCHTERMANN, Texas State University, Dept. of Mathematics, MCS 470, 601 University Drive, San Marcos, TX 78666 (USA)
E-mail : dochtermann@txstate.edu
Url : <https://dochtermann.wp.txstate.edu/>

BEN SMITH, University of Lancaster, School of Mathematical Sciences, Fylde College, Lancaster, LA1 4YF (United Kingdom)
E-mail : b.smith9@lancaster.ac.uk
Url : <https://sites.google.com/view/bsmithmathematics/>